

Problem 19

Use the Binomial Theorem to prove the following identities...

$$\text{a) } \sum_{j=0}^n \binom{n}{j} = 2^n \quad \text{b) } \sum_{k=1}^n (-1)^k \binom{n}{k} = -1$$

Key: $2 = 1 + 1$

$$(1+1)^n = \sum_{j=0}^n \binom{n}{j} 1^{n-j} 1^j$$

Key: $0 = 1 - 1$

$$(1-1)^n = \sum_{k=0}^n \binom{n}{k} 1^{n-k} (-1)^k$$

$$\begin{aligned} &= \sum_{k=0}^n \binom{n}{k} 1^{n-k} (-1)^k \\ &= \sum_{k=0}^n \binom{n}{k} (-1)^k \end{aligned}$$

□

The General Inclusion-Exclusion Principle

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder




Inclusion-Exclusion on 3 sets

$$|A \cup B| = |A| + |B| - |A \cap B|$$
$$|A \cup B \cup C| = |A| + |B| + |C| - |A \cap B| - |A \cap C| - |B \cap C| + |A \cap B \cap C|$$

<https://powcoder.com>

Add WeChat powcoder

Now everything
is counted exactly
once!



Problem 20

How many permutations of the 26 letters of the alphabet do not contain any of the strings "fish", "rat" or "bird"? Use complement.

Assignment Project Exam Help

F : Permutations that contain "fish" $\rightarrow |F| = 23!$

R : Permutations that contain "rat" $\rightarrow |R| = 24!$

B : Permutations that contain "bird" $\rightarrow |B| = 23!$

Goal: Count $|X \setminus (F \cup R \cup B)| = 26! - |F \cup R \cup B|$

$|F| = 23!$ (place "fish" in 24 positions, 23! permutations of other letters in order)

$|F \cap R|$: Place "fish", "rat" and 19 other letters $\Rightarrow 21!$ permutations.

$$|F \cap B| = 0$$

$$|R \cap B| = 0$$

$$|F \cap R \cap B| = 0$$

$$\Rightarrow \text{Answer: } 26! - (23! + 24! + 23! - 21! - 0 - 0 + 0)$$

$$= 26! - 2 \cdot 23! - 24! + 21! \quad \text{Fun simplifying}$$

Letter
in common

Inclusion-Exclusion on n sets

$$\begin{aligned}
 |A_1 \cup A_2 \cup \dots \cup A_n| &= |A_1| + |A_2| + \dots + |A_n| \\
 &\quad - |A_1 \cap A_2| - |A_1 \cap A_3| - \dots - |A_{n-1} \cap A_n| \\
 &\quad + |A_1 \cap A_2 \cap A_3| + |A_1 \cap A_2 \cap A_4| + \dots + |A_{n-2} \cap A_{n-1} \cap A_n| \\
 &\quad - \dots + (-1)^{n+1} |A_1 \cap A_2 \cap \dots \cap A_n|
 \end{aligned}$$

Handwritten notes:
 x is counted: $+k$ times
 $-\binom{k}{2}$ times
 $+\binom{k}{3}$ times
 $+\binom{k}{n}$

Assignment Project Exam Help

<https://powcoder.com>

→ Proof:

We have to make sure that any $x \in A_1 \cup A_2 \cup \dots \cup A_n$ is counted exactly once in the RHS.

Take x arbitrary. x appears in exactly k of the sets A_i .

$\Rightarrow x$ is counted:

$$\begin{aligned}
 \sum_{j=1}^n \binom{k}{j} (-1)^{j+1} &= \sum_{j=1}^k \binom{k}{j} (-1)^{j+1} \quad \text{because } \binom{k}{j} = 0 \text{ if } j > k \\
 &= - \sum_{j=1}^k \binom{k}{j} (-1)^j = -(-1)^k = 1 \quad \text{Prob 19} \quad \square
 \end{aligned}$$

Inclusion-Exclusion on n sets

Special case! When

$$N(k) = |A_{i_1} \cap \dots \cap A_{i_k}|$$

is a number that just depends on k then the formula becomes:

$$\begin{aligned} |A_1 \cup A_2 \cup \dots \cup A_n| &= \sum_{k=1}^n (-1)^{k+1} \sum_{1 \leq i_1 < \dots < i_k \leq n} N(k) \\ &= \sum_{k=1}^n (-1)^{k+1} N(k) \sum_{1 \leq i_1 < \dots < i_k \leq n} 1 \\ &= \sum_{k=1}^n (-1)^{k+1} \binom{n}{k} N(k) \end{aligned}$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Problem 21

(onto)

Let $|A| = m$ and $|B| = n$. Count the number of **surjective** functions $f: A \rightarrow B$.

*Recall: $\forall y \in B, \exists x \in A, f(x) = y$
for all y .
 $m = |A| \geq |B| = n$

(otherwise, answer = 0)

We want to avoid: $\exists y \in B, \forall x \in A, f(x) \neq y$.
(No surjective fnd.)

$X = \text{All functions } f: A \rightarrow B$

for $y \in B$: $F_y = \text{All functions that avoid the value } y \text{ (Range } \subseteq B \setminus \{y\})$.

A function is surjective iff it is in none of the F_y 's.

$$\Leftrightarrow f \in X \setminus \left(\bigcup_{y \in B} F_y \right) = X \setminus (F_1 \cup F_2 \cup \dots \cup F_n)$$

assuming $B = \{1, 2, 3, \dots, n\}$

Let's count $|X \setminus (F_1 \cup F_2 \cup \dots \cup F_n)| = |X| - |F_1 \cup F_2 \cup \dots \cup F_n|$.

Note: $|F_i| = \binom{n}{1} = \binom{n-1}{0} = \dots = \binom{n-1}{n-1}$ $\Rightarrow |F_i| = (n-1)^n$.

(i ≠ j) $|F_i \cap F_j| = (n-2)^n$ $\Rightarrow (n-2)^n$ options of $f(x)$, for each x .

<https://powcoder.com>

By incl-excl: $|F_1 \cup \dots \cup F_n| = \sum_{k=1}^n \binom{n}{k} (-1)^{k+1} N(k)$

Add WeChat powcoder

$N(0) = |X| = n^m$ (n options for each x). $= \sum_{k=1}^n \binom{n}{k} (-1)^{k+1} (n-k)^m$

Answer: $n^m \oplus \sum_{k=1}^n \binom{n}{k} (-1)^{k+1} (n-k)^m = \sum_{k=0}^n \binom{n}{k} (n-k)^m (-1)^k$

* Remark: When we really want to count the complement of a certain union,

Assignment Project Exam Help

$$\begin{aligned} &= N(0) - \sum_{k=1}^n \binom{n}{k} (-1)^{k+1} N(k) \\ &= \sum_{k=0}^n \binom{n}{k} (-1)^k N(k) \end{aligned}$$

Add WeChat powcoder

Shorter formula for the complement of the union!

* Def: $N(0) = |X|$.

Problem 22

The Hatcheck Problem

At a party with a *very large* number of guests, everyone has to leave their hat at the entrance. When they leave the party, everyone is too drunk to remember what hat is theirs, so they just pick one at random. **What is the probability** that no one picks their own hat?

<https://powcoder.com>

Add WeChat powcoder