

Problem 22

The Hatcheck Problem

At a party with a very large number of guests, everyone has to leave their hat at the entrance. When they leave the party, everyone is too drunk to remember what hat is theirs, so they just pick one at random. What is the probability that no one picks their own hat?

Probability = $\frac{n!}{n!}$ # ways that no one picks their own hat
<https://powcoder.com>

Add WeChat powcoder with no fixpoint
(Derangement).

- Def: A derangement of a set with n elements is a permutation of that set that keeps no element in its original position.

Assignment Project Exam Help

- Ex: $X = \{1, 2, 3, 4, 5\}$.

<https://powcoder.com>

$[2, 3, 4, 1, 5]$
 $\begin{matrix} 1^{\text{st}} & 2^{\text{nd}} & 3^{\text{rd}} & 4^{\text{th}} & 5^{\text{th}} \end{matrix}$

That is a derangement.

Add WeChat [powcoder](https://powcoder.com)

$[1, 3, 4, 2, 5]$
 $\begin{matrix} 1 & 2 & 3 & 4 & 5 \end{matrix}$

Not a derangement.

Notation: $!n$: "Subfactorial n "

= Number of derangements of n objects

Assignment Project Exam Help

-Ex: $!0 = 0$

$!1 = 0$

$!2 = 1 \rightarrow [1 \ 2] \quad [2 \ 1]$

$!3 = 2$

\vdots

$!n = ?$

(general formula)

1	2	3	4
2	1	3	4
3	2	1	4
4	3	2	1

<https://powcoder.com>

Add WeChat powcoder

Use inclusion-exclusion:

$X = \{ \text{All permutations of } n \text{ objects} \}$

$A_i = \{ \text{Permutations where } i \text{ is in position } i \}$

$D = \{ \text{Derangements of } n \text{ objects} \}$

<https://powcoder.com>

Ex: $n=5$
 $|A_2 \cap A_4| = ?$

☒ 3 ☐ 4 ☐ 5
3 options 2 opt. 1 opt.
 $= 3! = (5-2)!$

We know that: $|D| = \sum_{k=0}^n (-1)^k \binom{n}{k} N(k)$
Add WeChat powcoder

where $N(k) = |A_{i_1} \cap A_{i_2} \cap \dots \cap A_{i_k}| = (n-k)!$

$N(0) = |X| = n! = (n-0)!$

$n-k$ objects (all except i_1, \dots, i_k)
in $n-k$ positions.

$$!n = |D| = \sum_{k=0}^n (-1)^k \binom{n}{k} (n-k)!$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

⇒ Answer to the problem:

$$\text{Prob.} = \lim_{n \rightarrow \infty} \frac{!n}{n!} = \lim_{n \rightarrow \infty} \frac{n!}{n!} \sum_{k=0}^n \frac{(-1)^k}{k!} = \sum_{k=0}^{\infty} \frac{(-1)^k}{k!} = e^{-1} = \frac{1}{e}$$

Use the MacLaurin series $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$

Generating Functions (Counting with power series)

- Def: Given a sequence $(a_n)_{n=0}^{\infty} = (a_0, a_1, a_2, \dots)$,
the generating function of that sequence

is $A(x) = \sum_{n=0}^{\infty} a_n x^n$.
<https://powcoder.com>

Add WeChat powcoder

- Ex: $a_n = 2^n \Rightarrow A(x) = \sum_{n=0}^{\infty} 2^n x^n = \sum_{n=0}^{\infty} (2x)^n$

$b_n = n \Rightarrow B(x) = \sum_{n=0}^{\infty} n x^n$

-Def: Two formal power series are equal

$$A(x) = B(x)$$

Assignment Project Exam Help

$$\sum_{n=0}^{\infty} a_n x^n$$

$$\sum_{n=0}^{\infty} b_n x^n$$

<https://powcoder.com>

→ Operations: $A(x) + B(x) = \left(\sum_{n=0}^{\infty} a_n x^n \right) + \left(\sum_{n=0}^{\infty} b_n x^n \right)$

Addition

$$= \sum_{n=0}^{\infty} (a_n + b_n) x^n$$

→ Ex: $\left(\sum_{n=0}^{\infty} 2^n x^n \right) + \left(\sum_{n=0}^{\infty} n x^n \right) = \sum_{n=0}^{\infty} (2^n + n) x^n$

Multiplication: (Cauchy's Formula):

Assignment Project Exam Help

<https://powcoder.com>

→ Ex: Add WeChat powcoder

$$\begin{aligned}
 &= (1 + 2x + 4x^2 + 8x^3 + 16x^4 + \dots) (0 + x + 2x^2 + 3x^3 + 4x^4 + \dots) \\
 &= 0 + 1x + (2+2)x^2 + (3+4+4+0)x^3 + (4+6+8+8+0)x^4 + \dots \\
 &= x + 4x^2 + 11x^3 + 26x^4 + \dots
 \end{aligned}$$

* Note: With that structure $(+, \cdot)$, formal power series form a ring, denoted $R[[x]]$.

Assignment Project Exam Help

- Ex 2: $A(x) = \sum_{n=0}^{\infty} x^n \quad (a_n = 1, \forall n)$

<https://powcoder.com>

$$A(x)^2 = A(x) A(x)$$

Add WeChat powcoder

$$= \left(\sum_{h=0}^{\infty} x^h \right) \left(\sum_{n=0}^{\infty} x^n \right) = \sum_{h=0}^{\infty} \left(\sum_{k=0}^h a_k a_{h-k} \right) x^h$$

$$= \sum_{h=0}^{\infty} \left(\sum_{k=0}^h 1 \right) x^h = \sum_{h=0}^{\infty} (h+1) x^h$$

- Ex 3. $A(x) = \sum_{n=0}^{\infty} 1 \cdot x^n$

$B(x) = 1 - x + 0x^2 + 0x^3 + 0x^4 + \dots$

"Finite power series" $b_0 = 1$

$b_1 = -1$

$b_n = 0 : (n > 1)$

Assignment Project Exam Help

$A(x) B(x) = \left(\sum_{n=0}^{\infty} x^n \right) \left(\sum_{n=0}^{\infty} b_n x^n \right)$

<https://powcoder.com>

$= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n a_k b_{n-k} \right) x^n = 1x^0 + 0x + 0x^2 + \dots$

Add WeChat powcoder $\Rightarrow A(x)$ and $B(x)$ are inverses!

$n=0: a_0 b_0 = 1$

$n=1: a_0 b_1 + a_1 b_0 = 1(-1) + (-1)1 = 0$

$n \geq 1: a_0 b_n + a_1 b_{n-1} + \dots + a_{n-1} b_1 + a_n b_0 = 0$

$$\Rightarrow A(x) = (B(x))^{-1}$$

Equality between formal power series

$$\sum_{n=0}^{\infty} x^n = (1-x)^{-1}$$

* Recall

$$\sum_{n=0}^N x^n = \frac{1-x^{N+1}}{1-x}$$

Add WeChat

$$\sum_{n=0}^{\infty} x^n = \lim_{N \rightarrow \infty} \frac{1-x^{N+1}}{1-x}$$

Equality between real numbers $(-1 < x < 1)$

$$\frac{1}{1-x}$$

- Def: Let $F(x) = \sum_{n=0}^{\infty} a_n x^n$ be a formal

power series, and $f: (a, b) \rightarrow \mathbb{R}$ ($a < 0 < b$).

If there is a number $R > 0$ such that for

all $r \in (-R, R)$ the numerical series $F(x)|_{x=r} = F(r)$

converges to the value $f(r)$, then we say that

$F(x)$ represents the function $f(x)$ and

write

$$\underline{F(x)} \approx \underline{f(x)}$$

we will actually write $=$ later...

Note that \approx is compatible with $+$ and \cdot .

$F(x) + G(x) \approx f(x) + g(x)$
Assignment Project Exam Help

$$F(x) G(x) \approx f(x) g(x).$$

\Rightarrow <https://powcoder.com> is the same. We can go from one "world" to the other.

Add WeChat powcoder

Formal
Power
Series



Real
Functions.

We will then allow ourselves to write $=$ instead of \approx .

* Recall: Taylor's Theorem.

If $f(x) = \sum_{h=0}^{\infty} a_n x^n$ then f is infinitely

differentiable and,

<https://powcoder.com>

$$a_n = \frac{f^{(n)}(0)}{n!}$$

Power series representations of a function are unique!

Add WeChat powcoder

Moreover, for the "common calculus functions"
(powers, exp, log, trig, ...) $\sum_{h=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$ does represent $f(x)$.

=> Some power series we may use ...

$$\frac{1 - x^{N+1}}{1 - x} = 1 + x + x^2 + \dots + x^N$$

$$\frac{1}{1 - x} = \sum_{n=0}^{\infty} x^n$$

$$\binom{1/2}{2} = \frac{\frac{1}{2}(\frac{1}{2}-1)}{2!} = \frac{\frac{1}{2}(-\frac{1}{2})}{2} = -\frac{1}{8}$$

Assignment Project Exam Help

$$\frac{1}{1 - ax} = \sum_{n=0}^{\infty} a^n x^n$$

$$\frac{1}{1 - x^r} = \sum_{n=0}^{\infty} x^{nr}$$

*
Binomial
Theorem
(Extends)

$$(1 + x)^r = \sum_{n=0}^{\infty} \binom{r}{n} x^n \text{ where } \binom{m}{k} = \frac{m(m-1)\dots(m-k+1)}{k!}$$

Works for
any real
number!

$$\frac{1}{(1 - x)^r} = \sum_{n=0}^{\infty} \binom{n+r-1}{n} x^n$$

$$\sqrt{1+x} = \sum_{n=0}^{\infty} \binom{2n}{n} \frac{(-1)^{n+1}}{4^n(2n-1)} x^n$$

Add WeChat powcoder

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

$$\cosh(x) = \frac{e^x + e^{-x}}{2} = \sum_{n=0}^{\infty} \frac{x^{2n}}{(2n)!}$$

$$\sinh(x) = \frac{e^x - e^{-x}}{2} = \sum_{n=0}^{\infty} \frac{x^{2n+1}}{(2n+1)!}$$