

# Assignment Project Exam Help

## Predictive Analytics

Week 8: Linear Methods for Regression II

<https://powcoder.com>

Semester 2, 2018

Discipline of Business Analytics, The University of Sydney Business School

## Add WeChat powcoder

## Week 8: Linear Methods for Regression II

1. Introduction

2. Principal components regression

3. Partial least squares (optional)

4. Illustration and discussion

5. Considerations in high dimensions

6. Robust regression

Reading: Chapters 6.3 and 6.4 of ISL.

Exercise questions: Chapter 6.8 of ISL, Q5 and Q6.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

# Assignment Project Exam Help

**Introduction**  
<https://powcoder.com>

Add WeChat powcoder

## Dimension reduction methods (key concept)

# Assignment Project Exam Help

**Dimension reduction methods** consist of building  $M < p$  transformed variables which are linear combinations (projections) of the predictors. We then fit a linear regression of the response on the new variables.

<https://powcoder.com>  
Add WeChat powcoder

# Assignment Project Exam Help

Given the original predictors  $x_1, x_2, \dots, x_p$ , we let  $z_1, z_2, \dots, z_M$  represent  $M < p$  linear combinations of the original predictors, that is

<https://powcoder.com>

$$z_m = \sum_{j=1}^p \phi_{jm} x_j,$$

for some constants  $\phi_{1m}, \phi_{2m}, \dots, \phi_{pm}, m = 1, \dots, M$  to be determined.

Add WeChat powcoder

# Assignment Project Exam Help

We consider a linear regression model for the transformed predictors

$$Y_i = \theta_0 + \sum_{m=1}^M \theta_m z_{im} + \epsilon_i$$

<https://powcoder.com>

which we fit by OLS.

We therefore estimate only  $M+1 < p+1$  regression parameters, reducing variance compared to OLS.

## Dimension reduction methods

The model for the transformed predictors implies a model for the original predictors:

$$\sum_{m=1}^M \theta_m z_{im} = \sum_{m=1}^M \theta_m \left( \sum_{j=1}^p \phi_{jm} x_{ij} \right) = \sum_{j=1}^p \beta_j x_{ij},$$

where

$$\beta_j = \sum_{m=1}^M \theta_m \phi_{jm}.$$

Dimension reduction is therefore a constraint on the original linear regression model. The cost of imposing this restriction is bias.

# Assignment Project Exam Help

- The reduction in variance compared to OLS can be substantial when  $M \ll p$ .

- <https://powcoder.com>  
If  $M = p$  and  $Z_m$  are linearly independent, no dimension reduction occurs and dimension reduction methods are equivalent to OLS on original  $p$  predictors.

Add WeChat powcoder

- Dimension reduction methods can be useful when  $p > N$ .



# Assignment Project Exam Help

Principal components regression  
<https://powcoder.com>

Add WeChat powcoder

## Principal components analysis (key concept)

# Assignment Project Exam Help

**Principal Component Analysis (PCA)** is a popular way of deriving a set of low dimensional set of features from a large dimensional set of variables. In our setting, we want to use PCA to reduce the dimension of the  $N \times p$  design matrix  $\mathbf{X}$ .

In our discussion below, we assume that we first center and standardise all the predictors.

## Principal components (key concept)

# Assignment Project Exam Help

We define the first **principal component** of  $\mathbf{X}$  as the linear combination

$$z_1 = \phi_{11}x_1 + \phi_{21}x_2 + \dots + \phi_{p1}x_p,$$

such that  $z_1$  has largest sample variance among all linear combinations whose coefficients satisfy  $\|\phi_1\|_2^2 = \sum_{j=1}^p \phi_{j1}^2 = 1$ .

## Principal components (key concept)

# Assignment Project Exam Help

The  $m$ -th principal component of  $\mathbf{X}$  is the linear combination

$$z_m = \sum_{j=1}^p \phi_{jm} \mathbf{x}_j,$$

<https://powcoder.com>

such  $z_m$  has largest sample variance among all linear combinations that are orthogonal to  $z_1, \dots, z_{m-1}$  and satisfy  $\|\phi_m\|_2 = 1$ .

Add WeChat powcoder


## Principal components (key concept)

# Assignment Project Exam Help

The first  $m$  principal components of the design matrix  $\mathbf{X}$  provide the best  $m$ -dimensional linear approximation to it, in the sense of capturing variation in the predictor data.

<https://powcoder.com>  
Add WeChat powcoder


## Principal components analysis



The two axes represent predictors. The green line indicates the first principal component and the blue dashed line shows the second principal component.



## Principal components analysis

# Assignment Project Exam Help



## Principal components analysis

Assignment Project Exam Help




<https://powcoder.com>

Add WeChat powcoder



## Principal components analysis

Assignment Project Exam Help



# Assignment Project Exam Help

The **principal components regression** (PCR) method consists of running a regression of  $Y$  on the first  $m$  principal components of  $X$ . The PCR method implicitly assumes that directions of highest variance in  $X$  are the ones most associated with the response.

<https://powcoder.com>  
Add WeChat powcoder

## Principal components regression

---

**Algorithm 1** Principal components regression

---

- 1: Center and standardise the predictors.
  - 2: Use PCA to obtain  $z_1, \dots, z_p$ , the  $p$  principal components of the design matrix  $X$ .
  - 3: **for**  $m = 1, \dots, p$  **do**
  - 4:   Regress the response  $y$  on  $z_1, \dots, z_m$  (the first  $m$  principal components) by OLS and call it  $\mathcal{M}_m$ .
  - 5: **end for**
  - 6: Select the best model out of  $\mathcal{M}_1, \dots, \mathcal{M}_p$  by cross-validation.
-

## Principal Component Regression

- PCR can lead to substantial variance reduction compared to OLS when a small number of components account for a large part of the variation in the predictor data.

- Additional principal components leads to smaller bias, but larger variance.

- In PCR, the number of principal components is typically chosen by cross-validation.

- Try to sketch learning curve for PCR (Train & Validation (Test) MSE vs Number of Components).

- PCR does not perform variable selection.

## Comparison with ridge regression

- There is a close connection between the ridge regression and PCR methods.

Assignment Project Exam Help

- Ridge regression shrinks the coefficients of all principal components, with least shrinkage for the first component progressively smaller shrinkage factors for subsequent components.

<https://powcoder.com>

- PCR leaves the components with largest variance alone and discards the ones with smallest variance.

Add WeChat powcoder


- We can therefore think of ridge regression as a continuous version of PCR. Ridge may be preferred in most cases as it shrinks smoothly.

## Comparison with ridge regression

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



# Assignment Project Exam Help

Partial least squares (optional)  
<https://powcoder.com>

Add WeChat powcoder

## Partial Least Squares

# Assignment Project Exam Help

The partial least squares method (PLS) tries to identify the best linear combinations of predictors in a *supervised* way, in a sense that it takes into account the information in  $y$  to construct the new features. When constructing  $z_{1h}$ , PLS weights the predictors by the strength of their univariate effect on  $y$ .

That contrasts with PCA, which identifies principal directions in an *unsupervised* way.



## Partial Least Squares

---

### Algorithm 2 Partial Least Squares (Initialisation)

---

- 1: Center and standardise the predictors
- 2: Run  $p$  simple linear regressions of  $\mathbf{y}$  on each predictor  $\mathbf{x}_j$  and denote the associated coefficients as  $\phi_{1j}$ .
- 3: Compute the first direction  $\mathbf{z}_1 = \sum_{j=1}^p \phi_{1j} \mathbf{x}_j$ .
- 4: Run a SLR regression of  $\mathbf{y}$  on  $\mathbf{z}_1$  and let the coefficient be  $\hat{\theta}_1$ .  
Call this model  $\mathcal{M}_1$ .
- 5: Orthogonalise each predictor with respect to  $\mathbf{z}_1$ :  $\mathbf{x}_j^{(1)} = \mathbf{x}_j - \mathbf{x}_j \left[ (\mathbf{x}_j^T \mathbf{z}_1) / (\mathbf{x}_j^T \mathbf{x}_j) \right]$ . These are the residuals of a SLR of  $\mathbf{x}_j$  on  $\mathbf{z}_1$ . (continues on the next slide)

## Partial Least Squares

---

### Algorithm Partial Least Squares (continued)

---

1: **for**  $m = 2, \dots, p$  **do**

2: Run  $p$  simple linear regressions of  $y$  on each  $x_j^{(m-1)}$  and denote the associated coefficients as  $\phi_{mj}$ .

3: Compute the new direction  $z_m = \sum_{j=1}^p \phi_{mj} x_j$ .

4: Run a SLR regression of  $y$  on  $z_m$  and let the coefficient be  $\theta_m$ . Call the linear regression model with response  $y$ , inputs  $z_1, \dots, z_m$ , and estimated coefficients  $\hat{\theta}_1, \dots, \hat{\theta}_m$  model  $\mathcal{M}_m$ .

5: Orthogonalise each  $x_j^{(n-1)}$  with respect to  $z_m$ :  $x_j^{(m)} = x_j^{(m-1)} - x_j \left[ (x_j^T z_1) / (x_j^T x_j) \right]$ .

6: **end for**

7: Select the best model out of  $\mathcal{M}_1, \dots, \mathcal{M}_p$  by cross-validation.

---

## PCR and PLS: discussion

- While PCR seeks directions with high variance, PLS seeks directions with high variance and correlation with response.

- The variance aspect tends to dominate, such that PLS behaves similarly to PCR and ridge regression.

- Using  $y$  reduces bias but potentially increases variance. PLS shrinks low variance directions, but can actually inflate high variance ones.


- In practice, PLS often does no better or slightly worse than PCR and ridge regression.

# Assignment Project Exam Help

~~Illustration and discussion~~  
<https://powcoder.com>

Add WeChat powcoder

## Illustration: predicting the equity premium



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Since the cross-validation performance is nearly identical with 2 and 3 components, we select  $M = 2$  for the results below.

Equity premium prediction results

	Train $R^2$	Test $R^2$
OLS	0.108	0.014
PCR	0.039	0.048
PLS	0.085	0.036


Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

For this example, PCR has the best test performance among all linear methods that we have discussed.

## Comparison of shrinkage and selection methods



# Assignment Project Exam Help

~~Considerations in high dimensions~~  
<https://powcoder.com>

Add WeChat powcoder



# Assignment Project Exam Help

A **high-dimensional regime** occurs when the number of predictors is larger than the number of observations ( $p > N$ ). Similar issues occur when  $p \approx N$ .

<https://powcoder.com>

We cannot perform least squares in this setting, recall Lecture 2. If  $p = N$ , the training  $R^2$  is always one. OLS is too flexible when  $p > N$  and will overfit the data when  $p \approx N$ .

Add WeChat powcoder

## Example

# Assignment Project Exam Help

**Text analytics.** In type of analysis, the predictors are often a large number of binary variables indicating the presence of words in a document, search history, etc. This is called a **bag of words** model. With thousands of possible words, the the number of predictors is very large in this type of analysis.

<https://powcoder.com>

We can further extend the feature space to include **n-grams**, recording the appearance of words together in a sequence.

## Considerations in high dimensions

- We can apply variable selection, shrinkage, and dimension reduction methods with carefully tuned hyper-parameters to high-dimensional settings.

- However, even these methods are subject to marked deterioration in performance as the number of irrelevant or very weak predictors increases relative to  $N$ .

- Therefore, we cannot blindly rely on standard methods in high dimensional regimes. We need to carefully consider appropriate dimension reduction and penalisation schemes, preferably based on understanding of the substantive problem.

- The next slide shows an example.

## Supervised Principal Components

---

### Algorithm 3 Supervised Principal Components

---

- 1: Center and standardise the predictors.
  - 2: Run  $p$  separate simple linear regressions of  $y$  on each individual predictor and record the estimated coefficients.
  - 3: **for**  $\theta$  in  $0 \leq \theta_1 < \dots < \theta_K$  **do**
  - 4:   Form a reduced design matrix  $X_\theta$  consisting only of predictors whose SLR coefficient is higher than  $\theta$  in absolute value.
  - 5:   Use PCA to obtain  $z_1, \dots, z_m$ , the first  $m$  principal components of  $X_\theta$ .
  - 6:   Use these principal components to predict the response.
  - 7: **end for**
  - 8: Select  $\theta$  and  $m$  by cross-validation.
-

# Assignment Project Exam Help

**Robust regression**  
<https://powcoder.com>

Add WeChat powcoder

# Assignment Project Exam Help

All the linear regression methods that we have seen so far were based on the squared error loss function, which is equivalent to assuming a Gaussian likelihood for the data.

<https://powcoder.com>

However, estimation based on the squared error loss can result in poor fit when there are **outliers**. This is because the squared error penalises deviations quadratically, so that points with larger residuals have more effect on the estimation than points with low residuals (near the regression line).

Add WeChat powcoder

# Assignment Project Exam Help

One way to achieve robustness to outliers is to replace the squared error losses with other losses that are less influenced by unusual observations.

<https://powcoder.com>

Alternatively (and equivalently in some cases), we replace the Gaussian likelihood with that of a distribution with heavy tails. Such a distribution will assign higher likelihood to outliers, without having to adjust the regression fit to account for them.

Add WeChat powcoder

## Least absolute deviation

The **least absolute deviation** (LAD) estimator is

$$\beta_{\text{lad}} = \underset{\beta}{\operatorname{argmin}} \sum_{i=1}^n \left| y_i - \beta_0 - \sum_{j=1}^p \beta_j x_{ij} \right|$$

LAD estimation is equivalent to ML based on the Laplace distribution.

In the special case when we formulate the minimisation problem

$$\hat{m} = \underset{m}{\operatorname{argmin}} \sum_{i=1}^n |Y_i - m|$$

the LAD estimator  $\hat{m}$  is the sample median of the response.



## Huber loss

A popular method for robust regression is the **Huber loss**:

Assignment Project Exam Help

$$\hat{\beta}_{\text{huber}} = \underset{\beta}{\operatorname{argmin}} \left\{ \sum_{i=1}^N L_{\delta} \left( y_i - \beta_0 - \sum_{j=1}^p \beta_j x_{ij} \right) \right\}$$


<https://powcoder.com>

$$L_{\delta}(e) = \begin{cases} e^2 & \text{if } |e| \leq \delta \\ 2\delta|e| - \delta^2 & \text{if } |e| \geq \delta \end{cases}$$

Add WeChat powcoder

The Huber loss combines the good properties of squared and absolute errors.

## Loss functions



# Assignment Project Exam Help

- What are dimension reduction methods for regression?
- What is principal components analysis (PCR)?
- What is the relationship between PCR and ridge regression?
- What is the high-dimensional regime?
- Explain the purpose of robust regression.

<https://powcoder.com>

Add WeChat powcoder

## Technical appendix (Optional)

A column vector  $v$  is an **eigenvector** of a square matrix  $A$  if it satisfies the equation

$$Av = \lambda v,$$

where  $\lambda$  is a scalar known as the **eigenvalue** associated with  $v$ .

The eigenvectors of  $A$  do not change direction when multiplied by  $A$ .

A scalar  $\lambda$  is an eigenvalue of  $A$  if  $(A - \lambda I)$  is singular

$$\det(A - \lambda I) = 0.$$

## Principal components analysis

The **eigendecomposition** of a diagonalisable  $p \times p$  symmetric real square matrix  $\mathbf{A}$  has the form

$$\mathbf{A} = \mathbf{V} \mathbf{\Lambda} \mathbf{V}^T,$$

where  $\mathbf{\Lambda}$  is a  $p \times p$  diagonal matrix whose diagonal elements  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p \geq 0$  are the eigenvalues of  $\mathbf{A}$  and  $\mathbf{V}$  is a  $p \times p$  orthogonal matrix whose columns  $\mathbf{v}_j$  are the eigenvectors of  $\mathbf{A}$ .

If one or more eigenvalues  $\lambda_j$  are zero then  $\mathbf{A}$  is singular (non-invertible).

# Assignment Project Exam Help

A orthogonal matrix  $V$  is a square matrix whose columns and rows are orthonormal, i.e.,

<https://powcoder.com>

$$VV^T = V^TV = I,$$

such that  $V^{-1} = V^T$ .

Add WeChat powcoder

## Principal components analysis

# Assignment Project Exam Help

Now, let  $\mathbf{X}$  be the  $N \times p$  matrix of centered predictors. The sample variance-covariance matrix of  $\mathbf{X}$  is

$$\mathbf{S} = (\mathbf{X}^T \mathbf{X}) / N$$

<https://powcoder.com>

where  $\mathbf{X}^T \mathbf{X}$  has an eigendecomposition denoted as  $\mathbf{V} \mathbf{\Lambda} \mathbf{V}^T$ . The eigenvalues of  $\mathbf{X}^T \mathbf{X}$  are all positive provided that there is no perfect multicollinearity. Eigenvalues near zero indicate the presence of multicollinearity.

Add WeChat powcoder

## Principal components analysis

The first principal component of  $\mathbf{X}$  is

Assignment Project Exam Help

The sample variance of the first principal component is

$$s_{z_1}^2 = \frac{\mathbf{v}_1^T \mathbf{X}^T \mathbf{X} \mathbf{v}_1}{N} = \frac{\mathbf{v}_1 \mathbf{V} \mathbf{\Lambda} \mathbf{V}^T \mathbf{v}_1}{N} = \frac{\lambda_1}{N},$$

where  $\mathbf{v}_1^T \mathbf{v}_1 = 1$  and  $\mathbf{v}_1^T \mathbf{v}_j = 0$  ( $j \neq 1$ ) since  $\mathbf{V}$  is an orthogonal matrix. The first principal component is therefore the linear combination of the columns of  $\mathbf{X}$  that has the largest variance among all possible normalised linear combinations.



## Principal components analysis

The principal components of  $\mathbf{X}$  are

Assignment Project Exam Help

$$\mathbf{z}_m = \mathbf{X}\mathbf{v}_m.$$

for  $m = 1, \dots, p$ , with decreasing sample variance

<https://powcoder.com>

$$s^2_{\mathbf{z}_m} = \frac{\lambda_m}{N},$$

since  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p > 0$ . Since the eigenvectors are orthogonal, the principal components have sample correlation zero.

Add WeChat powcoder

The principal component  $\mathbf{z}_m$  is the direction of largest variance that is orthogonal to  $\mathbf{z}_1, \dots, \mathbf{z}_{m-1}$ .