

CS 593

Khasha Dehnad

Assignment Project Exam Help

Simple + Multiple Linear Regression

<https://powcoder.com>

Add WeChat powcoder

Class restarts

at 7:45

Simple Regression

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Introduction to Regression Analysis

- Regression analysis is used to:
 - Predict the value of a dependent variable based on the value of at least one independent variable
 - Explain the impact of changes in an independent variable on the dependent variable

Dependent variable: the variable we wish to explain
(also called the **endogenous variable**)

Independent variable: the variable used to explain
the dependent variable
(also called the **exogenous variable**)

Aims

- Describe the relationship between an independent variable X , and a continuous dependent variable Y as a straight line in R^2
 - Two Cases:
 - Fixed X : values of X are preselected by investigator
 - Variable X : a random sample of (X,Y) pairs
- Draw inferences regarding the relationship
- Predict the value of Y for a given X

Simple Linear Regression Model

The population regression model:

Assignment Project Exam Help
<https://powcoder.com>
Add WeChat powcoder

Dependent Variable

Population Y intercept

Population Slope Coefficient

Independent Variable

Random Error term

$$y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$$

Linear component

Random Error component

The diagram illustrates the Simple Linear Regression Model equation: $y_i = \beta_0 + \beta_1 x_i + \varepsilon_i$. The equation is presented in a light orange box. Above the box, several labels identify the components: 'Dependent Variable' points to y_i ; 'Population Y intercept' points to β_0 ; 'Population Slope Coefficient' points to β_1 ; 'Independent Variable' points to x_i ; and 'Random Error term' points to ε_i . Below the box, two purple brackets group the terms: the first bracket under $\beta_0 + \beta_1 x_i$ is labeled 'Linear component', and the second bracket under ε_i is labeled 'Random Error component'. The entire diagram is overlaid with red text: 'Assignment Project Exam Help', the URL 'https://powcoder.com', and 'Add WeChat powcoder'.

Linear Regression Assumptions

- The true relationship form is linear (Y is a linear function of X, plus random error)
- The error terms ε are independent of the x values
- The error terms are random variables with mean 0 and constant variance, σ^2

(the uniform variance property is called homoscedasticity)

$$E[\varepsilon_i] = 0 \quad \text{and} \quad E[\varepsilon_i^2] = \sigma^2 \quad \text{for } (i = 1, \dots, n)$$

- The random error terms, ε_i , are not correlated with one another, so that

$$E[\varepsilon_i \varepsilon_j] = 0 \quad \text{for all } i \neq j$$

Graphically (p 85)

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder





Figure 6.2: Simple Linear Regression Model for Fixed X 's

Simple Linear Regression Model

(continued)



α and β (p 86)




Figure 6.3: Theoretical Regression Line Illustrating α and β

Simple Linear Regression Equation

The simple linear regression equation provides an **estimate** of the population regression line

Assignment Project Exam Help
<https://powcoder.com>
Add WeChat powcoder

Estimated (or predicted) y value for observation i

Estimate of the regression intercept

Estimate of the regression slope

Value of x for observation i

$$\hat{y}_i = b_0 + b_1 x_i$$

The individual random error terms e_i have a mean of zero

$$e_i = (y_i - \hat{y}_i) = y_i - (b_0 + b_1 x_i)$$

Least Squares Coefficient Estimators

- b_0 and b_1 are obtained by finding the values of b_0 and b_1 that minimize the sum of the squared residuals (errors), SSE:

<https://powcoder.com>
Add WeChat: powcoder

$$\begin{aligned}\min \text{SSE} &= \min \sum_{i=1}^n e_i^2 \\ &= \min \sum (y_i - \hat{y}_i)^2 \\ &= \min \sum [y_i - (b_0 + b_1 x_i)]^2\end{aligned}$$

Differential calculus is used to obtain the coefficient estimators b_0 and b_1 that minimize SSE

Prediction

- The regression equation can be used to predict a value for y given a particular x

Assignment Project Exam Help

<https://powcoder.com>

- For a specified value, x_{n+1} , the predicted value is

$$\hat{y}_{n+1} = b_0 + b_1 x_{n+1}$$

Least Squares Coefficient Estimators

(continued)

- The slope coefficient estimator is

$$b_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2} = \frac{\text{Cov}(x, y)}{s_x^2} = r \frac{s_y}{s_x}$$


- And the constant or y-intercept is

$$b_0 = \bar{y} - b_1 \bar{x}$$

- The regression line always goes through the mean \bar{x} , \bar{y}

Analysis of Variance

(continued)



Explanatory Power of a Linear Regression Equation

- Total variation is made up of two parts:

$$SST = SSR + SSE$$

Total Sum of
Squares

Regression Sum of
Squares

Error (residual)
Sum of Squares

$$SST = \sum (y_i - \bar{y})^2$$

$$SSR = \sum (\hat{y}_i - \bar{y})^2$$

$$SSE = \sum (y_i - \hat{y}_i)^2$$

where:

\bar{y} = Average value of the dependent variable

y_i = Observed values of the dependent variable

\hat{y}_i = Predicted value of y for the given x_i value

Proof

$$\begin{aligned}\sum_{i=1}^n (Y_i - \bar{Y})^2 &= \sum_{i=1}^n (\hat{Y}_i - \bar{Y} + Y_i - \hat{Y}_i)^2 \\ &= \sum_{i=1}^n \{(\hat{Y}_i - \bar{Y})^2 + (Y_i - \hat{Y}_i)^2 + 2(\hat{Y}_i - \bar{Y})(Y_i - \hat{Y}_i)\}\end{aligned}$$

$$= SSR + SSE + 2 \sum_{i=1}^n (\hat{Y}_i - \bar{Y})(Y_i - \hat{Y}_i)$$

$$= SSR + SSE + 2 \sum_{i=1}^n (\hat{Y}_i - \bar{Y})e_i$$

$$= SSR + SSE + 2 \sum_{i=1}^n (b_0 + b_1 X_i - \bar{Y})e_i$$

$$= SSR + SSE + 2b_0 \sum_{i=1}^n e_i + 2b_1 \sum_{i=1}^n X_i e_i - 2\bar{Y} \sum_{i=1}^n e_i$$

$$= SSR + SSE$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Hypothesis Test for Population Slope Using the F Distribution

- F Test statistic:

$$F = \frac{MSR}{MSE}$$

where

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

$$MSR = \frac{SSR}{k}$$
$$MSE = \frac{SSE}{n - k - 1}$$

where F follows an F distribution with k numerator and $(n - k - 1)$ denominator degrees of freedom

(k = the number of independent variables in the regression model)

Computer Analysis

Results:

- estimates of slope (β_1) and intercept (β_0), using least squares
- residual mean square = estimate of variance (S^2)
- test if $\beta = \beta_0$
 - Usually, test $\beta = 0$, i.e. X has no effect on Y

Hypothesis Test for Population Slope Using the F Distribution

(continued)

- An alternate test for the hypothesis that the slope is zero:

Assignment Project Exam Help

$$H_0: \beta_1 = 0$$

$$H_1: \beta_1 \neq 0$$

<https://powcoder.com>

- Use the F statistic

Add WeChat powcoder

$$F = \frac{MSR}{MSE} = \frac{SSR}{s_e^2}$$

- The decision rule is

$$\text{reject } H_0 \text{ if } F \geq F_{1,n-2,\alpha}$$

Steps in Simple Regression


1. State the research hypothesis.
2. State the null hypothesis
3. Gather the data
4. Assess each variable separately first (obtain measures of central tendency and dispersion; frequency distributions; graphs); is the variable normally distributed?
5. Calculate the regression equation from the data
6. Calculate and examine appropriate measures of association and tests of statistical significance for each coefficient and for the equation as a whole
7. Accept or reject the null hypothesis
8. Reject or accept the research hypothesis
9. Explain the practical implications of the findings

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Effect of Outliers (p 102)



Leverage

Assignment Project Exam Help

$$h_i = \frac{1}{n} + \frac{(x_i - \bar{x})^2}{\sum (x_i - \bar{x})^2}$$

<https://powcoder.com>

Add WeChat powcoder

Influence Measures

- Cook's distance: “distance” between B with and without the i^{th} observation
- DFFITS: “distance” between \hat{Y} with and without the i^{th} observation

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Cook's Distance

Assignment Project Exam Help

$$D_i = \frac{(y_i - \hat{y}_i)^2}{(m-1)s^2} \frac{h_i}{(1-h_i)^2}$$

Influential observations

An observation is influential if:

- It is an outlier in X and Y
 - Cook's distance $> F_{0.5}(P+1, N-P-1)$
 - DFFITS $> \frac{2\sqrt{P+1}}{\sqrt{N-P-1}}$
- Assignment Project Exam Help
<https://powcoder.com>
Add WeChat powcoder

Try analysis with and without influential observations and compare results.

Explanatory Power of a Linear Regression Equation

- Total variation is made up of two parts:

$$SST = SSR + SSE$$

Total Sum of
Squares

Regression Sum of
Squares

Error (residual)
Sum of Squares

$$SST = \sum (y_i - \bar{y})^2$$

$$SSR = \sum (\hat{y}_i - \bar{y})^2$$

$$SSE = \sum (y_i - \hat{y}_i)^2$$

where:

\bar{y} = Average value of the dependent variable

y_i = Observed values of the dependent variable

\hat{y}_i = Predicted value of y for the given x_i value

Confidence & Prediction Intervals

- Confidence interval (CI) for mean of Y
- Prediction interval (PI) for individual Y

Assignment Project Exam Help

PI is wider than CI <https://powcoder.com>

Add WeChat powcoder

Confidence Interval for the Average Y, Given X

Confidence interval estimate for the **expected value of y** given a particular x_i

Assignment Project Exam Help

Confidence interval for $E(Y_{n+1} | X_{n+1})$:

<https://powcoder.com>

Add WeChat powcoder

$$\hat{y}_{n+1} \pm t_{n-2, \alpha/2} s_e \sqrt{\frac{1}{n} + \frac{(x_{n+1} - \bar{x})^2}{\sum (x_i - \bar{x})^2}}$$

Notice that the formula involves the term $(x_{n+1} - \bar{x})^2$ so the size of interval varies according to the distance x_{n+1} is from the mean, \bar{x}

Prediction Interval for an Individual Y, Given X

Confidence interval estimate for an **actual observed value of y** given a particular x_i

Assignment Project Exam Help

Confidence interval for \hat{y}_{n+1} :
<https://powcoder.com>

Add WeChat powcoder

$$\hat{y}_{n+1} \pm t_{n-2, \alpha/2} s_e \sqrt{1 + \frac{1}{n} + \frac{(x_{n+1} - \bar{x})^2}{\sum (x_i - \bar{x})^2}}$$

This extra term adds to the interval width to reflect the added uncertainty for an individual case

Estimating Mean Values and Predicting Individual Values

Goal: Form intervals around \hat{y} to express uncertainty about the value of y for a given x_i

Confidence Interval for the expected value of y , given x_i


$$\hat{y} = b_0 + b_1 x_i$$

Prediction Interval for an single observed y , given x_i

Assignment Project Exam Help

<https://powcoder.com>


Add WeChat powcoder




Relevant Data Range

- When using a regression model for prediction, only predict within the relevant range of data


Assignment Project Exam Help



Multiple Regression



Multiple Regression




Adjusted R-Sqr

Assignment Project Exam Help

$$R_{\text{adj}}^2 = 1 - (1 - R^2) \frac{n-1}{n-m-1}$$

<https://powcoder.com>
Add WeChat powcoder


VIF



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



VIF

Assignment Project Exam Help

<https://powcoder.com>

$$VIF_i = \frac{1}{1 - R_i^2}$$

Add WeChat powcoder

scan.pdf

file:///E:/HPSCANS/scan.pdf

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Indicates deviation, or residual

20

0 5 10 X

The number $N - 2$, called the **residual degrees of freedom**, is the sample size minus the number of parameters in the line (in this case, α and β). Using $N - 2$ as a divisor in computing S^2 produces an **unbiased estimate** of σ^2 . In example:

$$RE\ S, M = \frac{17}{1} = 17$$

The square root of the residual mean square is called the **standard error of the estimate** and is denoted by S .

Software regression programs will also produce the **standard errors** of A and B . These statistics are computed as

$$SE(A) = S \left[\frac{1}{N} + \frac{\bar{X}^2}{\sum (X - \bar{X})^2} \right]^{1/2}$$
$$SE(B) = \frac{S}{[\sum (X - \bar{X})^2]^{1/2}}$$

Correlation Coefficient - ρ

- Correlation coefficient measures the strength of linear association between X and Y in the population (ρ).
- it is estimated by sample (r)

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Correlation Analysis

- **Correlation** analysis is used to measure strength of the association (linear relationship) between two variables
 - Correlation is only concerned with strength of the relationship
 - No causal effect is implied with correlation
 - Correlation was first presented in Chapter 4

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Correlation Analysis

- The population correlation coefficient is denoted ρ (the Greek letter rho)
- The sample correlation coefficient is

<https://powcoder.com>

Add WeChat powcoder

$$r = \frac{s_{xy}}{s_x s_y}$$

where

$$s_{xy} = \frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{n-1}$$

Calculating the value of ρ

- $100 (1 - \rho^2)^{1/2} = \% \text{ of Standard Deviation}$
NOT “explained” by X

Assignment Project Exam Help

<https://powcoder.com>


$$\sigma^2 = \sigma_y^2 (1 - \rho^2)$$

Add WeChat powcoder

$$\Rightarrow \sigma = \sigma_y \sqrt{1 - \rho^2}$$

$$\Rightarrow \rho^2 = \frac{\sigma_y^2 - \sigma^2}{\sigma_y^2}$$

Graphically (p 92)



Calculating the value of ρ

- $100 (1 - \rho^2)^{1/2} = \% \text{ of Standard Deviation}$
NOT “explained” by X

Assignment Project Exam Help

<https://powcoder.com>

$$\sigma^2 = \sigma_y^2 (1 - \rho^2)$$

Add WeChat powcoder

$$\Rightarrow \sigma = \sigma_y \sqrt{1 - \rho^2}$$

$$\Rightarrow \rho^2 = \frac{\sigma_y^2 - \sigma^2}{\sigma_y^2}$$

Interpretation of ρ

- ρ^2 = reduction in variance of Y associated with knowledge of X/original variance of Y
 - $100\rho^2$ = % of variance of Y “explained by X”
- <https://powcoder.com>

Caveat: correlation vs causation

Add WeChat powcoder

Estimating the value of ρ (Pearson's Correlation Coefficient)

Assignment Project Exam Help
 $\rho = \frac{\sigma_{XY}}{\sigma_X \sigma_Y}$
<https://powcoder.com>

Add WeChat powcoder
 $r = \frac{S_{XY}}{S_X S_Y}$

$$S_{XY} = \sum (X - m(X))(Y - m(Y)) / (N - 1)$$

Interpretation of ρ

ρ	% of variance “explained”	% of variance not “explained”	% of SD “explained”	% of SD not “explained”
± 0.3	9%	91%	5%	95%
± 0.5	25%	75%	13%	87%
± 0.71	50%	50%	29%	71%
± 0.95	90%	10%	69%	31%

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Test for Zero Population Correlation

- To test the null hypothesis of no linear association,

Assignment Project Exam Help

$$H_0 : \rho = 0$$

<https://powcoder.com>

the test statistic follows the Student's t distribution with $(n - 2)$ degrees of freedom:

Add WeChat powcoder

$$t = \frac{r\sqrt{(n-2)}}{\sqrt{(1-r^2)}}$$

Example from Text: Lung Function

- Data from an epidemiological study of households
 - living in four areas with different amounts and types of air pollution (Appendix A)
- Data only on non-smoking fathers
 - X = height in inches
 - Y = forced expiratory volume in 1 second (FEV1)

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Scatter Plot (p 83)




Figure 6.1: Scatter Diagram and Regression Line of FEV1 Versus Height for Fathers

Example Results

- Least Squares Equation: $Y = -4.087 + 0.118X$

Assignment Project Exam Help

- Correlation $r = 0.504$

<https://powcoder.com>

Add WeChat powcoder

- Test $p = 0$,
 - $t = 7.1$ ($p = 94$), $p < 0.0001$
 - t test can be one or two sided

Analysis of Variance

- SST = total sum of squares
 - Measures the variation of the y_i values around their mean, \bar{y}
- SSR = regression sum of squares
 - Explained variation attributable to the linear relationship between x and y
- SSE = error sum of squares
 - Variation attributable to factors other than the linear relationship between x and y

Explanatory Power of a Linear Regression Equation

- Total variation is made up of two parts:

$$SST = SSR + SSE$$

Total Sum of
Squares

Regression Sum of
Squares

Error (residual)
Sum of Squares

$$SST = \sum (y_i - \bar{y})^2$$

$$SSR = \sum (\hat{y}_i - \bar{y})^2$$

$$SSE = \sum (y_i - \hat{y}_i)^2$$

where:


\bar{y} = Average value of the dependent variable

y_i = Observed values of the dependent variable

\hat{y}_i = Predicted value of y for the given x_i value

Analysis of Variance

(continued)



Coefficient of Determination, R^2

- The **coefficient of determination** is the portion of the total variation in the dependent variable that is explained by variation in the independent variable
- The coefficient of determination is also called **R-squared** and is denoted as R^2

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

$$R^2 = \frac{SSR}{SST} = \frac{\text{regression sum of squares}}{\text{total sum of squares}}$$

note:

$$0 \leq R^2 \leq 1$$

Correlation and R^2



- The coefficient of determination, R^2 , for a simple regression is equal to the simple correlation squared

<https://powcoder.com>



Add WeChat [powcoder](https://powcoder.com)

$$R^2 = r^2$$


Examples of Approximate r^2 Values



Examples of Approximate r^2 Values



Examples of Approximate r^2 Values



Estimation of Model Error Variance

- An estimator for the variance of the population model error is

$$\hat{\sigma}_e^2 = s_e^2 = \frac{\sum_{i=1}^n e_i^2}{n-2} = \frac{SSE}{n-2}$$


- Division by $n - 2$ instead of $n - 1$ is because the simple regression model uses two estimated parameters, b_0 and b_1 , instead of one

is called the **standard error of the estimate**

$$s_e = \sqrt{s_e^2}$$

Comparing Standard Errors

s_e is a measure of the variation of observed y values from the regression line



The magnitude of s_e should always be judged relative to the size of the y values in the sample data

Statistical Inference: Hypothesis Tests and Confidence Intervals

- The variance of the regression slope coefficient (b_1) is estimated by

Assignment Project Exam Help

$$s_{b_1}^2 = \frac{s_e^2}{\sum (x_i - \bar{x})^2} = \frac{s_e^2}{(n-1)s_x^2}$$

where:

s_{b_1} = Estimate of the standard error of the least squares slope

$$s_e = \sqrt{\frac{SSE}{n-2}} = \text{Standard error of the estimate}$$

Example Results

- Least Squares Equation: $Y = -4.087 + 0.118X$

Assignment Project Exam Help

- Correlation $r = 0.504$

<https://powcoder.com>

Add WeChat powcoder

- Test $p = 0$,
 - $t = 7.1$ (p 94), $p < 0.0001$
 - t test can be one or two sided

Hypothesis Test for Population Slope Using the F Distribution

- F Test statistic:

$$F = \frac{MSR}{MSE}$$

where

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

$$MSR = \frac{SSR}{k}$$
$$MSE = \frac{SSE}{n - k - 1}$$

where F follows an F distribution with k numerator and $(n - k - 1)$ denominator degrees of freedom

(k = the number of independent variables in the regression model)

Hypothesis Test for Population Slope Using the F Distribution

(continued)

- An alternate test for the hypothesis that the slope is zero:

Assignment Project Exam Help

$$H_0: \beta_1 = 0$$

$$H_1: \beta_1 \neq 0$$

<https://powcoder.com>

- Use the F statistic

Add WeChat powcoder

$$F = \frac{MSR}{MSE} = \frac{SSR}{s_e^2}$$

- The decision rule is

$$\text{reject } H_0 \text{ if } F \geq F_{1,n-2,\alpha}$$

ANOVA Overview

Table 6.1: ANOVA table for simple linear regression

Source of variation	Sums of squares	df	Mean square	F
Regression	$\sum(\hat{Y} - \bar{Y})^2$	1	$SS_{reg}/1$	MS_{reg}/MS_{res}
Residual	$\sum(Y - \hat{Y})^2$	$N - 2$	$SS_{res}/(N - 2)$	
Total	$\sum(Y - \bar{Y})^2$	$N - 1$		

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Table 6.2: ANOVA example from Figure 6.1

Source of variation	Sums of squares	df	Mean square	F
Regression	16.0532	1	16.0532	50.50
Residual	47.0451	148	0.3179	
Total	63.0983	149		

Test $\beta = 0$

- From ANOVA table: $F = 50.5$
 - Gives 2-sided test, p-value < 0.0001
 - One sided test is: $t = F^{1/2} = 7.1$
- <https://powcoder.com>
Add WeChat powcoder

Same as test for $\rho = 0$

Outliers

- Outlier in Y is studentized (or deleted studentized) residual > 2

Assignment Project Exam Help

- Leverage = $h = \frac{1}{N} + \frac{(X - \bar{X})^2}{\sum (X - \bar{X})^2}$
 - X's far from the mean of X have large leverage (h)
 - Observations with large leverage have large effect on the slope of the line.
- Outlier in X if $h > 4/N$

Residual Analysis

- Residual = $e = Y - \hat{Y}$

Assignment Project Exam Help

- Studentized residual = $e/S(1 - h)^{1/2}$
– h called “leverage”

<https://powcoder.com>

Add WeChat powcoder

- Deleted studentized residual = studentized residual with observation for computing regression and S deleted.

Influential observations

An observation is influential if:

- It is an outlier in X and Y
- Cook's distance $> F_{0.5}(2, N-2)$
- DFFITS $> \frac{2\sqrt{2}}{\sqrt{N-2}}$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Try analysis with and without influential observations and compare results.

Observations

- Point 1 is an outlier in Y with low leverage
 - impacts estimate of intercept but not slope
 - Tends to increase the estimates of S & SE of B
- Point 2 has high leverage; not an outlier in Y
 - doesn't impact estimate of B or A
- Point 3 has high leverage and is an outlier in Y
 - impacts the values of B, A, and S

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assumptions

- Homogeneity of variance (same σ^2)
 - Not extremely serious
 - Can be achieved through transformations if necessary
- Normal residuals
 - Slight departures ok
 - Can use transformations to achieve it
- Randomness
 - Serious
 - Can use hierarchical models for clustered samples

Checking Assumptions

- Plot residuals vs X or vs the predicted Y to check linearity and homogeneity of variance
- Create normal probability plots of residuals to check for normality

Assignment Project Exam Help

<https://powcoder.com>


Add WeChat powcoder

Residual Plots (p 98)

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



Transformations (p 105)

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder




Figure 6.9: Choice of Transformation: Typical Curves and Appropriate Transformation

Weighted Regression

- If σ^2 are not equal, use weight for each residual in the sum of squares used in Least Squares process.
- Weight = $1/\sigma^2$
- Gives unbiased estimate with smaller variance

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Weighted Regression - Caveat

- Solution,, standardize weight (w) to add up to the sample size (N)
 - e.g. $N = 5$, $w = 4, 1, 8, 2, 4$, sum of $w = 19$
 - define standardized weight (sw) = $w * 5 / 19$
 - sum of $sw = 5$
 - $= 1.05 + .26 + 2.11 + .53 + 1.05 = 5$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

What to watch for

- Need representative sample
- Range of prediction should match observed range in X in sample
- Use of nominal or ordinal, rather than interval or ration data
- Errors in variables
- Correlation does not imply causation
- Violation of assumptions
- Influential points
- Appropriate model

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

Multiple Linear Regression

<https://powcoder.com>

Add WeChat powcoder

Keywords for OUTPUT Statement

Keyword	Description
COOKD= <i>names</i>	Cook's influence statistic
COVRATIO= <i>names</i>	standard influence of observation on covariance of betas
DFFITS= <i>names</i>	standard influence of observation on predicted value
H= <i>names</i>	leverage,
LCL= <i>names</i>	lower bound of a % confidence interval for an individual prediction. This includes the variance of the error, as well as the variance of the parameter estimates.
LCLM= <i>names</i>	lower bound of a % confidence interval for the expected value (mean) of the dependent variable
PREDICTED P= <i>names</i>	predicted values
PRESS= <i>names</i>	th residual divided by , where is the leverage, and where the model has been refit without the th observation.
RESIDUAL R= <i>names</i>	residuals, calculated as ACTUAL minus PREDICTED
RSTUDENT= <i>names</i>	a studentized residual with the current observation deleted
STDl= <i>names</i>	standard error of the individual predicted value
STDP= <i>names</i>	standard error of the mean predicted value
STDR= <i>names</i>	standard error of the residual
STUDENT= <i>names</i>	studentized residuals, which are the residuals divided by their standard errors
UCL= <i>names</i>	upper bound of a % confidence interval for an individual prediction
UCLM= <i>names</i>	upper bound of a % confidence interval for the expected value (mean) of the dependent variable

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Aims

- Extend simple linear regression to multiple dependent variables.
- Describe a linear relationship between:
 - A single continuous Y variable, and
 - Several X variables
- Draw inferences regarding the relationship
- Predict the value of Y from X_1, X_2, \dots, X_p .
- Research Questions: To what extent does some combination of the IVs predict the DV?
- E.g. To what extent does age, gender, type/amount of food consumption predict low density lipid level.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assumptions

- Level of Measurement:
 - IVs – two or more, Continuous or dichotomous
 - DV - continuous
- Sample Size – Enough cases per IV
- Linearity: Are bivariate relationships linear
- Constant Variance (about line of best fit) – Homoscedasticity
- Multicollinearity: Between the IVs
- Multivariate outliers
- Normality of residuals about predicted value

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Approaches

- Direct: All IVs entered simultaneously
- Forward: IVs entered one by one until there are no significant IVs to be entered.
[Assignment Project Exam Help
https://powcoder.com](https://powcoder.com)
- Backward: IVs removed one by one until there are no significant IVs to be removed.
[Add WeChat powcoder](https://powcoder.com)
- Stepwise: Combination of Forward and Backward
- Hierarchical: IVs entered in steps.

Write ups

- Assumptions: How tested, extent met
- Correlations: What are they, what conclusions
- Regression coefficients: Report and interpret
- Conclusions and Caveats

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Steps in Multiple Regression


1. State the research hypothesis.
2. State the null hypothesis
3. Gather the data
4. Assess each variable separately first (obtain measures of central tendency and dispersion; frequency distributions; graphs); is the variable normally distributed?
5. Assess the relationship of each independent variable, one at a time, with the dependent variable (calculate the correlation coefficient; obtain a scatter plot); are the two variables linearly related?
6. Assess the relationships between all of the independent variables with each other (obtain a correlation coefficient matrix for all the independent variables); are the independent variables too highly correlated with one another?
7. Calculate the regression equation from the data
8. Calculate and examine appropriate measures of association and tests of statistical significance for each coefficient and for the equation as a whole
9. Accept or reject the null hypothesis
10. Reject or accept the research hypothesis
11. Explain the practical implications of the findings

Example (p 121)

Assignment Project Exam Help


<https://powcoder.com>

Add WeChat powcoder



Assignment Project Exam Help

Add WeChat powcoder



Mathematical Model

- The mean of Y values at a given X is:

$$\alpha + \beta_1 X_1 + \beta_2 X_2 + \dots + \beta_p X_p$$

Assignment Project Exam Help

- Variance of Y values at any set of X's is σ^2

<https://powcoder.com>

(For all X)

Add WeChat powcoder

- Y values are normally distributed at each X
(needed for inference)

Types of X (independent) variables

- Fixed: selected in advance
- Variable: as in most studies
- X's can be continuous or discrete (categorical)
- X's can be transformations of other X's, e.g., polynomial regression.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Computer Analysis

- Estimates of: $\alpha, \beta_1, \beta_2, \dots, \beta_p$ using least-squares.
- Residual mean square (S^2) is estimate of variance σ^2
- Confidence intervals for mean of Y
- Prediction intervals for individual Y

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Example of Bonferroni

- Test 3 hypotheses
- P-values are: 0.014, 0.036, 0.075
- Let nominal significance level = 0.15
 - \therefore first 2 are significant
- Bonferroni Adjusted p-values multiply by 3, giving: 0.042, 0.108, 0.225
 - Only first is significant
 - Probability of at rejecting at least 1 out of m hypotheses

$$FWER = Pr \left\{ \bigcup_{i_o} (p_i \leq \frac{\alpha}{m}) \right\} \leq \sum_{i_o} \{Pr(p_i \leq \frac{\alpha}{m})\} \leq m_0 \frac{\alpha}{m} \leq m \frac{\alpha}{m} = \alpha$$

Analysis of variance (p 132)

- Does regression plane help in predicting values of Y ?

Assignment Project Exam Help

- Test hypothesis that all β_i 's = 0

<https://powcoder.com>

Table 7.1: ANOVA Table for multiple regression

Source of variation	Sums of squares	df	Mean square	F
Regression	$\sum(\hat{Y} - \bar{Y})^2$	P	SS_{reg}/P	$MS_{\text{reg}}/MS_{\text{res}}$
Residual	$\sum(Y - \hat{Y})^2$	$N - P - 1$	$SS_{\text{reg}}/(N - P - 1)$	
Total	$\sum(Y - \bar{Y})^2$	$N - 1$		

Example: Reg of FEV1 on height and weight (p 132)

Table 7.2: ANOVA example from the lung function data (fathers)

Source of variation	Sums of squares	df	Mean square	<i>F</i>
Regression	21.0570	2	10.5285	36.81
Residual	42.0413	147	0.2860	
Total	63.0983			

Assignment Project Exam Help


<https://powcoder.com>

Add WeChat powcoder

- $F = 36.81$; $df = 2, 147$; $p\text{-value} < 0.0001$
- Use percentile link from web site:
<http://faculty.vassar.edu/lowry/tabs.html#f>

Venn Diagrams

- Multiple R^2
- Bivariate Correlation between IV1 and DV
- Bivariate Correlation between IV2 and DV
- Correlation between IV1 and IV2
- Target: IV's that highly correlate with the DV, but don't highly correlate with each other



Correlation Coefficient

- The multiple correlation coefficient (R) measures the strength of association between Y, and the set of X's in the population. <https://powcoder.com>
- It is estimated as the simple correlation coefficient between the Y's and their predicted values (\hat{Y} 's)

Coefficient of Determination

- R^2 = Coefficient of determination
= SS due to regression / SS total
- R^2 = (reduction in variance of Y due to X's) / (original variance of Y).
- Therefore $100R^2$ = % of variance of Y “explained by X’s”.
- And $100(1 - R^2)^{1/2}$ = % of Standard Deviation NOT “explained” by X’s


Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Regression

Assignment Project Exam Help
<https://powcoder.com>
Add WeChat powcoder



Standard Deviation of bet1

Assignment Project Exam Help

<https://powcoder.com>

$$\sigma_{b_1} = \frac{\sigma}{\sqrt{\sum x^2 - (\sum x)^2 / n}}$$

Add WeChat powcoder

Confidence Interval Mean Value

CONFIDENCE INTERVAL FOR THE MEAN VALUE OF y FOR A GIVEN VALUE OF x

$$\hat{y}_p \pm t_{n-2}(s) \sqrt{\frac{1}{n} + \frac{(x_p - \bar{x})^2}{\sum (x_i - \bar{x})^2}}$$

where \hat{y}_p is the point estimate of y for a particular value of x , t_{n-2} a multiplier associated with the sample size and confidence level, s the standard error of the estimate, and x_p the particular value of x for which the prediction is being made.

Confidence Interval for prediction

Assignment Project Exam Help

<https://powcoder.com>

PREDICTION INTERVAL FOR A RANDOMLY CHOSEN VALUE OF y FOR A GIVEN VALUE OF x

Add WeChat powcoder

$$\hat{y}_p \pm t_{n-2}(s) \sqrt{1 + \frac{1}{n} + \frac{(x_p - \bar{x})^2}{\sum (x_i - \bar{x})^2}}$$

Adjusted R-square

Assignment Project Exam Help

<https://powcoder.com>

$$R_{\text{adj}}^2 = 1 - (1 - R^2) \frac{n - 1}{n - m - 1}$$

Add WeChat powcoder

135 of 340

MULTICOLLINEARITY 117

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder




Figure 3.8 When the predictors x_1 and x_2 are uncorrelated, the response surface y rests on a solid basis, providing stable coefficient estimates.

5:09 PM 3/22/2018

118 CHAPTER 3 MULTIPLE REGRESSION AND MODEL BUILDING

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder




Figure 3.9 Multiple Regression: A 3D Plot of the Fitted Regression Surface

Sequential SS vs. Partial SS

Assignment Project Exam Help

TABLE 3.14 Difference Between Sequential and Partial SS

Variable	Sequential SS	Partial SS
x_1	$SS(x_1)$	$SS(x_1 x_2, x_3, x_4)$
x_2	$SS(x_2 x_1)$	$SS(x_2 x_1, x_3, x_4)$
x_3	$SS(x_3 x_1, x_2)$	$SS(x_3 x_1, x_2, x_4)$
x_4	$SS(x_4 x_1, x_2, x_3)$	$SS(x_4 x_1, x_2, x_3)$

Assignment Project Exam Help

<https://powcoder.com>

120 CHAPTER 3 MULTIPLE REGRESSION AND MODEL BUILDING

Add WeChat powcoder

What effect do these changes in VIF_i have on s_{b_i} , the variability of the i th coefficient? We have

$$s_{b_i} = s c_i = s \sqrt{\frac{1}{(n-1) s_i^2} \frac{1}{1 - R_i^2}} = s \sqrt{\frac{VIF_i}{(n-1) s_i^2}}$$

If x_i is uncorrelated with the other predictors, $VIF_i = 1$, and the standard error of the coefficient s_{b_i} will not be inflated. However, if x_i is correlated with the other

VIF



large when x_i is highly correlated with the other predictors. The first factor, $1 / ((n - 1)s_i^2)$, measures the variance of the i th predictor, x_i . It is the second factor, $1 / (1 - R_i^2)$, which measures the variance inflation between the i th predictor and the remaining predictors. The second factor is denoted as the *variance inflation factor*.

Add WeChat powcoder

$$VIF_i = \frac{1}{1 - R_i^2}$$

behavior of the VIF? Suppose that x_i is uncorrelated with the other predictors, so that $R_i^2 = 0$. Then we will have



Interpretation of R

R	% of variance “explained”	% of variance not “explained”	% of SD “explained”	% of SD not “explained”
± 0.3	9%	91%	5%	95%
± 0.5	25%	75%	13%	87%
± 0.71	50%	50%	29%	71%
± 0.95	90%	10%	69%	31%

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Partial Correlation

- The correlation coefficient measuring the degree of dependence between two variables
 - after adjusting for the linear effect of one or more of the other X variables

Example: T_1 and T_2 are test scores


- Find partial R between T_1 and T_2 after adjusting for IQ

Visually (p 130)

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



- Partial R = simple R between the two residuals

Interpretation of regression coefficients

- In the model: $\alpha + \beta_1 X_1 + \beta_2 X_2 + \dots + \beta_p X_p$ if ρ is the partial correlation between Y and X_1 , given X_2, \dots, X_p , then
- Testing that $\beta_1 = 0$ is equivalent to testing that $\rho = 0$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Hence, β_1 is called the partial regression coefficient of Y on X_1 , given X_2, \dots, X_p

Values of regression coefficients

- Problem: Values of β_i 's are not directly comparable
- Hence: Standardized coefficients:
 - Standardized $\beta_i = \beta_i * (SD(X_i) / SD(Y))$
- Standardized β_i are directly comparable.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Multicollinearity

$$[\text{SE}(B_i)]^2 = \frac{S^2}{(N-1)(S_i)^2} \times \frac{1}{1-(R_i)^2}$$

- The case where some of the X variables are highly correlated
- This will impact estimates and their SE's (p 143)
- Consider Tolerance, and its inverse, Variance Inflation Factor
- Target Tolerance < 0.01, or VIF > 100
- Remedy: use variable selection to delete some X variables, or a dimension reduction techniques such as Principal Components.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Misleading Correlations

- Example (Lung Function data, Appendix A):
FEV1 vs height and age
- Depends on gender

Assignment Project Exam Help


<https://powcoder.com>

Add WeChat powcoder


Total vs Stratified Correlation

Gender	Correlation between FEV1 and	
	Height	Age
Total	0.739	-0.073
Male	0.504	-0.310
Female	0.465	-0.267


FEV1 vs height




FEV1 vs height – Regression lines



FEV1 vs age



FEV1 vs age– Regression lines



Residual Analysis

- Residual = $e = Y - \hat{Y}$

Assignment Project Exam Help

- Studentized residual = $e/S(1 - h)^{1/2}$
– h called “leverage”

<https://powcoder.com>

Add WeChat powcoder

- Deleted studentized residual = studentized residual with observation for computing regression and S deleted.

Outliers

- Outlier in Y is studentized (or deleted studentized) residual > 2 (same as simple case)
[Assignment Project Exam Help](#)
- Outlier in X if $h_i > 2(P+1)/N$
<https://powcoder.com>
Add WeChat powcoder

Some Caveats

- See list for simple regression
- Need representative sample
- Violations of assumptions, outliers
- Multicollinearity: coefficient of any one variable can vary widely, depending on what others are included in the model
- Missing values
- Number of observations in the sample should be large enough relative to number of variables in the model.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Outline

- Matrix Review: $(A - \lambda I) X = 0$; Eigenvalues
- Simple linear regression
- Visit **Assignment Project Exam Help**
<http://www.atucla.edu/stat/sas/output/reg.htm>
<https://powcoder.com>
- Assign HW 6.1,2,5 for next week
Add WeChat powcoder
- If we get to Chapter 7, assign HW 7.2, 7.4, 7.5, 7.6
(Hand in 7.2,4,5) 7.7 Will be assigned next week.
- Start Multiple Regression Lecture
- Go over Multiple Regression Example – 7.1

Quick Matrix Review

$$(A - \lambda I) X = 0$$

Assignment Project Exam Help
<https://powcoder.com>

$$A = \begin{pmatrix} 3 & 1 \\ 2 & 2 \end{pmatrix} \quad (A - \lambda I) = \begin{pmatrix} 3 - \lambda & 1 \\ 2 & 2 - \lambda \end{pmatrix}$$

$\lambda = 1, 4$ Add WeChat powcoder

$$\lambda = 1 \Rightarrow y = -2x$$

$$\lambda = 4 \Rightarrow y = x$$

Quick Matrix Review

$$(A - \lambda I) X = 0$$

$$A = \begin{bmatrix} 3 & 1 & 2 \\ 2 & 2 & 5 \\ 1 & 3 & 2 \end{bmatrix} \quad (A - \lambda I) = \begin{bmatrix} 3-\lambda & 1 & 2 \\ 2 & 2-\lambda & 5 \\ 1 & 3 & 2-\lambda \end{bmatrix}$$

Add WeChat powcoder


$$(3-\lambda)(2-\lambda)(2-\lambda) + 1*5*1 + 3*3*2 - ((1*(2-\lambda)*3) + (2*1*(2-\lambda)) + ((3-\lambda)*5*3)) = 0$$

$$(12 - 16\lambda + 7\lambda^2 - \lambda^3 + 5 + 18) - ((6 - 3\lambda) + (4 - 2\lambda) + (45 - 15\lambda)) = 0$$

$$-20 + 4\lambda + 7\lambda^2 - \lambda^3 = 0$$

$$\lambda = 7.17, -1.76, 1.59$$

Analysis of Variance



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder