

<https://powcoder.com>

Assignment Project Exam Help

Add WeChat powcoder

ST227 Survival Models - Part III

Assignment Project Exam Help
Estimation in the Markov Model

<https://powcoder.com>

Add WeChat powcoder

George Tzougas

Department of Statistics,
London School of Economics

January 31, 2019

<https://powcoder.com>

1 Markov Models: Data and Estimation

Assignment Project Exam Help
Add WeChat powcoder

1.1 Introduction

Assignment Project Exam Help

- We now have mathematical **models for various situations that arise in insurance**, e.g. a model of mortality with states $\{Alive, Dead\}$, a model for PHI with states $\{Well, Ill, Dead\}$.
<https://powcoder.com>
Add WeChat powcoder
- These **models depend on transition intensities** μ_x^{gh} , $g \neq h$ between states g and h .
- The values of these **transition intensities are unknown** - they are parameters in the models.

<https://powcoder.com>

Assignment Project Exam Help

- Thus, we need some data which we can use to estimate the values of the transition intensities μ_{gh}

Add WeChat powcoder

- We will start with the 2-state model of mortality with states $\{Alive, Dead\}$, and work through this example in detail.

Assignment Project Exam Help

<https://powcoder.com>

- It turns out that the answers for the 2-state model give a very good guide to what happens in the multi-state case.

Add WeChat powcoder

<https://powcoder.com>

Assignment Project Exam Help

1.2 Data and notation for the 2-state model

Add WeChat powcoder

- Suppose we have data on n independent lives all aged between x and $x + 1$. We suppose for life i :

Assignment Project Exam Help

$x + \alpha_i$ = age at which observation begins, $0 < \alpha_i < 1$

$x + b_i$ = age at which observation ends, if life i does not die, $0 \leq b_i < 1$

$x + t_i$ = age at which observation stops, by death or censoring

It will often be useful to work with the time spent under observation, so we define

v_i = $t_i - \alpha_i$, the observed waiting time

Also, we define

d_i = 1, if life i is observed to die

d_i = 0, if life i is not observed to die

<https://powcoder.com>

Assignment Project Exam Help

Note:

Add WeChat powcoder

- *Censoring is a form of missing data problem in which time to event is not observed for reasons such as termination of study before all recruited subjects have shown the event of interest or the subject has left the study prior to experiencing an event.*

<https://powcoder.com>

- *Censoring is the key feature of survival data (indeed survival analysis might be defined as the analysis of censored data) and the mechanisms which give rise to censoring play an important part in statistical inference.*
- *Censoring is present when we do not observe the exact length of a lifetime, but observe only that its length falls within some interval. This can happen in several ways which we are going to discuss in detail in the following lectures.*

<https://powcoder.com>

Assignment Project Exam Help

The following diagrams may be helpful.

Add WeChat powcoder

Case 1. The i th **life does not die**, i.e. reaches the end of the observation period at $x + b_i$.

Assignment Project Exam Help

<https://powcoder.com>




In this case, $d_i = 0$, $t_i = b_i$, $v_i = b_i - a_i$.

<https://powcoder.com>

Assignment Project Exam Help

Case 2. The i th **life dies** at $x + t_i$ before the end of the observation period.

Add WeChat powcoder



Add WeChat powcoder

In this case, $d_i = 1$, $\alpha_i < t_i < b_i$, $v_i = t_i - \alpha_i$.

<https://powcoder.com>

1.3 The random variables D_i and V_i

Add WeChat powcoder

In the simple two state model the observations on the i th life are: whether the life died or not, i.e. the value d_i , and the observed waiting time, i.e. the value v_i . We can think of the observations d_i and v_i as realisations of random variables D_i and V_i . Inference for the transition intensity μ_x will depend on the joint distribution of D_i and V_i . Before deriving this distribution we make two important points:

- D_i and V_i are not independent, since as it was previously shown
- V_i has a mixed distribution, i.e. the probability distribution has a **continuous density** for values of v_i which correspond to **death**, i.e. for

<https://powcoder.com>

Assignment Project Exam Help

$v_i < b_i - \alpha_i$, and a mass of probability (i.e. discrete distribution) for the value of v_i which corresponds to survival, i.e. for $v_i = b_i - \alpha_i$.

Add WeChat powcoder

Assumption: Before deriving the joint distribution of D_i and V_i we will make a further assumption. We assume that $\mu_{x+t} = \mu_x = \mu$ for $0 \leq t \leq 1$, i.e. the transition intensity μ_{x+t} is constant from age x to age $x + 1$. We can now prove the following result:

Assignment Project Exam Help

<https://powcoder.com>

Result: The joint probability function of D_i and V_i can be written

Add WeChat powcoder

$$f_i(d_i, v_i) = e^{-\mu v_i} \mu^{d_i} \quad (1)$$

where $\mu_{x+t} = \mu$ for $0 \leq t \leq 1$.

<https://powcoder.com>

Assignment Project Exam Help

Proof: We consider the two cases: $D_i = 0$ and $D_i = 1$.

Add WeChat powcoder

Case 1: $D_i = 0$. We need the probability the life survives from $x + \alpha_i$ to $x + b_i$. We have

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

<https://powcoder.com>

Assignment Project Exam Help

Add WeChat powcoder

$$P_r(D_i = 0, V_i = b_i - \alpha_i)$$

$$= P_r(\text{Life } i \text{ survives from } x + \alpha_i \text{ to } x + b_i)$$

Assignment Project Exam Help

$$= \exp\left(-\int_0^{v_i} \mu_{x+\alpha_i+s} ds\right), \text{ where } v_i = b_i - \alpha_i$$

<https://powcoder.com>

$$= \exp(-\mu v_i), \text{ since } \mu_{x+\alpha_i+s} = \mu \text{ is constant}$$

$$= e^{-\mu v_i} \mu^{d_i}, \text{ since } d_i = 0 \quad (2)$$

Case 2: $D_i = 1$. We need the probability density function that a life aged $x + \alpha_i$ dies at time $x + \alpha_i + v_i$. We have

<https://powcoder.com>

Assignment Project Exam Help

$$\begin{aligned} P_r(D_i = 1, v_i \leq V_i < v_i + dv_i) &= \int_{v_i}^{v_i + dv_i} p_{x+\alpha_i} \times q_{x+\alpha_i+v_i} dv_i \\ &= \exp\left(-\int_0^{v_i} \mu_{x+\alpha_i+s} ds\right) \times (\mu_{x+\alpha_i+v_i} dv_i + o(dv_i)) \\ &= \exp(-\mu v_i) \times (\mu dv_i + o(dv_i)) \text{ since } \mu_{x+t} = \mu \text{ is constant} \\ &= e^{-\mu v_i} \mu^{d_i} dv_i + o(dv_i), \text{ since } d_i = 1. \end{aligned} \quad (3)$$

Dividing Eq.(3) by dv_i and taking the limit as $dv_i \rightarrow 0$, we get:

$$f_i(d_i, v_i) = e^{-\mu v_i} \mu^{d_i}. \quad (4)$$

<https://powcoder.com>

Assignment Project Exam Help

- Thus, in both cases, Eq.(2) and Eq.(4), we can write the probability function as

$$f_i(d_i, v_i) = e^{-\mu v_i} \mu^{d_i} \quad (5)$$

Assignment Project Exam Help


- But what does the mixed distribution of D_i and V_i look like? Here is a simple example: take $a_i = 0$, $b_i = 1$, i.e. the observation period is the full year of life, unless death occurs.
- Suppose $\mu = 0.004$, the approximate force of mortality for a male life aged 50.
- The probability of survival is $e^{-0.004} = 0.996$. The density function for a death is $\mu e^{-\mu t} = 0.004 e^{-0.004 t}$, $0 \leq t \leq 1$.

<https://powcoder.com>

Assignment Project Exam Help

- In the plot of the mixed distribution below note that the probability mass for survival at 1 is drawn on a different scale from the density function for death.

Assignment Project Exam Help



<https://powcoder.com>

Assignment Project Exam Help

1.4 The maximum likelihood estimate of μ

Add WeChat powcoder

Suppose that the waiting times of the n lives are $v' = (v_1, v_2, \dots, v_n)$ and the record of deaths is $d' = (d_1, d_2, \dots, d_n)$. The n lives are independent so, from (5), the total likelihood is

Assignment Project Exam Help

<https://powcoder.com>

$$L(\mu; D, v) = \prod_{i=1}^n f_i(d_i, v_i) = \prod_{i=1}^n \mu e^{-\mu v_i} \mu^{d_i} e^{-\mu \sum v_i} \mu^{\sum d_i}. \quad (6)$$

We are interested in maximizing $L(\mu; D, v)$ with respect to μ .

<https://powcoder.com>

Assignment Project Exam Help

Firstly, we write $L(\mu; D, v)$ in more compact form as:

Add WeChat powcoder

$$L(\mu) = e^{-\mu v} \mu^d, \quad (7)$$

Assignment Project Exam Help

where

<https://powcoder.com>

Add WeChat powcoder

$$\begin{aligned} v &= \sum v_i = \text{Total waiting time,} \\ d &= \sum d_i = \text{Total number of deaths.} \end{aligned} \quad (8)$$

<https://powcoder.com>

Assignment Project Exam Help

Then, the log-likelihood will be:

Add WeChat powcoder

$$\begin{aligned}l(\mu) &= \log(L(\mu)) = \log(e^{-\mu v} \mu^d) \\&= \log(e^{-\mu v}) + \log(\mu^d) \\&= -\mu v + d \log(\mu)\end{aligned}\tag{9}$$

Assignment Project Exam Help

We need to find the maximum by finding the derivative:

Add WeChat powcoder

$$\frac{\partial l(\mu)}{\partial \mu} = -v + \frac{d}{\mu} = 0,\tag{10}$$

The maximum likelihood estimate of μ follows from (10) as

<https://powcoder.com>

Assignment Project Exam Help

$$\hat{\mu} = \frac{d}{v} = \frac{\text{Total number of deaths}}{\text{Total waiting time}} \quad (11)$$

- We are interested in the **statistical properties of $\hat{\mu}$** . For this purpose, we **think of $L(\mu)$ and $\hat{\mu}$** , see Eqs (7 and 11) respectively, **as random variables.**

<https://powcoder.com>

- Corresponding to Eq.(7) we define the random variable

$$L(\mu; D, V) = e^{-\mu V} \mu^D, \quad (12)$$

where $D = \sum D_i$ is the random total number of deaths and $V = \sum V_i$ is the random *total waiting time*.

<https://powcoder.com>

Assignment Project Exam Help

- Corresponding to Eq.(11) we define the estimator

Add WeChat powcoder

$$\tilde{\mu} = \frac{D}{V}. \quad (13)$$

Assignment Project Exam Help

Comment: The previous **maximum likelihood estimation** procedure concerned **lives all aged between x and $x + 1$** , e.g. from 20 to 21. An obvious question is: "How do we extend the estimation process to a wider range of ages, e.g. from 20 to 70?" One way of doing this is to split the age range into a number of short age ranges in such a way that the μ_x can be assumed constant within each sub-range. Unfortunately, this solution creates another problem: there is no reason to expect that the large number of estimates of μ_x for different x -values will be consistent. For example, we expect mortality to increase, at least from age 20, say. We will be forced to smooth (or **graduate**) these raw rates.

<https://powcoder.com>

Assignment Project Exam Help

1.5 The Maximum Likelihood Theorem (MTL)

Add WeChat powcoder

- As was previously mentioned we are interested in investigating the **statistical properties of $\hat{\mu}$** , such as **asymptotic unbiasedness, asymptotic efficiency and asymptotic normality**. For this purpose, we defined $L(\mu; D, V)$ and μ as the random variables corresponding to $L(\mu)$ and $\hat{\mu}$.

<https://powcoder.com>

Add WeChat powcoder

- In order to be able to carry on with our investigation, we have make use of two important general results from mathematical statistics:
 - properties of the **score function**, and
 - the **Maximum Likelihood Theorem**.

<https://powcoder.com>

Assignment Project Exam Help

Here we give a brief review of this material. First, we consider the one parameter case, we maximize the log-likelihood with respect to the unknown parameter θ .

Add WeChat powcoder

Definition: The **score function**. Suppose $\mathbf{X}' = (X_1, X_2, \dots, X_n)$ is a vector of random variables where the X_i are independent random variables with probability function $f_i(x, \theta)$ which depends on a parameter μ

Assignment Project Exam Help

<https://powcoder.com>

Let

Add WeChat powcoder

$$l = l(\theta) = l(\theta, \mathbf{X}) = \sum_{i=1}^n \log f_i(X_i, \theta) \quad (14)$$

be the **log-likelihood function (viewed as a random variable)**. Then

<https://powcoder.com>

Assignment Project Exam Help

Add WeChat powcoder

$$U = U(\theta) = \frac{\partial l}{\partial \theta} \quad (15)$$

is called the **score function**.

Assignment Project Exam Help

Note: The score function is **additive** as follows: let

<https://powcoder.com>

Add WeChat powcoder

$$U_i = U_i(\theta) = \frac{\partial l_i}{\partial \theta} = \frac{\partial}{\partial \theta} \log f_i(X_i, \theta) \quad (16)$$

be the score function for the i th observation, $i = 1, \dots, n$. Then

<https://powcoder.com>

Assignment Project Exam Help

Add WeChat [powcoder](https://powcoder.com)

(17)

Result: The score function satisfies.

<https://powcoder.com>

(a) $E(U) = 0$, and

Add WeChat [powcoder](https://powcoder.com)

(b) $Var(U) = -E\left(\frac{\partial U}{\partial \theta}\right) = -E\left(\frac{\partial^2 l}{\partial \theta^2}\right).$

<https://powcoder.com>

Assignment Project Exam Help

Proof: First, take $n = 1$ and without loss of generality $f_i(x, \theta) = f(x, \theta)$ (or for simplicity's sake f)

Add WeChat powcoder

(a) Consider the identity

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

$$\int f(x, \theta) dx = 1.$$

(18)

Differentiate (18) with respect to θ . We get

$$\int \frac{\partial f}{\partial \theta} dx = 0.$$

<https://powcoder.com>

Assignment Project Exam Help

Rewrite this equation as

Add WeChat powcoder

$$\int \left(\frac{\partial}{\partial \theta} \log f \right) f dx = 0. \quad (19)$$

Assignment Project Exam Help

Note that, $U = \frac{\partial}{\partial \theta} \log f$ and $E(U) = E\left(\frac{\partial}{\partial \theta} \log f\right)$. Thus, the left hand side of this equation gives $E(U) = 0$ for $n = 1$. Hence $E(U) = 0$ for all n by the additivity property, see Eq. (17).

(b) The proof uses the same trick. In particular we differentiate Eq. (19), with respect to θ and we get

<https://powcoder.com>

Assignment Project Exam Help

Add WeChat powcoder

$$\frac{\partial}{\partial \theta} \left\{ \int \left(\frac{\partial}{\partial \theta} \log f \right) f dx \right\} = 0 \quad (20)$$

Hence, using the product rule we have that

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

<https://powcoder.com>

Assignment Project Exam Help

$$\int \left\{ \frac{\partial}{\partial \theta} \left(\frac{\partial}{\partial \theta} \log f \right) f + \left(\frac{\partial}{\partial \theta} \log f \right) \frac{\partial f}{\partial \theta} \right\} dx = 0$$

$$\Rightarrow \int \frac{\partial}{\partial \theta} \left(\frac{\partial}{\partial \theta} \log f \right) f dx + \int \left(\frac{\partial}{\partial \theta} \log f \right) \left(\frac{\partial}{\partial \theta} \log f \right) f dx = 0$$

$$\Rightarrow E \left(\frac{\partial U}{\partial \theta} \right) + E(U^2) = 0, \text{ since } U = \frac{\partial}{\partial \theta} \log f$$

Add WeChat powcoder

$$\Rightarrow E(U^2) = -E \left(\frac{\partial U}{\partial \theta} \right)$$

$$\Rightarrow \text{Var}(U) = E(U^2) - E^2(U) = -E \left(\frac{\partial U}{\partial \theta} \right), \text{ since } E(U) = 0$$

$$\text{or } \text{Var}(U) = -E \left(\frac{\partial^2 l}{\partial \theta^2} \right), \text{ since } \frac{\partial U}{\partial \theta} = \frac{\partial^2}{\partial \theta^2} \log f = \frac{\partial^2 l}{\partial \theta^2}$$

<https://powcoder.com>

Assignment Project Exam Help

and (b) is proved for $n = 1$. The general case follows by the additivity of the score function, see Eq.(17) and the independence of the observations.

- It is important to note that the score function U is a fundamental quantity in statistics and it is connected with the properties of the maximum likelihood estimate of θ . In particular,
 - The MLE satisfies $\partial l / \partial \theta = 0$ i.e. $U = 0$
 - The asymptotic variance of the MLE is found from $Var(U)$.
- In what follows we will define the **Fisher information function** (sometimes simply called information function) based on U .

<https://powcoder.com>

Assignment Project Exam Help

- The **Fisher information function** is a way of measuring the amount of information that an observable random variable X carries about an unknown parameter θ of a distribution that models X . Formally, **it is the variance of the score** $Var(U)$, or $-E\left(\frac{\partial U}{\partial \theta}\right)$, since as was proved in (b) $Var(U) = -E\left(\frac{\partial U}{\partial \theta}\right)$.

Assignment Project Exam Help

<https://powcoder.com>

Definition: Fisher's information function $I(\theta)$ is

Add WeChat powcoder

$$I(\theta) = -E\left(\frac{\partial^2 l}{\partial \theta^2}\right) = -E\left(\frac{\partial U}{\partial \theta}\right) = Var(U) \quad (21)$$

We usually just call $I(\theta)$ the information function. We can now state (without proof) the **Maximum Likelihood Theorem (MLT)**.

<https://powcoder.com>

Assignment Project Exam Help

Theorem: Suppose $\hat{\theta}$ satisfies $U = 0$ and that θ_0 is the true (but unknown) value of θ . If $\tilde{\theta}$ is the random variable corresponding to $\hat{\theta}$, then $\tilde{\theta}$ is

(i) asymptotically unbiased, i.e. $E(\tilde{\theta}) \rightarrow \theta_0$,

Assignment Project Exam Help

(ii) asymptotically efficient, i.e. $\text{Var}(\tilde{\theta}) \rightarrow I(\theta_0)^{-1}$,

Add WeChat powcoder

(iii) asymptotically normal, i.e.

$$\tilde{\theta} \approx \mathcal{N}(\theta_0, I(\theta_0)^{-1}). \quad (22)$$

Of course, θ_0 is unknown so how do we calculate $I(\theta_0)^{-1}$? There are two main approximations.

<https://powcoder.com>

Assignment Project Exam Help

- Replace θ_0 with θ , i.e.

Add WeChat powcoder

$$I(\theta_0) \approx I(\hat{\theta}) = -E \left(\frac{\partial U}{\partial \theta} \right) \Big|_{\theta=\hat{\theta}} = -E \left(\frac{\partial^2 l}{\partial \theta^2} \right) \Big|_{\theta=\hat{\theta}}$$

<https://powcoder.com>

- If we can't work out the expectation in $E \left(\frac{\partial U}{\partial \theta} \right)$ then try

Add WeChat powcoder

$$I(\theta_0) \approx - \left(\frac{\partial^2 l}{\partial \theta^2} \right) \Big|_{\theta=\hat{\theta}} \quad (23)$$

<https://powcoder.com>

Assignment Project Exam Help

1.6 Properties of $\tilde{\mu} = D/V$ in the 2-state model

Add WeChat powcoder

The maximum likelihood theorem is a most powerful and general result and we apply it now to the problem of determining the properties of the MLE of the force of mortality in the 2-state model

Assignment Project Exam Help

We know from section 1.4 that

<https://powcoder.com>

Add WeChat powcoder

$$\begin{aligned} L(\mu) &= e^{-\mu V} \mu^D \\ \Rightarrow l(\mu) &= -\mu V + D \log \mu \\ \Rightarrow U(\mu) &= -V + \frac{D}{\mu} \\ \Rightarrow \tilde{\mu} &= \frac{D}{V} \end{aligned} \tag{24}$$

<https://powcoder.com>

Assignment Project Exam Help

Now apply the Maximum Likelihood Theorem. Suppose that μ_0 is the true, but unknown, value of the unknown force of mortality.

(i) First, $\tilde{\mu}$ is asymptotically unbiased so

Assignment Project Exam Help

<https://powcoder.com>

$E(\tilde{\mu}) \rightarrow \mu_0$
Add WeChat powcoder

(25)

(ii) Second, $\tilde{\mu}$ is asymptotically efficient so

<https://powcoder.com>

Assignment Project Exam Help

$$\begin{aligned} \text{Var}(\tilde{\mu}) &\longrightarrow I(\mu_0)^{-1} \\ &= \left\{ -E \left(\frac{\partial U}{\partial \mu} \right) \right\}^{-1} \Big|_{\mu=\mu_0} \\ &= - \left\{ -E \left(-\frac{D}{\mu^2} \right) \right\}^{-1} \Big|_{\mu=\mu_0}, \quad \text{by differentiating (24)} \\ &= \left(\frac{E(D)}{\mu^2} \right) \Big|_{\mu=\mu_0} \\ &= \frac{\mu^2}{E(D)} \Big|_{\mu=\mu_0} \end{aligned}$$

<https://powcoder.com>

Assignment Project Exam Help

Hence,

$$Var(\mu) \longrightarrow \frac{\mu^2}{E(D)} \Big|_{\mu=\mu_0}. \quad (26)$$

Now $E(U) = 0$ so from (24) we get

$$E\left(\frac{D}{\mu} - V\right) = 0 \Rightarrow \frac{E(D)}{\mu} = E(V) \Rightarrow E(D) = \mu E(V). \quad (27)$$

Thus, from Eqs(26 and 27) we get

$$Var(\tilde{\mu}) \longrightarrow \frac{\mu}{E(V)} \Big|_{\mu=\mu_0} = \frac{\mu_0}{E(V)}. \quad (28)$$

<https://powcoder.com>

Assignment Project Exam Help

(iii) Third, $\tilde{\mu}$ is asymptotically normal.

Add WeChat powcoder

Our final result is

Assignment Project Exam Help

$$\tilde{\mu} \approx N\left(\mu_0, \frac{\mu_0}{E(V)}\right) \quad (29)$$

Add WeChat powcoder

Comments: We note the following

- The observed waiting time $v = \sum v_i$ is usually called the **central exposed to risk** E_x^c in actuarial language.

<https://powcoder.com>

Assignment Project Exam Help

- In practice, we evaluate $Var(\tilde{\mu}) = \mu_0/E(V)$ by

Add WeChat powcoder

$$Var(\tilde{\mu}) = \frac{\mu_0}{E(V)} \approx \frac{\hat{\mu}}{v} = \frac{d/v}{v} = \frac{d}{v^2} \quad (30)$$

Assignment Project Exam Help

How does the estimation result (29) for the 2-state model extend to more complicated multi-state models such as Illness/Death models?

Add WeChat powcoder


1.7 Estimation in the 3-state model

In Part II we considered the following 3-state model for working, retiring and dying.

<https://powcoder.com>

Assignment Project Exam Help

Add WeChat powcoder



Suppose we have data on n lives and for the i th life we observe

<https://powcoder.com>

Assignment Project Exam Help

$$\begin{aligned} d_i &= \begin{cases} 0 & \text{when } i\text{th life does not die} \\ 1 & \text{when } i\text{th life does die} \end{cases} \\ r_i &= \begin{cases} 0 & \text{when } i\text{th life does not retire} \\ 1 & \text{when } i\text{th life does retire} \end{cases} \\ v_i &= t_i - a_i, \quad \text{the } i\text{th waiting time} \end{aligned} \tag{31a}$$

<https://powcoder.com>

- Our observation is the triple (v_i, d_i, r_i) with corresponding underlying random variable (V_i, D_i, R_i) . The distribution of (V_i, D_i, R_i) is given by

$$Pr(D_i = d_i, R_i = r_i, v_i < V_i < v_i + dv_i). \tag{32}$$

<https://powcoder.com>

Assignment Project Exam Help

- Notice that once more this is a mixed distribution with a mass of probability corresponding to remaining in the working state and a continuous density function corresponding to retirement or death.

- The key result is ${}_tP_x^{ww} = {}_tP_x^{ww}$ (occupation probability) since in this case state w cannot be re-entered once it has been left).

$${}_tP_x^{ww} = \exp\left(-\int_0^t (\mu + \nu) ds\right) = e^{-(\mu + \nu)t} \quad (33)$$

We discuss the three cases: (i) remaining working, (ii) dying, (iii) retiring.

Case 1: Remaining working. In this case $d_i = 0$, $r_i = 0$ and the life remains in the working state from $x + \alpha_i$ to $x + b_i$ so we have

<https://powcoder.com>

Assignment Project Exam Help

$$\begin{aligned} & P_r(D_i = 0, R_i = 0, V_i = b_i - \alpha_i) \\ &= P_r(\text{Neither died nor retired during } x + \alpha_i \longrightarrow x + b_i) \\ &= {}^{ww}_{b_i - \alpha_i} p_x \\ &= e^{-(\mu + \nu)(b_i - \alpha_i)}, \text{ from (35) for } t = b_i - \alpha_i \\ &= e^{-(\mu + \nu)v_i} \end{aligned}$$

since $v_i = b_i - \alpha_i$ when $d_i = 0, r_i = 0$.

Hence, the probability density function (pdf) in this case is

$$f_i(0, 0, v_i) = e^{-(\mu + \nu)v_i}. \quad (34)$$

<https://powcoder.com>

Assignment Project Exam Help

Case 2: Dying. In this case $d_i = 1$, $r_i = 0$, $v_i < V_i < v_i + dv_i$ so we have

Add WeChat powcoder

$$\begin{aligned} Pr(D_i &= 1, R_i = 0, v_i < V_i < v_i + dv_i) \\ &= Pr(\text{Neither died nor retired during } (x + \alpha_i, x + \alpha_i + v_i) \dots \\ &\quad \dots \text{ and then died in } (x + \alpha_i + v_i, x + \alpha_i + v_i + dv_i)) \\ &= v_i p_{x+\alpha_i}^{ww} \times_{dv_i} q_{x+\alpha_i+v_i} \\ &= e^{-(\mu+\nu)v_i} \mu dv_i \end{aligned}$$

Dividing by dv_i and taking the limit we find the pdf is

$$f_i(1, 0, v_i) = e^{-(\mu+\nu)v_i} \mu. \quad (35)$$

<https://powcoder.com>

Assignment Project Exam Help

Add WeChat powcoder

Case 3: Retiring. In this case $d_i = 0$, $r_i = 1$, $v_i < V_i < v_i + dv_i$ so by symmetry with case 2 we have

Assignment Project Exam Help

<https://powcoder.com>

$$Pr(D_i = 0, R_i = 1, v_i < V_i < v_i + dv_i) = e^{-(\mu+\nu)v_i} \times (\nu dv_i + o(dv_i))$$

Add WeChat powcoder

and the pdf in this case is

$$f_i(0, 1, v_i) = e^{-(\mu+\nu)v_i} \nu. \quad (36)$$

In summary, see Eqs(34,35 and 36) we have

<https://powcoder.com>

Assignment Project Exam Help

- Case 1: $d_i = 0, r_i = 0 \Rightarrow f_i(0, 0, v_i) = e^{-(\mu+\nu)v_i},$

Add WeChat powcoder

- Case 2: $d_i = 1, r_i = 0 \Rightarrow f_i(1, 0, v_i) = e^{-(\mu+\nu)v_i}\mu,$

Assignment Project Exam Help

- Case 3: $d_i = 0, r_i = 1 \Rightarrow f_i(0, 1, v_i) = e^{-(\mu+\nu)v_i}\nu.$

<https://powcoder.com>

All three cases can conveniently be packaged as

Add WeChat powcoder

$$f_i(d_i, r_i, v_i) = e^{-(\mu+\nu)v_i} \times \mu^{d_i} \times \nu^{r_i} \quad (37)$$

and so, assuming that all n lives are independent, we obtain the likelihood as

<https://powcoder.com>

Assignment Project Exam Help

Add WeChat powcoder

$$L(\mu, \nu) = \prod_{i=1}^n e^{-(\mu+\nu)v_i} \times \mu^{d_i} \times \nu^{r_i}$$

$$L(\mu, \nu) = e^{-\mu v} \mu^d \times e^{-\nu v} \nu^r \quad (38)$$

Assignment Project Exam Help

where $v = \sum v_i$ is the total waiting time, $d = \sum d_i$ is the total number of deaths and $r = \sum r_i$ is the total number of retirements.

Add WeChat powcoder

Notice that $L(\mu, \nu)$ **factorises** into a product of two functions, one of μ and one of ν . As a result the log-likelihood has the particularly simple form

$$l(\mu, \nu) = -\mu v + d \log \mu - \nu v + r \log \nu \quad (39)$$

<https://powcoder.com>

Assignment Project Exam Help

and the estimators of μ and ν are easily obtained by computing the partial derivatives of the log-likelihood with respect to μ and ν and setting them equal to 0:

Assignment Project Exam Help

$$\tilde{\mu} = \frac{D}{V}, \quad \tilde{\nu} = \frac{R}{V}.$$

(40)

<https://powcoder.com>

Comment: There are obvious parallels between

- (37) and the likelihood $L(\mu) = e^{\mu\nu} \mu^d$ in the 2-state model, and
- (39) and the estimator $\tilde{\mu} = D/V$ in the 2-state model.

<https://powcoder.com>

Assignment Project Exam Help


1.8 More estimation in the 3-state model

Add WeChat powcoder

Our second example of the 3-state model is the model for PHI discussed in Part II. Suppose that the transition intensities are constant for the n lives in the investigation and consider the diagram:

Assignment Project Exam Help

<https://powcoder.com>



<https://powcoder.com>

Assignment Project Exam Help

Suppose that the observation on the i th life is

Add WeChat powcoder

$$v_i = \text{total waiting time in state } h \quad (41)$$

$$\omega_i = \text{total waiting time in state } i$$

$$s_i = \text{number of transitions from } h \text{ to } i$$

$$r_i = \text{number of transitions from } i \text{ to } h$$

$$d_i = \begin{cases} 1 & \text{if } i \longrightarrow d, 0 \text{ otherwise} \end{cases}$$

$$u_i = \begin{cases} 1 & \text{if } i \longrightarrow d, 0 \text{ otherwise.} \end{cases}$$

A straightforward extension of the argument in the previous example for the i th life gives the likelihood. An example may help.

<https://powcoder.com>

Assignment Project Exam Help

Example: Consider a life that is initially healthy, falls ill after time t_1 , recovers after a further time t_2 and then dies after a further time t_3 . As always we consider the events falling ill, recovering and dying as occurring in a short interval of time, as in the diagram



The probability of observing this sequence of events is

$$e^{-(\mu+\sigma)t_1} [\sigma\delta t_1 + o(\delta t_1)] \times e^{-(\rho+\nu)t_2} [\rho\delta t_2 + o(\delta t_2)] \times e^{-(\mu+\sigma)t_3} [\mu\delta t_3 + o(\delta t_3)] .$$

<https://powcoder.com>

Assignment Project Exam Help

Dividing by $\delta t_1 \delta t_2 \delta t_3$ and taking the limit as $\delta t_i \rightarrow 0$, $i = 1, 2, 3$ we find the contribution to the likelihood from this life is

$$e^{-(\mu+\sigma)(t_1+t_3)} e^{-(\rho+\nu)t_2} \mu^s \sigma^r \nu^u \quad (42)$$

<https://powcoder.com>

The general formula for the likelihood $L(\mu, \nu, \sigma, \rho)$ follows by multiplying up over all lives:

Add WeChat powcoder

$$\begin{aligned} & \prod_{i=1}^n e^{-(\sigma+\mu)v_i} \times e^{-(\rho+\nu)\omega_i} \times \sigma^{s_i} \times \rho^{r_i} \times \mu^{d_i} \times \nu^{u_i} \\ &= e^{-(\sigma+\mu)v} e^{-(\rho+\nu)\omega} \sigma^s \rho^r \mu^d \nu^u. \end{aligned}$$

<https://powcoder.com>

Assignment Project Exam Help

Once more the likelihood factorises into separate functions and so the estimators are easily obtained by computing the partial derivatives of the log-likelihood with respect to μ and ν and setting them equal to 0:

Assignment Project Exam Help

$$\tilde{\mu} = \frac{D}{V}, \quad \tilde{\nu} = \frac{U}{W}, \quad \tilde{\sigma} = \frac{S}{V}, \quad \tilde{\rho} = \frac{R}{W} \quad (43)$$

<https://powcoder.com>

The estimates in all these examples have the form

$$\frac{\text{MLE of transition intensity } g \rightarrow h}{=} \frac{\text{No.transfers from } g \text{ to } h}{\text{Total waiting time in } g}. \quad (44)$$

<https://powcoder.com>

Assignment Project Exam Help

1.9 The likelihood in multi-state models

Add WeChat powcoder


It is important to write down the likelihood for a multi-state model. Thus, to emphasise this we collect the results of all our examples.

Assignment Project Exam Help

Example 1: Simple two state model; see 1.3 and 1.4.

<https://powcoder.com>

Add WeChat powcoder



where v is the total waiting time in the alive state and d is the total number of deaths. Then

<https://powcoder.com>

Assignment Project Exam Help

Add WeChat powcoder


(45)

Example 2: Retiral; see 1.7.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



where v is the total waiting time in the working state, d is the total number of deaths and r is the total number of retirals. Then

<https://powcoder.com>


Assignment Project Exam Help

Add WeChat powcoder

(46)

Example 3: PHI; see 1.8.

Assignment Project Exam Help



where v and w are total waiting times and r, s, d and u are total numbers of transfers between states as shown. Then

<https://powcoder.com>

Assignment Project Exam Help

$$L(\mu, \nu) = e^{-\mu} e^{-\nu} \times e^{-\mu} e^{-\nu} \times \mu^d \times \nu^u \times \sigma^s \times \rho^r. \quad (47)$$

1.10 The likelihood methodology

<https://powcoder.com>

These examples allow us to formulate a general methodology as follows:

[Add WeChat powcoder](https://powcoder.com)

(a) Each **waiting time** ω_i in state i contributes a factor

$$e^{-\left(\sum_j \mu_{ij}\right) \omega_i}, \quad (48)$$

<https://powcoder.com>

Assignment Project Exam Help

where $\sum_j \mu_{ij}$ is the sum of the transition intensities **out of state i** .

Add WeChat powcoder

(b) Each **transfer number** t_{ij} from state i to state j contributes a factor

Assignment Project Exam Help

μ_{ij}^{tj} ,
<https://powcoder.com>

(49)

where μ_{ij} is the transition intensity from state i to state j

Add WeChat powcoder

(c) The total likelihood is the product of all the above contributions.

A tip for the exam: You should be able to apply this rule and derive the likelihood for simple examples, as done in lectures/classes.

<https://powcoder.com>

1.11 Assignment Project Exam Help Properties of the MLE's

Add WeChat powcoder

We saw in the simple two state model



<https://powcoder.com>

Add WeChat powcoder

that

$$\tilde{\mu} = \frac{D}{V} = \frac{\text{No.deaths}}{\text{Total waiting time}} \quad (50)$$

<https://powcoder.com>

Assignment Project Exam Help

and that asymptotically

Add WeChat powcoder

$$\tilde{\mu} \approx \mathcal{N}(\mu_0, \frac{\mu_0}{E(V)}). \quad (51)$$

Assignment Project Exam Help

The proof consisted of applying the Maximum Likelihood Theorem (MLT) together with some properties of the score function.


Add WeChat powcoder

<https://powcoder.com>

Assignment Project Exam Help

For the model for retiral we have the likelihood $L(\mu, \nu)$

Add WeChat powcoder



Add WeChat powcoder

$$L(\mu, \nu) = e^{-\mu v} \mu^d \times e^{-\nu v} \nu^r \quad (52)$$

where v is the total waiting time in the working state, d is the number of deaths and r is the number of retireals.

<https://powcoder.com>

Assignment Project Exam Help

What is the **joint distribution** of $\tilde{\mu}$ and $\tilde{\nu}$? We need to **extend the MLT to the multiparameter case**

Add WeChat powcoder

The following parallels the one parameter case for the simple two state model in 1.5.

Assignment Project Exam Help

<https://powcoder.com>

1.12 Multi-parameter version of the Maximum Likelihood Theorem (MTL)

Add WeChat powcoder

Definition: The **score function**. Suppose $\mathbf{X}' = (X_1, X_2, \dots, X_n)$ is a vector of random variables where the X_i are independent random variables with probability function $f_i(x, \theta)$ which depends on a parameter $\theta' = (\theta_1, \dots, \theta_r)$ with r components. Let

<https://powcoder.com>

Assignment Project Exam Help

Add WeChat powcoder

$$l = l(\theta) = l(\theta, \mathbf{X}) = \sum_{i=1}^n \log f_i(X_i, \theta) \quad (53)$$

be the log-likelihood function (viewed as a random variable). Also, let

<https://powcoder.com>

Add WeChat powcoder

$$U_i = U_i(\theta) = \frac{\partial l}{\partial \theta_i}, i = 1, \dots, r \quad (54)$$

Then $\mathbf{U}' = (U_1, \dots, U_r)$ is called the **score function**.

Result: The score function satisfies:

<https://powcoder.com>

Assignment Project Exam Help

(a) $E(\mathbf{U}) = 0$, and

Add WeChat powcoder

(b) The variance-covariance matrix is $Var(\mathbf{U}) = -E\left(\frac{\partial \mathbf{U}}{\partial \boldsymbol{\theta}}\right) = (v_{ij})_{r \times r}$,
where $v_{ij} = -E\left(\frac{\partial^2 l}{\partial \theta_i \partial \theta_j}\right)$ are its elements.

Assignment Project Exam Help

For example, when we have two parameters θ_1 and θ_2 , we get

<https://powcoder.com>

Add WeChat powcoder

$$Var(\mathbf{U}) = \begin{bmatrix} -E\left(\frac{\partial^2 l}{\partial \theta_1^2}\right) & -E\left(\frac{\partial^2 l}{\partial \theta_1 \partial \theta_2}\right) \\ -E\left(\frac{\partial^2 l}{\partial \theta_2 \partial \theta_1}\right) & -E\left(\frac{\partial^2 l}{\partial \theta_2^2}\right) \end{bmatrix}$$

Proof: The proof for (a) and (b) is pretty much the same as the one-parameter case. Firstly, we take $n = 1$ and write $f_i = f$.

<https://powcoder.com>

Assignment Project Exam Help

(a) Consider the identity

Add WeChat powcoder

$$\int f(x, \theta) dx = 1. \quad (55)$$

Differentiate with respect to θ_i . We get

Add WeChat powcoder

$$\int \frac{\partial f}{\partial \theta_i} dx = 0. \quad (56)$$

Rewrite this equation as

<https://powcoder.com>

Assignment Project Exam Help

Add WeChat powcoder

$$\int \left(\frac{\partial}{\partial \theta_i} \log f \right) f dx = 0 \quad (57)$$

The left hand side of this equation involves $E(U_i)$, so $E(U) = 0$ as required.

<https://powcoder.com>

- (b) The proof uses the same trick as in the one-parameter case. Differentiate (57) with respect to θ_i and use the product rule. We get

Add WeChat powcoder

<https://powcoder.com>

Assignment Project Exam Help

$$\Rightarrow \int \left\{ \frac{\partial}{\partial \theta_j} \left(\frac{\partial}{\partial \theta_i} \log f \right) f + \left(\frac{\partial}{\partial \theta_i} \log f \right) \frac{\partial f}{\partial \theta_j} \right\} dx = 0$$

$$\Rightarrow \int \frac{\partial}{\partial \theta_j} \left(\frac{\partial}{\partial \theta_i} \log f \right) f dx + \int \left(\frac{\partial}{\partial \theta_i} \log f \right) \left(\frac{\partial}{\partial \theta_j} \log f \right) f dx = 0$$

<https://powcoder.com>

$$\Rightarrow \int \frac{\partial U_i}{\partial \theta_j} f dx + \int U_i U_j f dx = 0$$

$$\Rightarrow E(U_i U_j) = -E \left(\frac{\partial U_i}{\partial \theta_j} \right), \text{ since } U_i = \frac{\partial}{\partial \theta_i} \log f$$

$$\Rightarrow \text{Cov}(U_i, U_j) = E(U_i U_j) - E(U_i)E(U_j) = -E \left(\frac{\partial^2 l}{\partial \theta_i \partial \theta_j} \right), \text{ since } E(U_i) = 0$$

<https://powcoder.com>

Assignment Project Exam Help

Hence, (b) is proved.

Add WeChat powcoder

- The case of general n follows by the additivity of the score function and the independence of the X_i

Assignment Project Exam Help

- The information function $I(\theta)$ extends as follows:

Add WeChat powcoder

Definition: Fisher's **information function** $I(\theta)$ is

$$I(\theta) = -E \left(\frac{\partial^2 l}{\partial \theta_i \partial \theta_j} \right) = -E \left(\frac{\partial \mathbf{U}}{\partial \theta} \right) = \text{Var}(\mathbf{U}). \quad (58)$$

<https://powcoder.com>

Assignment Project Exam Help

As before we usually call $I(\theta)$ the information function. We can now state (without proof) the multi-parameter version of the **MLT**.

Theorem: Suppose $\hat{\theta}$ satisfies $U = 0$ and that θ_0 is the true (but unknown) value of θ . If $\tilde{\theta}$ is the random variable corresponding to $\hat{\theta}$, then $\tilde{\theta}$ is

- <https://powcoder.com>
Add WeChat powcoder
- (i) asymptotically unbiased, i.e. $E(\tilde{\theta}) \longrightarrow \theta_0$,
 - (ii) asymptotically efficient, i.e. $Var(\tilde{\theta}) \longrightarrow I(\theta_0)^{-1}$,
 - (iii) asymptotically normal, i.e we have

$$\tilde{\theta} \approx \mathcal{N}(\theta_0, I(\theta_0)^{-1}) \quad (59)$$

<https://powcoder.com>

1.13 Assignment Project Exam Help Examples of MLE's

Add WeChat powcoder

Example 1: Retiral/Death model. We found the likelihood for this model in 1.7 as

Assignment Project Exam Help

$$L(\mu, \nu) = e^{-\mu v} \mu^d \times e^{-\nu v} \nu^r \quad (60)$$

Add WeChat powcoder

which gives

$$l(\mu, \nu) = -\mu v + d \log \mu - \nu v + r \log \nu \quad (61)$$

<https://powcoder.com>

Assignment Project Exam Help

as the log-likelihood and

Add WeChat powcoder

$$\tilde{\mu} = \frac{D}{V}, \quad \tilde{\nu} = \frac{R}{V} \quad (62)$$

Assignment Project Exam Help

as the estimators of μ and ν . Now the asymptotic distribution of $\tilde{\mu}$ and $\tilde{\nu}$ is determined by the information matrix $I(\mu, \nu)$.

Add WeChat powcoder

Observe that in this case is given by

$$I(\mu, \nu) = \begin{bmatrix} -E \left(\frac{\partial^2 l}{\partial \mu^2} \right) & -E \left(\frac{\partial^2 l}{\partial \mu \partial \nu} \right) \\ -E \left(\frac{\partial^2 l}{\partial \nu \partial \mu} \right) & -E \left(\frac{\partial^2 l}{\partial \nu^2} \right) \end{bmatrix}$$

<https://powcoder.com>

Assignment Project Exam Help

Add WeChat powcoder

$$U_{\mu} = \frac{\partial l}{\partial \mu} = -v + \frac{d}{\mu}, \text{ thus} \quad (63)$$

$$\mu E(v) = E(d), \text{ since } E(U_{\mu}) = 0,$$

Assignment Project Exam Help

from (63) $\frac{\partial^2 l}{\partial \mu^2} = -\frac{d}{\mu^2}$, thus

<https://powcoder.com>

Add WeChat powcoder

$$-E\left(\frac{\partial^2 l}{\partial \mu^2}\right) = -E\left(-\frac{d}{\mu^2}\right) = \frac{E(d)}{\mu^2} = \frac{\mu E(v)}{\mu^2} = \frac{E(v)}{\mu}, \quad (64)$$

from (63) $\frac{\partial^2 l}{\partial \mu \partial \nu} = 0$, thus

$$-E\left(\frac{\partial^2 l}{\partial \mu \partial \nu}\right) = 0. \quad (65)$$

<https://powcoder.com>

Assignment Project Exam Help

Similarly, since

Add WeChat powcoder

$$U_v = \frac{\partial W}{\partial v} = -v + \frac{r}{v}, \text{ we have} \quad (66)$$

$$-E \left(\frac{\partial^2 l}{\partial v^2} \right) = \frac{E(v)}{v} \text{ and} \quad (67)$$

Assignment Project Exam Help

$$-E \left(\frac{\partial^2 l}{\partial \mu \partial v} \right) = 0 \quad (68)$$

<https://powcoder.com>

Hence, using Eqs(64, 65 and 67, 68) we find that

Add WeChat powcoder

$$I(\mu, \nu) = \begin{bmatrix} \frac{E(V)}{\mu} & 0 \\ 0 & \frac{E(V)}{v} \end{bmatrix} \quad (69)$$

Thus, $I(\mu, \nu)$ is **diagonal** and so $\tilde{\mu}$ and $\tilde{\nu}$ are **asymptotically independent**.

<https://powcoder.com>

Assignment Project Exam Help

N.B. The estimators will always be asymptotically independent whenever the log-likelihood can be expressed as the sum of separate functions of the parameters as in Eq.(61).

Therefore,

Assignment Project Exam Help

<https://powcoder.com>

$$\tilde{\mu} \approx \mathcal{N}\left(\mu_0, \frac{\mu_0}{E(V)}\right) \text{ and} \quad (70)$$

$$\tilde{\nu} \approx \mathcal{N}\left(\nu_0, \frac{\nu_0}{E(V)}\right), \text{ thus} \quad (71)$$

$\tilde{\mu}$ and $\tilde{\nu}$ are asymptotically independent

Example 2: PHI model. We found the likelihood for this model in 1.8 as

<https://powcoder.com>

Assignment Project Exam Help

$$L(\mu, \nu, \rho, \sigma) = e^{-(\sigma + \nu)\nu} e^{-(\sigma + \nu)\omega} \sigma^s \rho^r \mu^d \nu^u \quad (72)$$

which gives

Assignment Project Exam Help

$$\tilde{\mu} = \frac{D}{V}, \quad \tilde{\nu} = \frac{U}{W}, \quad \tilde{\sigma} = \frac{S}{V}, \quad \tilde{\rho} = \frac{R}{W}. \quad (73)$$

as the estimators of μ, ν, σ and ρ .

Once more the likelihood factorises into separate functions and so the asymptotic distribution of the estimators is easily obtained. First, since $I(\mu, \nu, \rho, \sigma)$ is diagonal, $\tilde{\mu}, \tilde{\nu}, \tilde{\sigma}$ and $\tilde{\rho}$ are asymptotically independent. Moreover,

<https://powcoder.com>

Assignment Project Exam Help

Add WeChat powcoder
 $\mu \approx \mathcal{N}\left(\mu_0, \frac{\mu_0}{E(V)}\right)$

with corresponding results for $\tilde{\nu}$, $\tilde{\sigma}$ and $\tilde{\rho}$.

Assignment Project Exam Help

Conclusion: The Markov model gives simple estimates of the transition intensities with simple properties. We need to keep track of only two things:

<https://powcoder.com>

Add WeChat powcoder

- the number of transfers between states, and
- the total time spent in each state.

This simplicity is part of the appeal of this class of models.