

Assignment Project Exam Help

Bayesian Learning

<https://powcoder.com>

Prof. Dan A. Simovici

UMB
Add WeChat powcoder

1 Probability Spaces

Assignment Project Exam Help

2 Random Variables

3 Conditional Probabilities

<https://powcoder.com>

4 ML and Conditional Probabilities

Add WeChat powcoder

5 Bayes Theorem and Concept Learning

Assignment Project Exam Help


Suppose that (Ω, \mathcal{E}, P) is a probability space, \mathcal{E} is a family of subsets of Ω known as **events**, and P is a probability. The elements of Ω are **elementary events**.

In many cases, \mathcal{E} consists of all subsets of Ω , and we will make this assumption unless a special statement says otherwise.

Add WeChat powcoder

Example

Rolling two dice is described by a finite probability space that consists of 36 elementary events: $(1, 1), (1, 2), \dots, (6, 6)$.



An **event** in the previous example is a subset of $\{1, \dots, 6\} \times \{1, \dots, 6\}$, that is, a subset of the set of pairs $\{(x, y) \mid 1 \leq x \leq 6, 1 \leq y \leq 6\}$.

Example

- throws that have the same number of both dice:

$$S = \{(1, 1), (2, 2), (3, 3), (4, 4), (5, 5), (6, 6)\}$$

- throws such that the sum of the numbers is greater than 8:

$$B = \{(2, 6), (3, 5), (3, 6), (4, 4), (4, 5), (4, 6), (5, 3), (5, 4), (5, 6), (6, 6), (6, 5)\}$$

Note that Ω consists of 36 elementary events and there are $2^{36} \approx 10^{12}$ events in this very simple probability space

Probability of an event V in this context is the number $P(V)$ given by

$$P(V) = \frac{|V|}{|\Omega|}.$$

Example

We have

$$P(S) = \frac{6}{36} = \frac{1}{6}.$$

and

$$P(B) = \frac{18}{36} = \frac{1}{2}$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

Informally, Borel sets of \mathbb{R} are the sets that can be constructed from open or closed sets by repeatedly taking countable unions and intersections.

Definition

Let (Ω, \mathcal{E}, P) be a probability space. A function $X: \Omega \rightarrow \mathbb{R}$ is a **random variable** if $X^{-1}(U) \in \mathcal{E}$ for every Borel subset of \mathbb{R} .

Add WeChat powcoder

Definition

A simple random variable is defined by a table:

$$X : \begin{pmatrix} x_1 & x_2 & \cdots & x_n \\ p_1 & p_2 & \cdots & p_n \end{pmatrix},$$

where x_1, \dots, x_n are the values assumed by X and $p_i = P(X = x_i)$ for $1 \leq i \leq n$. We always have $p_1 + \cdots + p_n = 1$.

The expected value of X is

$$E[X] = x_1 p_1 + x_2 p_2 + \cdots + x_n p_n.$$

Assignment Project Exam Help

Example

A random variable X whose distribution is:

$$X \sim \begin{pmatrix} 0 & 1 \\ q & p \end{pmatrix},$$

<https://powcoder.com>

where $p + q = 1$ is said to have a **Bernoulli distribution** with parameter p .

Note that $E[X] = p$ and $\text{var}(X) = pq$.

Add WeChat powcoder

Example

Let $p, q \in [0, 1]$ be two numbers such that $p + q = 1$. Consider the random variable defined by

$$X: \left(0, \binom{n}{1} q^{n-1} p, \dots, \binom{n}{k} q^{n-k} p^k, \dots, n \right),$$

We refer to a random variable with this distribution as a *binomial random variable*. Note that

$$q^n + \binom{n}{1} q^{n-1} p + \dots + \binom{n}{k} q^{n-k} p^k + \dots + p^n = (q + p)^n = 1.$$

Example cont'd

Assignment Project Exam Help

The expectation of a binomial variable is

$$E[X] = np.$$

<https://powcoder.com>

The variance of a random variable X is

$$\text{var}(X) = E[(X - E(X))^2] = E[X^2] - (E[X])^2.$$

Add WeChat powcoder

In the case of a binomial variable the variance is $\text{var}(X) = npq$.

The Characteristic Function of an Event

Assignment Project Exam Help

If A is an event, then the function $1_A: \Omega \rightarrow \{0, 1\}$ defined by

$$1_A(\omega) = \begin{cases} 1 & \text{if } \omega \in A, \\ 0 & \text{otherwise,} \end{cases}$$

<https://powcoder.com>

is a random variable,

Add WeChat (1 - P(A), P(A)) powcoder

Note that $E(1_A) = P(A)$ and $\text{var}(1_A) = P(A)(1 - P(A))$.

The event $A \wedge B$ takes place when both A and B occur; the event $A \vee B$ takes place when at least one of A and B occur.

Example

The event $S \wedge B$ takes place when the result of throwing the dice results in a pair of numbers (n, n) whose sum is greater than 8 and consists of the pairs:

$$(4, 4), (5, 5), (6, 6)$$

Therefore $P(S \wedge B) = \frac{3}{36} = \frac{1}{12}$.

Definition

If B is an event such that $P(B) > 0$ one can define the probability of an event A conditioned on B as

$$P(A|B) = \frac{P(A \cap B)}{P(B)}.$$

Example

<https://powcoder.com>

The probability of the event S conditioned on B is

$$P(S|B) = \frac{P(S \cap B)}{P(B)} = \frac{\frac{1}{12}}{\frac{1}{2}} = \frac{1}{6}$$

and

$$P(B|S) = \frac{P(S \cap B)}{P(S)} = \frac{\frac{1}{12}}{\frac{1}{6}} = \frac{1}{2}.$$

Definition

Two events A, B are independent if $P(A \wedge B) = P(A)P(B)$.

If A, B are independent events, then

$$P(A|B) = \frac{P(A \wedge B)}{P(B)} = \frac{P(A)P(B)}{P(B)} = P(A)$$

and

$$P(B|A) = \frac{P(B \wedge A)}{P(A)} = \frac{P(B)P(A)}{P(A)} = P(B).$$

Note that B and S are independent events because

$$P(B \wedge S) = \frac{1}{12} = P(B)P(S).$$

- The **product rule** or the **Bayes theorem**:

$$P(A \wedge B) = P(A|B)P(B) = P(B|A)P(A).$$

Assignment Project Exam Help

- The **sum rule**:

$$P(A \vee B) = P(A) + P(B) - P(A \wedge B).$$

- The **total probability rule**: if A_1, \dots, A_n are mutually exclusive and $\sum_{i=1}^n P(A_i) = 1$, then

$$P(B) = \sum_{i=1}^n P(B|A_i)P(A_i).$$

Assignment Project Exam Help

In ML we are often interested in determining the best hypothesis from some space H given the observed data S .

“Best” means in this context, the most probable hypothesis given

- the data S and
- any initial knowledge of **prior** probabilities of hypotheses in H .

Add WeChat powcoder

Assignment Project Exam Help

- “Prior probabilities” (or a priori probabilities) mean probabilities of hypotheses **before** seeing the data S .
- “Posterior probabilities” mean probabilities of hypotheses **after** seeing the data S .

<https://powcoder.com>

If no prior knowledge exist all hypotheses have the same probability.
In ML we are interested to compute $P(h|S)$ that h holds given the observed training data S .

Add WeChat powcoder

Bayes' Theorem in ML

Assignment Project Exam Help

For a sample S and a hypothesis h we have

$$P(h|S) = \frac{P(S|h)P(h)}{P(S)}$$

<https://powcoder.com>

Note that:

- $P(h|S)$ increases with $P(h)$ and with $P(S|h)$.
- $P(h|S)$ decreases with $P(S)$, because the more probable is that S will be observed independent of h , the less evidence S provides for h .

Learning Scenario

Consider some set of candidate hypotheses H and seek the most probable hypothesis given the observed data S .

Any such maximally probable hypothesis is called a **maximum a posteriori hypothesis**, **MAP**.

h_{MAP} is <https://powcoder.com>

$$h_{MAP} = \operatorname{argmax}_{h \in H} P(h|S)$$

$$= \operatorname{argmax}_{h \in H} \frac{P(S|h)P(h)}{P(S)}$$

$$= \operatorname{argmax}_{h \in H} P(S|h)P(h)$$

because $P(S)$ is a constant.

Maximum Likelihood Hypothesis

Assignment Project Exam Help

In some cases we assume that every hypothesis of H is **apriori equally probable**, that is, $P(h_i) = P(h_j)$ for all $h_i, h_j \in H$.

Now,

$$h_{MAP} = \operatorname{argmax}_{h \in H} P(S|h).$$

$P(S|h)$ is known as the **likelihood** of S given h .

Add WeChat powcoder

Example

A medical diagnosis problem:

The hypothesis space contains two hypotheses:

- h_0 : patient has no cancer;
- h_1 : patient has cancer.

An imperfect diagnosis test that has two outcomes, \oplus and \ominus .

$$P(\oplus|h_1) = 0.98 \quad P(\oplus|h_0) = 0.03$$

$$P(\ominus|h_1) = 0.02 \quad P(\ominus|h_0) = 0.97$$

Prior knowledge: Only 0.08% of population has cancer; 99.2% does not.

Example (cont'd)

Assignment Project Exam Help

The test returns \oplus . Should we conclude that the patient has cancer?

The **MAP** hypothesis is obtained as

$$h_{MAP} = \operatorname{argmax}_{h \in H} P(\mathcal{I}|h)P(h).$$

<https://powcoder.com>

$$\begin{aligned} P(\oplus|h_1)P(h_1) &= 0.98 * 0.008 = 0.0078, \\ P(\oplus|h_0)P(h_0) &= 0.03 * 0.992 = 0.0298. \end{aligned}$$

Add WeChat powcoder

The **MAP** hypothesis is h_0 ; the patient has no cancer.

Brute-Force Bayes Concept Learning

Assignment Project Exam Help

- For each hypothesis $h \in H$ calculate the posterior probability:

$$P(h|S) = \frac{P(D|h)P(h)}{P(S)}$$

- Output the hypothesis h_{MAP} with

$$h_{MAP} = \arg\max_{h \in H} P(h|S).$$

Assignment Project Exam Help

Assumption for the Brute-Force Bayes Concept Learning

- Training data is $S = \{(x_1, y_1), \dots, (x_m, y_m)\}$, where $y_i = f(x_i)$ for $1 \leq i \leq m$ and it is noise-free.
- The target hypothesis is contained in H .
- We have no apriori reason to believe that any hypothesis is more probable than the other

<https://powcoder.com>
Add WeChat powcoder

Consequences

Assignment Project Exam Help

- $P(h) = \frac{1}{|H|}$;
- The probability of S given h is 1 if S is consistent with h and 0 otherwise.

<https://powcoder.com>

$$P(S|h) = \begin{cases} 1 & \text{if } y_i = h(x_i) \text{ for } 1 \leq i \leq m \\ 0 & \text{otherwise;} \end{cases}$$

Add WeChat powcoder

Assignment Project Exam Help

- Let $VS_{H,S}$ be the subset of hypotheses of H that is consistent with S .
- If S is inconsistent with h then $P(h|S) = \frac{0 \cdot P(h)}{P(S)} = 0$.
 - If S is consistent with h then

<https://powcoder.com>

$$P(h|S) = \frac{1}{P(S)} = \frac{1}{\frac{|VS_{H,S}|}{|H|}} = \frac{1}{|VS_{H,S}|}$$

Add WeChat powcoder

Since the hypotheses are mutually exclusive (that is, $P(h_i \wedge h_j) = 0$ if $i \neq j$), by the total probability law:

$$P(S) = \sum_{h_i \in H} P(S|h_i)P(h_i)$$

$$= \sum_{h \in VS_{H,S}} 1 \cdot \frac{1}{|H|} + \sum_{h \notin VS_{H,S}} 0 \cdot \frac{1}{|H|}$$

$$= \sum_{h \in VS_{H,S}} 1 \cdot \frac{1}{|H|} = \frac{|VS_{H,S}|}{|H|}.$$

Add WeChat powcoder

Note that under this setting every consistent hypothesis is a MAP hypothesis.