

Regression
(Module 5)

Assignment Project Exam Help

<https://powcoder.com>



Add WeChat powcoder

Statistics (MAST20005) &
Elements of Statistics
(MAST90038)

School of Mathematics and Statistics
University of Melbourne

Semester 2, 2022

Aims of this module

- Introduce the concept of **regression**
- Show a simple model for studying the relationship between two variables
- Discuss correlation and how it relates to regression

<https://powcoder.com>

Add WeChat powcoder

Outline

Introduction

Regression

Simple linear regression

Point estimation of the mean

Interlude: Analysis of variance

Point estimation of the variance

Standard errors of the estimates

Confidence intervals

Prediction intervals

R example

Model checking

Further regression models

Correlation

Definitions

Point estimation

Relationship to regression

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Relationships between two variables

We have studied how to do estimation for some simple scenarios:

- iid samples from a single distribution (X_i)
- comparing iid samples from two different distributions (X_i & Y_j)
- differences between paired measurements ($X_i - Y_i$)

We now consider how to analyse bivariate data more generally, i.e. two variables, X and Y , measured at the same time, i.e. as a pair.


The data consist of pairs of data points, (x_i, y_i) .

These can be visualised using a **scatter plot**.

Example data

x_i	y_i
1.80	9.18
1.40	7.66
2.10	6.33
0.30	4.51
3.60	14.04
0.70	4.94
1.10	4.24
2.10	8.19
0.90	4.55
3.80	11.57

$n = 10$



Outline

Introduction

Regression

Simple linear regression

Point estimation of the mean

Interlude: Analysis of variance

Point estimation of the variance

Standard errors of the estimates

Confidence intervals

Prediction intervals

R example

Model checking

Further regression models

Correlation

Definitions

Point estimation

Relationship to regression

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Regression

Often interested in how Y depends on X . For example, we might want to use X to predict Y .

In such a setting, we will assume that the X values are known and fixed (henceforth, x instead of X), and look at how Y varies given x .

Example: Y is a student's final mark for Statistics, and x is their mark for the prerequisite subject Probability. Does x help to predict Y ?

The regression of Y on x is the conditional mean, $\mathbb{E}(Y | x) = \mu(x)$.

The regression can take any form. We consider **simple linear regression**, which has the form of a **straight line**:

$$\mathbb{E}(Y | x) = \alpha + \beta x \quad \text{and} \quad \text{var}(Y | x) = \sigma^2.$$

Example: simple linear regression model


Assignment Project Exam Help

$$E(Y | x) = \alpha + \beta x$$

$$\text{var}(Y | x) = \sigma^2$$

<https://powcoder.com>

Add WeChat powcoder



Terminology

- Y is called a **response** variable. Can also be called an **outcome** or **target** variable. Please do **not** call it the 'dependent' variable.
- x is called a **predictor** variable. Can also be called an **explanatory** variable. Please do **not** call it an 'independent' variable.
- $\mu(x)$ is called the (linear) **predictor function** or sometimes the **regression curve** or the **model equation**.
- The parameters in the predictor function are called **regression coefficients**.

Why 'regression'?

It is strange terminology, but it has stuck.

Refers to the idea of 'regression to the mean':

if a variable is extreme on its first measurement, it will tend to be closer to the average on its second measurement, and vice versa.

First described by Sir Francis Galton when studying the inheritance of height between fathers and sons. In doing so, he invented the technique of simple linear regression.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Linearity

A regression model is called **linear** if it is linear in the coefficients.

It doesn't have to define a straight line!

Complex and non-linear functions of x are allowed, as long as the resulting predictor function is a linear combination (i.e. an additive function) of them, with the coefficients 'out the front'.

For example, the following are linear models:

$$\mu(x) = \alpha + \beta x + \gamma x^2$$

$$\mu(x) = \frac{\alpha}{x} + \frac{\beta}{x^2}$$

$$\mu(x) = \alpha \sin x + \beta \log x$$

The following are NOT linear models:

$$\mu(x) = \alpha \sin(\beta x)$$

Assignment Project Exam Help

$$\mu(x) = \frac{\alpha}{1 + \beta x}$$

$$\mu(x) = \alpha x^\beta$$

<https://powcoder.com>

... but the last one can be re-expressed as a linear model on a log scale (by taking logs of both sides),

Add WeChat powcoder

$$\mu^*(x) = \alpha^* + \beta^* \log x$$

Outline

Introduction

Regression

Simple linear regression

Point estimation of the mean

Interlude: Analysis of variance

Point estimation of the variance

Standard errors of the estimates

Confidence intervals

Prediction intervals

R example

Model checking

Further regression models

Correlation

Definitions

Point estimation

Relationship to regression

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Estimation goals

Back to our simple linear regression model:

$$\mathbb{E}(Y | x) = \alpha + \beta x \quad \text{and} \quad \text{var}(Y | x) = \sigma^2.$$

- We wish to estimate the slope (β), the intercept (α), the variance of the errors (σ^2), their standard errors and construct confidence intervals for these quantities.
- Often want to use the fitted model to make predictions about future observations (i.e. predict Y for a new x).
- Note: the Y_i are not iid. They are independent but have different means, since they depend on x_i .
- We have not (yet) assumed any specific distribution for Y , only a conditional mean and variance.

Reparameterisation

Changing our model slightly.

Let $\alpha_0 = \alpha + \beta\bar{x}$, which gives:

$$\begin{aligned}\mathbb{E}(Y | x) &= \alpha + \beta x \\ &= \alpha_0 + \beta(x - \bar{x})\end{aligned}$$

<https://powcoder.com>

Now our model is in terms of α_0 and β .

Add WeChat powcoder

This will make calculations and proofs simpler.

Least squares estimation

Choose α_0 and β to minimize the sum of squared deviations:

$$H(\alpha_0, \beta) = \sum_{i=1}^n (y_i - \alpha_0 - \beta(x_i - \bar{x}))^2$$

Solve this by finding the partial derivatives and setting to zero:

$$0 = \frac{\partial H(\alpha_0, \beta)}{\partial \alpha_0} = 2 \sum_{i=1}^n [y_i - \alpha_0 - \beta(x_i - \bar{x})](-1)$$
$$0 = \frac{\partial H(\alpha_0, \beta)}{\partial \beta} = 2 \sum_{i=1}^n [y_i - \alpha_0 - \beta(x_i - \bar{x})](-(x_i - \bar{x}))$$

These are called the **normal equations**.

Least squares estimators

Some algebra yields the least square estimators,

$$\hat{\alpha}_0 = \bar{Y}, \quad \hat{\beta} = \frac{\sum_{i=1}^n (x_i - \bar{x}) Y_i}{\sum_{i=1}^n (x_i - \bar{x})^2}.$$

Another expression for $\hat{\beta}$ is

$$\hat{\beta} = \frac{\sum_{i=1}^n (x_i - \bar{x})(Y_i - \bar{Y})}{\sum_{i=1}^n (x_i - \bar{x})^2}.$$

These are equivalent, due to the following result:

$$\sum (x_i - \bar{x})(Y_i - \bar{Y}) = \sum (x_i - \bar{x}) Y_i.$$

Can also then get an estimator for α :

$$\hat{\alpha} = \hat{\alpha}_0 - \hat{\beta}\bar{x}$$

Assignment Project Exam Help

And also an estimator for the predictor function,

$$\hat{\mu}(x) = \hat{\alpha} + \hat{\beta}x$$

$$= \hat{\alpha}_0 + \hat{\beta}(x - \bar{x})$$

$$= \bar{Y} + \hat{\beta}(x - \bar{x}).$$

Add WeChat powcoder

Ordinary least squares

This method is sometimes called ordinary least squares or OLS.

Other variants of least squares estimation exist, with different names.

For example, 'weighted least squares'.

<https://powcoder.com>

Add WeChat powcoder

Example: least squares estimates

For our data:

$$\bar{x} = 1.78$$

$$\bar{y} = 7.52 = \hat{\alpha}_0$$

$$\hat{\alpha}_1 = 2.91$$

$$\hat{\beta} = 2.59$$

The fitted model equation is then:

$$\hat{\mu}(x) = 2.91 + 2.59x$$

```
> rbind(y, x)
      [,1] [,2] [,3] [,4] [,5] [,6] [,7] [,8] [,9] [,10]
y 9.18 7.66 6.33 4.51 14.04 4.94 4.24 8.19 4.55 11.57
x 1.30 1.40 2.70 0.30 3.60 0.70 1.10 2.10 0.90 3.30
```

```
> model1 <- lm(y ~ x)
```

```
> model1
```

Call:

```
lm(formula = y ~ x)
```

Coefficients:

(Intercept)	x
2.911	2.590

Properties of these estimators

What do we know about these estimators?

They are all linear combinations of the Y_i ,

$$\hat{\alpha} = \sum_{i=1}^n \left(\frac{1}{n} \right) Y_i$$

$$\hat{\beta} = \sum_{i=1}^n \left(\frac{x_i - \bar{x}}{K} \right) Y_i$$

where $K = \sum_{i=1}^n (x_i - \bar{x})^2$.

This allows us to easily calculate means and variances.

Means?

$$\mathbb{E}(\hat{\alpha}_0) = \mathbb{E}(\bar{Y}) = \frac{1}{n} \sum_{i=1}^n \mathbb{E}(Y_i) = \frac{1}{n} \sum_{i=1}^n [\alpha_0 + \beta(x_i - \bar{x})] = \alpha_0$$

Assignment Project Exam Help

$$\begin{aligned} \mathbb{E}(\hat{\beta}) &= \sum_{i=1}^n \frac{(x_i - \bar{x})}{K} \mathbb{E}(Y_i) = \frac{1}{K} \sum_{i=1}^n (x_i - \bar{x})(\alpha_0 + (x_i - \bar{x})\beta) \\ &= \frac{1}{K} \sum_{i=1}^n (x_i - \bar{x})\alpha_0 + \frac{K}{K}\beta = \beta \end{aligned}$$

Add WeChat powcoder

This also implies, $\mathbb{E}(\hat{\alpha}) = \alpha$ and $\mathbb{E}(\hat{\mu}(x)) = \mu(x)$, and so we have that all of the estimators are unbiased.

Variances?

$$\text{var}(\hat{\alpha}_0) = \text{var}(\bar{Y}) = \frac{1}{n^2} \sum_{i=1}^n \text{var}(Y_i) = \frac{\sigma^2}{n}$$

Assignment Project Exam Help

$$\begin{aligned} \text{var}(\hat{\beta}) &= \text{var}\left(\sum_{i=1}^n \frac{(x_i - \bar{x})}{K} Y_i\right) = \sum_{i=1}^n \left(\frac{x_i - \bar{x}}{K}\right)^2 \text{var}(Y_i) \\ &= \frac{1}{K^2} \sum_{i=1}^n (x_i - \bar{x})^2 \text{var}(Y_i) = \frac{1}{K^2} \sum_{i=1}^n (x_i - \bar{x})^2 \sigma^2 \end{aligned}$$

$$= \frac{1}{K^2} \sigma^2 \sum_{i=1}^n (x_i - \bar{x})^2 = \frac{1}{K^2} \sigma^2 K$$

$$= \frac{\sigma^2}{K}$$

Similarly,

$$\text{var}(\hat{\alpha}) = \left(\frac{1}{n} + \frac{\bar{x}^2}{K} \right) \sigma^2$$

$$\text{cov}(\hat{\alpha}, \hat{\beta}) = 0$$

$$\text{var}(\hat{\mu}(x)) = \left(\frac{1}{n} + \frac{(x - \bar{x})^2}{K} \right) \sigma^2$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Can we get their standard errors?

We need an estimate of σ^2 .

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Analysis of variance: iid model

For $X_i \sim N(\mu, \sigma^2)$ iid,

Assignment Project Exam Help

$$\sum_{i=1}^n (X_i - \mu)^2 = \sum_{i=1}^n (X_i - \bar{X})^2 + n(\bar{X} - \mu)^2$$

<https://powcoder.com>

Add WeChat powcoder

Analysis of variance: regression model

Assignment Project Exam Help

$$\sum_{i=1}^n (Y_i - \alpha_0 - \beta(x_i - \bar{x}))^2$$

$$= \sum_{i=1}^n (Y_i - \alpha_0 - \beta(x_i - \bar{x}) + \hat{\alpha}_0 + \hat{\beta}(x_i - \bar{x}) - \alpha_0 - \beta(x_i - \bar{x}))^2$$

$$= \sum_{i=1}^n (Y_i - \hat{\alpha}_0 - \hat{\beta}(x_i - \bar{x}) + (\hat{\alpha}_0 - \alpha_0) + (\hat{\beta} - \beta)(x_i - \bar{x}))^2$$

$$= \sum_{i=1}^n (Y_i - \hat{\alpha}_0 - \hat{\beta}(x_i - \bar{x}))^2 + n(\hat{\alpha}_0 - \alpha_0)^2 + K(\hat{\beta} - \beta)^2$$

Note that the cross-terms disappear. Let's see...

The cross-terms...

Assignment Project Exam Help

$$t_1 = 2 \sum_{i=1}^n (Y_i - \hat{\alpha}_0 - \hat{\beta}(x_i - \bar{x}))(\hat{\alpha}_0 - \alpha_0)$$

$$t_2 = 2 \sum_{i=1}^n (Y_i - \hat{\alpha}_0 - \hat{\beta}(x_i - \bar{x}))(\hat{\beta} - \beta)(x_i - \bar{x})$$

$$t_3 = 2 \sum_{i=1}^n (x_i - \bar{x})(\hat{\beta} - \beta)(\hat{\alpha}_0 - \alpha_0)$$

Since $\sum_{i=1}^n (x_i - \bar{x}) = 0$ and $\sum_{i=1}^n (Y_i - \hat{\alpha}_0) = \sum_{i=1}^n (Y_i - \bar{Y}) = 0$,
the first and third cross-terms are easily shown to be zero.

For the second term,

$$\frac{t_2}{2(\hat{\beta} - \beta)} = \sum_{i=1}^n (Y_i - \bar{Y})(x_i - \bar{x}) - \hat{\beta} \sum_{i=1}^n (x_i - \bar{x})^2$$

$$= \sum_{i=1}^n (Y_i - \bar{Y})(x_i - \bar{x}) - \hat{\beta}K$$

$$= \sum_{i=1}^n Y_i(x_i - \bar{x}) - \sum_{i=1}^n \bar{Y}(x_i - \bar{x})$$

$$= 0$$

Therefore, all the cross-terms are zero.

Back to the analysis of variance formula. . .

Assignment Project Exam Help

$$\sum_{i=1}^n (Y_i - \alpha_0 - \beta(x_i - \bar{x}))^2$$

$$= \sum_{i=1}^n (Y_i - \hat{\alpha}_0 - \hat{\beta}(x_i - \bar{x}))^2 + n(\hat{\alpha}_0 - \alpha_0)^2 + n(\hat{\beta} - \beta)^2$$

Taking expectations gives,

$$\sigma^2 = \mathbb{E}(D^2) + n\sigma^2 + n\sigma^2$$
$$\Rightarrow \mathbb{E}(D^2) = (n-2)\sigma^2$$

where

$$D^2 = \sum_{i=1}^n (Y_i - \hat{\alpha}_0 - \hat{\beta}(x_i - \bar{x}))^2.$$

Variance estimator

Based on these results, we have an unbiased estimator of the variance,

$$\hat{\sigma}^2 = \frac{1}{n-2} D^2.$$

The inferred mean for each observation is called its fitted value,
 $\hat{Y}_i = \hat{\alpha}_0 + \hat{\beta}(x_i - \bar{x}).$

The deviation from each fitted value is called a residual, $R_i = Y_i - \hat{Y}_i.$

The variance estimator is based on the sum of squared residuals,
 $D^2 = \sum_{i=1}^n R_i^2.$

Example: variance estimate

For our data:

$$\hat{d}^2 = 16.12$$

$$\hat{\sigma}^2 = 2.015$$

$$\hat{\sigma} = 1.42$$

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Standard errors

We can substitute $\hat{\sigma}^2$ into the formulae for the standard deviation of the estimators in order to calculate standard errors

For example,

$$\text{var}(\hat{\beta}) = \frac{\hat{\sigma}^2}{K}$$

$$\Rightarrow \text{se}(\hat{\beta}) = \frac{\hat{\sigma}}{\sqrt{K}}$$

Add WeChat powcoder

Example: standard errors

For our data:

$$se(\hat{\alpha}_0) = \frac{\hat{\sigma}}{\sqrt{n}} = 0.449$$

$$se(\hat{\beta}) = \frac{\hat{\sigma}}{\sqrt{K}} = 0.404$$

$$se(\hat{\mu}(x)) = \hat{\sigma} \sqrt{\frac{1}{n} + \frac{(x - \bar{x})^2}{K}} = 1.42 \times \sqrt{\frac{1}{10} + \frac{(x - 1.78)^2}{1234}}$$

Add WeChat powcoder

Maximum likelihood estimation

Want to also construct confidence intervals. This requires further assumptions about the population distribution

Assignment Project Exam Help

Let's assume a normal distribution:

$$Y_i \sim N(\alpha + \beta x_i, \sigma^2)$$

<https://powcoder.com>

Alternative notation (commonly used for regression/linear models):

$$Y_i = \alpha + \beta x_i + \epsilon_i, \quad \text{where } \epsilon_i \sim N(0, \sigma^2).$$

Add WeChat powcoder

Let's maximise the likelihood...

Since the Y_i 's are independent, the likelihood is:

$$L(\alpha, \beta, \sigma^2) = \prod_{i=1}^n \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left\{ -\frac{(y_i - \alpha - \beta x_i)^2}{2\sigma^2} \right\}$$
$$= \left(\frac{1}{\sqrt{2\pi\sigma^2}} \right)^n \exp \left\{ -\frac{\sum_{i=1}^n (y_i - \alpha - \beta x_i)^2}{2\sigma^2} \right\}$$

$$-\ln L(\alpha, \beta, \sigma^2) = \frac{n}{2} \ln(2\pi\sigma^2) + \frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - \alpha - \beta x_i)^2$$
$$= \frac{n}{2} \ln(2\pi\sigma^2) + \frac{1}{2\sigma^2} H(\alpha, \beta)$$

The α_0 and β that maximise the likelihood (minimise the log-likelihood) are the same as those that minimise the sum of squares, H .

The OLS estimates are the same as the MLEs!

What about σ^2 ?

Differentiate by σ , set to zero, solve...

Assignment Project Exam Help
 $\hat{\sigma}_{MLE}^2 = \frac{1}{n} D^2$

This is biased. Prefer to use the previous, unbiased estimator,

<https://powcoder.com>
 $\hat{\sigma}^2 = \frac{1}{n-2} D^2$

Add WeChat powcoder

Sampling distributions

The Y_1, \dots, Y_n are independent normally distributed random variables.

Except for $\hat{\sigma}^2$, our estimators are linear combinations of the Y_i so will also have normal distributions, with mean and variance as previously derived

For example,

$$\hat{\beta} \sim N\left(\beta, \frac{\sigma^2}{K}\right)$$

Moreover, we know $\hat{\alpha}_0$ and $\hat{\beta}$ are independent, because they are bivariate normal rvs with zero covariance.

Using the analysis of variance decomposition (from earlier), we can show that,

$$\frac{(n-2)\hat{\sigma}^2}{\sigma^2} \sim \chi_{n-2}^2.$$

Assignment Project Exam Help

Therefore, we can define pivots for the various mean parameters. For example,

$$\frac{\hat{\beta} - \beta}{\sigma / \sqrt{K}} \sim t_{n-2}$$

and

$$\frac{\hat{\mu}(x) - \mu(x)}{\sigma \sqrt{\frac{1}{n} + \frac{(x-\bar{x})^2}{K}}} \sim t_{n-2}$$

This allows us to construct confidence intervals.

Example: confidence intervals

For our data, a 95% CI for β is:

$$\hat{\beta} \pm c \frac{\sigma}{\sqrt{K}} = 2.59 \pm 2.31 \times 0.404 = (1.66, 3.52)$$

where c is the 0.975 quantile of t_{n-2}

A 95% CI for $\mu(3)$ is:

$$\hat{\mu}(3) \pm c \times \text{se}(\hat{\mu}(3)) = 10.68 \pm 2.31 \times 0.667 = (9.14, 12.22)$$

Deriving prediction intervals

Use the same trick as we used for the simple model,

$$Y^* \sim N(\mu(x^*), \sigma^2)$$

$$\hat{\mu}(x^*) \sim N\left(\mu(x^*), \left(\frac{1}{n} + \frac{(x^* - \bar{x})^2}{K}\right) \sigma^2\right)$$

$$Y^* - \hat{\mu}(x^*) \sim N\left(0, \left(1 + \frac{1}{n} + \frac{(x^* - \bar{x})^2}{K}\right) \sigma^2\right)$$

Add WeChat powcoder

A 95% PI for Y^* is given by:

$$\hat{\mu}(x^*) \pm c \hat{\sigma} \sqrt{1 + \frac{1}{n} + \frac{(x^* - \bar{x})^2}{K}}$$

Example: prediction interval

A 95% PI for Y^* corresponding to $x^* = 3$ is:

Assignment Project Exam Help

$$10.68 \pm 2.31 \times 1.42 \times \sqrt{1 + \frac{1}{10} + \frac{(3 - 1.78)^2}{12.34}} = (7.06, 14.30)$$

<https://powcoder.com>

Much wider than the corresponding CI, as we've seen previously.

Add WeChat powcoder

```
> model1 <- lm(y ~ x)
> summary(model1)
```

Call:

```
lm(formula = y ~ x,
```

Residuals:

	Min	1Q	Median	3Q	Max
	-2.01970	-1.05963	0.02808	1.04774	1.80580

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	2.9114	0.8479	3.434	0.008908 **
x	2.5897	0.4041	6.408	0.000107 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.419 on 8 degrees of freedom

Multiple R-squared: 0.8369, Adjusted R-squared: 0.8166

F-statistic: 41.06 on 1 and 8 DF, p-value: 0.0002074

```
> # Confidence intervals for mean parameters
```

```
> confint(model1)
```

```
2.5 %    97.5 %
```

```
(Intercept) 0.9560629 4.867703
```

```
x1 1.6577220 3.521621
```

```
> # Data to use for prediction.
```

```
> data2 <- data.frame(x = 3)
```

```
> # Confidence interval for mu(3).
```

```
> predict(model1, newdata = data2, interval = "confidence")
```

```
fit      lwr      upr
```

```
1 10.6804 9.142823 11.21798
```

```
> # Prediction interval for y when x = 3.
```

```
> predict(model1, newdata = data2, interval = "prediction")
```

```
fit      lwr      upr
```

```
1 10.6804 7.064 14.2968
```

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

R example explained

- The `lm` (linear model) command fits the model.
- `model1` is an object that contains all the results of the regression needed for later calculations.
- `summary(model1)` acts on `model1` and summarizes the regression.
- `predict(car, calculate CIs and PIs).`
- R provides more detail than we need at the moment. Much of the output relates to hypothesis testing that we will get to later.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Plot data and fitted model

Assignment Project Exam Help

```
> plot(x, y, col = "blue")  
> abline(model1, col = "blue")
```

The command `abline(model1)` adds the fitted line to a plot.


<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



Fitted values and CIs for their means

```
predict(model, interval = "confidence")  
      fit      lwr      upr
```

```
1  7.572793  6.537531  8.608056  
2  6.536924  5.442924  7.630925  
3  8.349695  7.272493  9.426895  
4  3.688288  1.963799  5.412771  
5 12.234204 10.247160 14.221248  
6  4.724154  3.280382  6.167925  
7  5.780023  4.546338  6.973707  
8  8.349695  7.272496  9.426895  
9  5.242088  3.921478  6.562699  
10 12.752138 10.603796 14.900481
```

Confidence band for the mean

```
data3 <- data.frame(x = seq(-1, 5, 0.05))  
> y.conf <- predict(model1, data3, interval = "confidence")  
> head(cbind(data3, y.conf))
```


	x	fit	lwr	upr
1	-1.00	0.3217104	-2.468233	3.111653
2	-0.95	0.4811941	-2.295531	3.197919
3	-0.90	0.5806777	-2.122943	3.284298
4	-0.85	0.7101613	-1.950472	3.370794
5	-0.80	0.8396449	-1.778124	3.457414
6	-0.75	0.9691286	-1.605506	3.544164

```
> matplot(data3$x, y.conf, type = "l", lty = c(1, 2, 2),  
+         lwd = 2, xlab = "x", ylab = "y")  
> points(x, y, col = "blue")
```

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



Prediction bands for new observations

Assignment Project Exam Help

```
y.pred <- predict(model1, data3, interval = "prediction")  
> head(cbind(data3, y.pred))
```


	x	fit	lwr	upr
1	-1.00	0.3217104	-3.979218	4.622639
2	-0.95	0.5119411	-3.831827	4.724215
3	-0.90	0.5806777	-3.664763	4.826119
4	-0.85	0.7101613	-3.508034	4.928357
5	-0.80	0.8396449	-3.351646	5.030936
6	-0.75	0.9491286	-3.195606	5.133863

```
> matplot(data3$x, y.pred, type = "l", lty = c(1, 3, 3),  
+         lwd = 2, xlab = "x", ylab = "y")  
> points(x, y, col = "blue")
```

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



Both bands plotted together

```
> matplot(data3$x, cbind(y.conf, y.pred[, 1-4]),  
+         type = "l", lwd = 2, lty = c(1, 2, 2, 3, 3),  
+         xlab = "x", ylab = "y")  
> points(x, y)
```


<https://powcoder.com>

Add WeChat powcoder

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



Checking our assumptions

What modelling assumptions have we made?

- Linear model for the mean
- Equal variances for all observations (**homoscedasticity**)
- Normally distributed residuals

Ways to check these:

- Plot the data and fitted model together (done!)
- Plot residuals vs fitted values
- QQ plot of the residuals



In R, the last two of these are very easy to do:

```
> plot(model1, 1:2)
```


Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



Outline

Introduction

Regression

Assignment Project Exam Help

Point estimation of the mean

Interlude: Analysis of variance

Point estimation of the variance

Standard errors of the estimates

Confidence intervals

Prediction intervals

R example

Model checking

Further regression models

Correlation

Definitions

Point estimation

Relationship to regression

<https://powcoder.com>

Add WeChat powcoder

Multiple regression

- What if we have more than one predictor?

- Observe x_{i1}, \dots, x_{ik} as well as y_i (for each i)

- Can fit a **multiple regression model**:

$$\mathbb{E}(Y \mid x_1, \dots, x_k) = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \dots + \beta_k x_k$$

- This is linear in the coefficients, so is still a linear model

- Fit by method of least squares by minimising:

$$II = \sum_{i=1}^n (y_i - \beta_0 - \beta_1 x_{i1} - \beta_2 x_{i2} - \dots - \beta_k x_{ik})^2$$

- Take partial derivatives, etc., and solve for β_0, \dots, β_k .
- The subject Linear Statistical Models (MAST30025) looks into these types of models in much more detail.

Two-sample problem

- The two-sample problem can be expressed as a linear model!
- Sample $Y_1, \dots, Y_n \sim N(\mu_1, \sigma^2)$ and $Y_{n+1}, \dots, Y_{n+m} \sim N(\mu_2, \sigma^2)$
- Define **indicator variables** (x_{i1}, x_{i2}) where $(x_{i1}, x_{i2}) = (1, 0)$ for $i = 1, \dots, n$ and $(x_{i1}, x_{i2}) = (0, 1)$ for $i = n + 1, \dots, n + m$.
- Observed data: (y_i, x_{i1}, x_{i2})
- Then Y_1, \dots, Y_n each have mean $1 \times \beta_1 + 0 \times \beta_2 = \mu_1$ and Y_{n+1}, \dots, Y_{n+m} each have mean $0 \times \beta_1 + 1 \times \beta_2 = \mu_2$.
- This is in the form a multiple regression model
- The **general linear model** unifies many different types of models together into a common framework. The subject MAST30025 covers this in more detail.

Outline

Introduction

Regression

Assignment Project Exam Help

Point estimation of the mean

Interlude: Analysis of variance

Point estimation of the variance

Standard errors of the estimates

Confidence intervals

Prediction intervals

R example

Model checking

Further regression models

Correlation

Definitions

Point estimation

Relationship to regression

<https://powcoder.com>

Add WeChat powcoder

Correlation coefficient

(Revision) for two rvs X and Y , the correlation coefficient, or simply the correlation, is defined as:

$$\rho = \rho_{XY} = \frac{\text{cov}(X, Y)}{\sqrt{\text{var } X \text{ var } Y}} = \frac{\sigma_{XY}}{\sigma_X \sigma_Y}$$

This is a quantitative measure of the strength of relationship, or association, between X and Y .










We will now consider inference on ρ based on an iid sample of pairs (X_i, Y_i) .

Note: unlike in regression, X is now considered as a random variable.

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder



Sample covariance

To estimate $\text{cov}(X, Y)$ we use the sample covariance:

$$S_{XY} = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y}) = \frac{1}{n-1} \left(\sum_{i=1}^n X_i Y_i - n\bar{X}\bar{Y} \right)$$

<https://powcoder.com>

You can check that this is unbiased, $\mathbb{E}(S_{XY}) = \sigma_{XY} = \text{cov}(X, Y)$.

Add WeChat powcoder

Sample correlation coefficient

To estimate ρ we use the sample correlation coefficient (also known as Pearson's correlation coefficient)

$$R = R_{XY} = \frac{S_{XY}}{S_X S_Y} = \frac{\sum_{i=1}^n (X_i - \bar{X})(Y_i - \bar{Y})}{\sqrt{\sum_{i=1}^n (X_i - \bar{X})^2 \sum_{i=1}^n (Y_i - \bar{Y})^2}}$$

You can check that $|R| \leq 1$, just like $|\rho| \leq 1$.

This gives a point estimate of ρ .

For further results, we make some more assumptions. . .

Bivariate normal

Assume X and Y have correlation ρ and follow a bivariate normal distribution

$$\begin{bmatrix} X \\ Y \end{bmatrix} \sim N_2 \left(\begin{bmatrix} \mu_X \\ \mu_Y \end{bmatrix}, \begin{bmatrix} \sigma_X^2 & \rho\sigma_X\sigma_Y \\ \rho\sigma_X\sigma_Y & \sigma_Y^2 \end{bmatrix} \right)$$


<https://powcoder.com>

In this case, the regressions are linear,

$$\mathbb{E}(X | Y = y) = \mu_X + \frac{\rho\sigma_X}{\sigma_Y}(y - \mu_Y) = \alpha' + \beta'y$$

$$\mathbb{E}(Y | X = x) = \mu_Y + \frac{\rho\sigma_Y}{\sigma_X}(x - \mu_X) = \alpha + \beta x$$

Note: $\beta' \neq 1/\beta$



Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder

Variance explained

An alternative analysis of variance decomposition:

$$\begin{aligned}\sum (Y_i - \bar{Y})^2 &= \sum (Y_i - \hat{\alpha} - \hat{\beta}x_i)^2 + \hat{\beta}^2 \sum (x_i - \bar{x})^2 \\ &= (1 - R^2) \sum (Y_i - \bar{Y})^2 + R^2 \sum (Y_i - \bar{Y})^2\end{aligned}$$

This implies that R^2 is the proportion of the variation in Y 'explained' by x .

In this usage R^2 is called the coefficient of determination.

Remarks

- For simple linear regression, the coefficient of determination is the same as the square of the sample correlation, with both being denoted by R^2 .
- Also, the proportion of Y explained by x is the same as the proportion of X explained by y . Both are equal to R^2 , which is a symmetric expression of both X and Y .
- For more complex models, the coefficient of determination is more complicated: it needs to be calculated using all predictor variables together.

Approximate sampling distribution

Define:

Assignment Project Exam Help

$$g(r) = \frac{1}{2} \ln \left(\frac{1+r}{1-r} \right)$$

This function has a standard name, $g(r) = \text{artanh}(r)$, and so does its inverse, $g^{-1}(r) = \tanh(r)$. The function $g(r)$ is also known as the Fisher transformation.

<https://powcoder.com>

The following is a widely used approximation:

Add WeChat powcoder

$$g(R) \approx N \left(g(\rho), \frac{1}{n-3} \right)$$

We can use this to construct approximate confidence intervals.

Example: correlation

For our data:

$$r = 0.91$$

$$r^2 = 0.84$$

An approximate 95% CI for $g(\rho)$ is:

$$g(r) \pm \frac{c}{\sqrt{n-3}} = 1.56 \pm 1.96 \times 0.378 = (0.819, 2.30)$$

where $c = \Phi^{-1}(1 - \alpha/2)$. Transforming this to an approximate 95% CI for ρ :

$$(\tanh(0.819), \tanh(2.30)) = (0.67, 0.98)$$

```
> cor(x, y)
[1] 0.9148421
```

```
> cor(x, y)^2
[1] 0.836936
```

```
> cor.test(x, y)
```

Pearson's product-moment correlation

data: x and y

t = 6.4078, df = 8, p-value = 0.0002074

alternative hypothesis: true correlation is not equal to 0

95 percent confidence interval:

0.6726924 0.9799873

sample estimates:

cor

0.9148421

Assignment Project Exam Help

<https://powcoder.com>

Add WeChat powcoder


```
> model1 <- lm(y ~ x)
> summary(model1)
```

Call:

```
lm(formula = y ~ x,
```

Residuals:

Min	1Q	Median	3Q	Max
-2.01970	-1.05963	0.02808	1.04774	1.80580

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	2.9114	0.8479	3.434	0.008908 **
x	2.5897	0.4041	6.408	0.000107 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 1.419 on 8 degrees of freedom

Multiple R-squared: 0.8369, Adjusted R-squared: 0.8166

F-statistic: 41.06 on 1 and 8 DF, p-value: 0.0002074