

## Lecture 12 Stationary Distribution (cont)

Example 1. Let  $\{X_n: n=0,1,2,\dots\}$  be a Markov chain with  $S=\{0,1,2,3\}$  and

$$P = \begin{pmatrix} \frac{1}{2} & \frac{1}{2} & 0 & 0 \\ \frac{1}{6} & \frac{1}{2} & \frac{1}{3} & 0 \\ 0 & \frac{1}{3} & \frac{1}{2} & \frac{1}{6} \\ 0 & 0 & \frac{1}{2} & \frac{1}{2} \end{pmatrix}$$

Find the stationary distribution of the chain.

Assignment Project Exam Help

Solution: this is a birth-death

Markov chain. Visit <https://powcoder.com> Add WeChat powcoder

$$d=3 \quad p_0 = \frac{1}{2}, p_1 = \frac{1}{3}, p_2 = \frac{1}{6} \\ \varepsilon_1 = \frac{1}{6}, \varepsilon_2 = \frac{1}{3}, \varepsilon_3 = \frac{1}{2}$$

$$\Rightarrow \pi(k) = \frac{\frac{p_0 \dots p_{k-1}}{\varepsilon_1 \dots \varepsilon_k}}{1 + \frac{p_0}{\varepsilon_1} + \frac{p_0 p_1}{\varepsilon_1 \varepsilon_2} + \frac{p_0 p_1 p_2}{\varepsilon_1 \varepsilon_2 \varepsilon_3}}$$

$$\Rightarrow 1 + \frac{p_0}{\varepsilon_1} + \frac{p_0 p_1}{\varepsilon_1 \varepsilon_2} + \frac{p_0 p_1 p_2}{\varepsilon_1 \varepsilon_2 \varepsilon_3} \\ = 1 + 3 + 3 + 1 = 8$$

$$\Rightarrow \pi(0) = \frac{1}{8}, \quad \pi(1) = \frac{3}{8}, \quad \pi(2) = \frac{3}{8} \\ \pi(3) = \frac{1}{8}$$

Example 2. Suppose we have two boxes containing  $d$  red balls and  $d$  blue balls in total. Initially  $d$  balls are placed in box I. At each trial one ball is chosen at random from each box and then the two balls are put back in the opposite boxes. Set

$X_0$  = number of red balls in box I initially

$X_n$  = number of red balls in box I after the  $n^{\text{th}}$  trial

Find ① IP ;

② the stationary distribution

Solution:  $S = \{0, 1, 2, \dots, d\}$

$$\begin{aligned} P(x, x) &= \frac{x}{d} \cdot \frac{d-x}{d} + \frac{d-x}{d} \cdot \frac{x}{d} \\ &= 2 \cdot \frac{x(d-x)}{d^2} \end{aligned}$$

$$P(x, x+1) = \frac{d-x}{d} \cdot \frac{d-x}{d} = \left(\frac{d-x}{d}\right)^2$$

$$P(x, x-1) = \frac{x}{d} \cdot \frac{x}{d} = \left(\frac{x}{d}\right)^2$$

Assignment Project Exam Help

$$P(x, y) = 0 \text{ for } y \neq x, x \pm 1$$

<https://powcoder.com>

$\Rightarrow \{X_n : n=0, 1, 2, \dots\}$  is a birth-death

Markov chain.

$$P = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 & 0 \\ \left(\frac{1}{d}\right)^2 & 2\frac{d-1}{d^2} & \left(\frac{d-1}{d}\right)^2 & \dots & \dots & \dots \\ 0 & 0 & \dots & \dots & 1 & 0 \end{pmatrix}$$

② For any  $k = 1, \dots, d$ ,

$$\frac{p_0 \dots p_{k-1}}{\sum_1 \dots \sum_k} = \frac{\left(\frac{d}{a}\right)^2 \left(\frac{d-1}{a}\right)^2 \dots \left(\frac{d-k+1}{a}\right)^2}{\left(\frac{1}{a}\right)^2 \dots \left(\frac{k}{a}\right)^2}$$

$$= \left( \frac{d(d-1)(d-2) \dots (d-k+1)}{1 \cdot 2 \dots k} \right)^2$$

$$= \binom{d}{k}^2$$

Assignment Project Exam Help

$$\Rightarrow \pi(k) = \frac{\binom{d}{k}}{1 + \binom{d}{1}^2 + \dots + \binom{d}{a}^2}$$

Add WeChat powcoder

$$= \frac{\binom{d}{k}}{\binom{2d}{a}}$$

$$1 + \binom{d}{1}^2 + \dots + \binom{d}{a}^2 = \sum_{i=0}^d \binom{d}{i} \binom{d}{a-i}$$

$$\binom{2d}{a} //$$

For birth-death chain with

$S = \{0, 1, 2, \dots\}$  the stationary distribution exists if and only if

$$1 + \sum_{k=1}^{\infty} \frac{p_0 \cdots p_{k-1}}{q_1 \cdots q_k} < \infty$$

If the above holds, then

$$\pi(k) = \frac{p_0 \cdots p_{k-1}}{q_1 \cdots q_k} \cdot \frac{1}{1 + \sum_{i=1}^{\infty} \frac{p_0 \cdots p_{i-1}}{q_1 \cdots q_i}}$$

Assignment Project Exam Help  
<https://powcoder.com>  
 Add WeChat powcoder

- Number and Average Number of Visit to a State

For any  $n \geq 1$ ,  $y \in S$ , set

$$N_n(y) = \sum_{k=1}^n \mathbb{1}_y(X_k)$$

$$G_n(x, y) = E_x[N_n(y)] = \sum_{k=1}^n P^k(x, y)$$

Theorem 1:

① For any transient state  $y$

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = 0 \quad \text{almost surely}$$

$$\lim_{n \rightarrow \infty} \frac{G_n(x, y)}{n} = 0$$

② For any recurrent state  $y$

Assignment Project Exam Help

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{1}{\mathbb{E}_y[T_y]} \quad \{T_y < \infty\}$$

<https://powcoder.com>

$$\lim_{n \rightarrow \infty} \frac{G_n(x, y)}{n} = \frac{P_{xy}}{m_y}$$

Add WeChat powcoder

Where  $m_y = \mathbb{E}_y[T_y]$ .

Proof: ① By Theorem 1 of Lecture 6,

$N_n(y) \rightarrow N(y) < \infty$  with probability one

$\Rightarrow \frac{N_n(y)}{n} \rightarrow 0$  with probability one

$$\frac{G_n(x, y)}{n} = \frac{E_x[N_n(y)]}{n}$$

$$= E_x\left[\frac{N_n(y)}{n}\right]$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{G_n(x, y)}{n} = E_x\left[\lim_{n \rightarrow \infty} \frac{N_n(y)}{n}\right]$$

$$= 0$$

Dominated Convergence theorem

## Assignment Project Exam Help

② Assume that  $y$  is recurrent.

<https://powcoder.com>

Let  $T_y^0$  denote the hitting time of  $y$  and assume this occurs.

Add WeChat powcoder

$w_y^1$  = additional number of steps for the first return to  $y$

$w_y^r$  = additional number of steps for the  $r$ th return to  $y$

then  $w_y^1, \dots, w_y^r$  are i.i.d with mean  $m_y$

By the Strong law of large numbers,

$$\frac{w_y^1 + \dots + w_y^r}{r} \xrightarrow{r \rightarrow \infty} m_y \text{ with}$$

probability one.

Let  $T_y^k$  denote the time of the  $k^{\text{th}}$  visit to  $y$ . Then

Assignment Project Exam Help

$$\frac{T_y^k}{k} = \frac{T_y^0 + w_y^1 + \dots + w_y^k}{k} \xrightarrow{k \rightarrow \infty} m_y$$

<https://powcoder.com>

with probability one

Add WeChat powcoder

Choosing  $k = N_n(y)$ , it follows that

$$T_y^{N_n(y)} \leq n \leq T_y^{N_n(y)+1}$$

$$\frac{N_n(y)^{\text{th visit}}}{n} \leq \frac{n}{N_n(y)+1} \leq \frac{T_y^{N_n(y)+1}}{N_n(y)+1}$$

$$\Rightarrow \frac{T_y^{N_n(y)}}{N_n(y)} \leq \frac{n}{N_n(y)} \leq \frac{T_y^{N_n(y)+1}}{N_n(y)+1}$$

$$= \frac{N_n(y)+1}{N_n(y)} \frac{T_y^{N_n(y)+1}}{N_n(y)+1}$$



$$\Rightarrow \lim_{n \rightarrow \infty} \frac{T_y^{N_n(y)}}{N_n(y)} \leq \lim_{n \rightarrow \infty} \frac{n}{N_n(y)} \leq \lim_{n \rightarrow \infty} \frac{T_y^{N_n(y)+1}}{N_n(y)+1}$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{n}{N_n(y)} = m_y$$

$$\Rightarrow \lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{1}{m_y}$$

Assignment Project Exam Help

If  $y$  is not accessible, then

<https://powcoder.com>

$N_n(y) = 0$ . Hence

Add WeChat powcoder

$$\lim_{n \rightarrow \infty} \frac{N_n(y)}{n} = \frac{\mathbb{1}_{\{T_y < \infty\}}}{m_y}$$

$$\lim_{n \rightarrow \infty} \frac{G_n(x, y)}{n} = \lim_{n \rightarrow \infty} E_x \left[ \frac{N_n(y)}{n} \right]$$

$$= E \left[ \lim_{n \rightarrow \infty} \frac{N_n(y)}{n} \right]$$

$$= \frac{P_{xy}}{m_y}$$

## Null Recurrent and Positive Recurrent States

Definition 1. For any recurrent state  $x \in S$ , let  $m_x = E_x[T_x]$

$x$  is null recurrent if  $m_x = \infty$

positive recurrent if  $m_x < \infty$

## Assignment Project Exam Help

Observation:

<https://powcoder.com>

$\lim_{n \rightarrow \infty} \frac{G_n(x, y)}{n} = 0$  if

Add WeChat powcoder

$y$  is transient or null recurrent.

Thus Null Recurrent is between  
Transient and Positive Recurrent.