spinal accessory and hyoglossal; but physiologists are not in general agreed upon the classification and names of the cranial nerves.

The nerves are mostly cells, or tubules, composed of membranous sacs or sheaths, lined with nervous matter, inclosing nerve-filaments in the form of bundles, which stretch away, divide, and subdivide, but never lose the original quality of sensation or motion which they derive from their roots. In every membranous sheath there are two distinct sets of filaments, one of which communicates with the cerebral hemispheres, and the other with the spinal cord.

The spinal cord is a long, irregular column of nerve-substance, inclosed in a sheath, composed of the cranial membranes prolonged down the spine. It extends from the medulla oblongata to the second lumbar vertebra of the spine. (See Fig. 4.)

The spinal cord, although composed of gray and white nervous matter like the brain, presents a different arrangement of the layers,—the white matter being on the exterior, and the gray or cineritious mass forming a band which traverses the interior of the column.

Like the medulla oblongata, the spinal cord is divided into two symmetrical halves, united in the middle by a commissure, or converging lines of nerve-fibre; it is also separated into an anterior and posterior column by a vertical fissure. Each lateral half is traversed by two longitudinal tracts, which separate it by distinct furrows. Thus there are on either side three well-marked columns, called the anterior, posterior, and lateral.