mencing at the connection with the uterus, it gradually widens and terminates in a bell-like orifice with a series of fringe-like processes termed fimbriæ, one of which is connected with the outer end of the ovary, as may be seen in Fig. 12, No. 1.

The ovaries are oval-shaped, elongated bodies, one on each side of the uterus, in the back part of the broad ligament, under and back of the Fallopian tubes.

The ovaries are connected to the uterus by ligaments, associated with their inner extremity; and at the outer extremity, to the above-mentioned fimbriated extremity of the Fallopian tubes by a short cord or ligament; in the posterior part to the broad ligament.

Each of the ovaries is about an inch and a half in length, and three-quarters of an inch in width, and about one-third of an inch thick—their weight from one to two drachms. Imbedded in the tissue of the internal surface of the ovaries are the Graafian vesicles, containing the ova. The fluid contained in these vesicles is albuminous, and holds the ovum. The Graafian vesicles gradually moving, work out to the surface of the ovaries, burst, and liberate the fluid and ova, passing into the Fallopian tubes; the fimbriated extremities are supposed to grasp the ovaries, and hence the mature ova are conducted through the tube into the Fallopian tubes, and conveyed to the uterus, and at regular periods are eliminated from the uterus, producing menstruation, unless the ovum is fecundated. (Fig. 12, No. 2.)

The two round ligaments of the uterus are two