ward in C, No. 6 electrode across the stomach in E. If a decided improvement is noted continue for twenty minutes. If, however, no perceptible effect is produced in five minutes, apply B–D twenty minutes.

Second. No. 5 electrode, base of brain, on spine downward in E, No. 7 electrode under the feet in H, twenty minutes.

Both of these applications may be completed by placing the round electrode between the hands in C, having the other electrode on the stomach or spine in E, five minutes. Apply C–G also in the same manner.

CONGESTIVE CHILLS.

First. No. 5 electrode, base of brain, on spine downward in B, No. 6 electrode across abdomen in D, until improvement in the congestion; then use C–E for B–D until normal temperature is restored. (This treatment may be continued for an hour.)

Second. If the congestion is cephalic, or of the brain, with chill, apply No. 5 electrode, base of brain, on spine downward in B, No. 7 electrode under the feet in D, twenty to thirty minutes. Use C–G also.

Third. If the congestion is pulmonic, or of the lungs, round electrode in hot foot-bath in D, No. 5 electrode from base of brain, on spine downward in B, ten minutes; change No. 5 electrode, same position, to C; place No. 6 electrode across upper abdomen in E, ten to fifteen minutes.

Fourth. If the congestion is of the kidneys, place No. 5 electrode from coccyx upward on the spine in