hemorrhage, etc., as for the same maladies otherwise caused. The vapor-bath is to be guardedly employed, if needed, remembering the debility existing in these cases, and given with a mild tonic current (F–G), not to exceed ten minutes.

Ulceration of the bowels in fevers.

Apply No. 6 electrode across abdomen, just covering the navel, in B, No. 5 electrode on the spine from coccyx up in D, fifteen minutes, then, for any soreness (see note), ten minutes. Repeat once or twice a day, meeting other symptoms, if needful, at any time during the second daily treatment, but aim in every manipulation to implicate the affected parts.

Hemorrhage of the bowels in fevers.

No. 2 electrode at the base of spine in B, sponge-cup or No. 4 electrode over the right iliac region (the flank enclosing the small intestines), in C, this being the supposed origin of the flow.

The principle of this application to be observed in all kinds of hemorrhage. B–E, reversed, and F–H, however, being potent in paretic conditions of the blood-vessels.

YELLOW FEVER.

Have apartment disinfected and as thoroughly ventilated as possible. Keep the patient warm. Sponge-cup rubbed over liver and gall-bladder to the stomach in B, No. 4 electrode between the kidneys in D, fifteen minutes; change immediately No. 4 electrode over stomach in D, No. 8 electrode on the back from coccyx, long point up, in E, ten minutes; also place feet in hot foot-bath, with round