NUMBNESS AND HEAVINESS OF THE LEGS.

No. 6 electrode over one hip, then the other, ten minutes each in B. No. 7 electrode under the feet in D, twenty minutes. Also use D–F in the same manner.

CEREBRO-SPINAL MENINGITIS—SPOTTED FEVER.

Treat for chill and fever as usual.

First. No. 5 electrode from base of brain down the spine in B, round electrode covered with sponge pressed against the coccyx in E, fifteen minutes.

Second. No. 2 electrode at base of brain in B. No. 5 electrode on spine, from coccyx extending up in C, twenty minutes; every second day, if practicable, administer vapor-bath—not too hot, or too long—with round electrode in D in the foot-bath, sponge-cup in A from base of brain down the spine.

SEVERE INFLAMMATIONS IN GENERAL.

Inflammation must be rapidly overcome by the use of the proper currents mildly administered, usually more or less complication may be expected; in which case the several symptoms must be successively or, if practicable, simultaneously met.

Thus, for example, we will suppose a bad case of dyspepsia, with hæmorrhoids, to be aggravated into a violent inflammatory condition. The typical course of treatment for such a case would be, as usual, to attack the most distressing part first. Proceed thus:

First. Introduce the rectal electrode well lubricated with glycerine or fine soap (as all similar