aeter exceedingly uncomfortable, and is thus prejudicial to speedy recovery. It may appear as local or general irritability, or may even be a product of active electrization, disturbing the functional equilibrium. In such cases use round electrode in A between the hands, No. 6 cleetrode over stomach in C, five to fifteen minutes. Any faulty electrization should of course be remedied, either by changing to the proper current or different locations.

If a child is cross and irritable, place No. 5 electrode on spine from the base of brain downwards in A, and No. 7 electrode under feet in B, ten minutes, change B to hands five minutes. Irritability in adult, A–C instead of the preceding current, five or ten minutes longer duration. In abdominal discomfort, No. 5 electrode on spine from waist up in D, No. 6 electrode aeross abdomen in B, ten minutes, in either ease end with A base of brain, C in hands five minutes.

### VERTIGO.

The cranial diagnosis will enable the physician to proceed with confidence. The cause of this trouble is frequently gastric derangement, or it may be due to eerebral or other nervous affections. If gastrie, treat as for dyspepsia.

If cerebral, place

First. No. 5 cleetrode on spine from base of brain downward in B, No. 6 electrode over stomach, liver, and spleen in D, ten to fifteen minutes, remove No. 6 electrode to hands five minutes, alternate with

Second. No. 5 electrode on spine from coccyx up-