First. The patient should lie upon the back. No. 5 electrode from the base of brain on the spine downward in C, No. 6 electrode under the calves of the legs in A, for ten minutes; then remove No. 5 electrode from the base of brain and place it across the stomach (to implicate the diaphragm) ten minutes.

Second. No. 5 electrode between the shoulders, on the spine, downward in H, No. 6 electrode across the lower abdomen in E, fifteen minutes; then remove No. 6 electrode in E to the feet for ten minutes.

The round electrodes held in the hands, in E–G, five minutes; can be used to complete either of the preceding applications.

Third. No. 6 electrode under the calves of legs in E, No. 5 electrode from coccyx up between the kidneys in H, ten minutes. Transfer No. 5 electrode to base of brain, on spine downward, in A, No. 6 electrode to under calves of legs in C, ten minutes. Transfer No. 6 electrode in C to under the hands five minutes.

Fourth. The vapor-bath, with No. 7 electrode placed under the towel on the seat in D, round electrode in the foot-bath in A, first having applied the sponge-cup down the spine in B in the usual way.

Two or three hours after the bath use No. 3 electrode between kidneys in E, No. 2 electrode on the upper part of the spine, below the back of neck, in G, fifteen minutes.

Repeat the direct applications for a child every day, for an adult twice a day, until better. Local affections may be treated in the interval.