applications must be regular, until complete restoration is effected.

Having already given the points for localized treatment, it remains only to give those for a whole limb or other region. If in the upper part of the body, the cheek, tongue, larynx, arm, or hand, apply along the paralyzed tract the negative with the brush, sponge-cup, spatula, or the operator's hand; place the positive upon the roots of the corresponding nerves, in the spine, graduating the current to the electro-sensibility and electro-contractility of the patient, using the SILVER HELIX, or A–C, E–F, and F–H currents, ten or fifteen minutes, or until some slight sensation or motion is perceptible, administering frequent but not very long applications of this kind, never over twenty minutes.

If the parts are deficient in animal heat, use C–E current from the base of brain to the throat, or the hands.

Paralysis of the lower extremities is to be managed on very similar principles, if no visceral paralysis, as of the bladder, rectum, or anus, is present. Often, however, it coexists, owing to lumbar or other injury.

For this apply No. 6 electrode or the brush along the perineum in F. The sponge-cup upon the nerve-roots in D, vibrating as directed, for fifteen or twenty minutes. Or, if the sphincters are particularly affected, introduce the rectal instrument into the rectum, or the rubber insulated bladder electrode to the neck of the bladder in A, No. 5 electrode on the spine between the kidneys in D, five or ten minutes' treatments, alternate days.