the hair downward in D. Sponge-cup in C, resting half a minute on bridge of the nose, and then stroked over the eyes outward, keeping the sponge under the arch of the brow, five minutes each eye. Conditional, on extension of the catarrh to the throat, complete the treatment with the sponge-cup stroked around the throat and downward over the breast-bone in B, No. 4 electrode small of back on the spine in D, ten minutes.

Fourth. Conditional, on the anterior passages being the principal seat of disease, use the nasal electrode therein, with E, No. 2 electrode base of brain in C, five or ten minutes.

Nasal polypus.

First. To induce a discharge use the bodkin similar to the nasal electrode with a sponge wrapped and tied around the end, applied within the nasal passage to the polypus in H, No. 2 electrode base of brain in C, ten minutes. Alternate with A–D current ten minutes in the same manner.

Second. After two or three applications a fine silver wire may be placed around the polypus in D, A at the base of brain ten minutes; the tumor will probably become detached during the second or third application; when it discharges, use No. 2 electrode base of brain in A, sponge-cup base of nose, between the eyes, in D, ten minutes.

Bleeding of the nose.

The round electrode or sponge-cup between the hands in A, No. 1 electrode at the base of brain in C. If not relieved in five minutes change to E-F current, holding the hands over the head.