apply through the operator, with the hand placed on the external enlargement in B, ten minutes.

Second. No. 2 electrode base of brain in A, sponge-cup externally over tonsils in C, five minutes each side.

Third. No. 2 electrode base of brain in B, the spatula on the tongue in D, ten minutes.

Fourth. No. 1 electrode back of neck in C, electrode between the hands in E, ten minutes.

If chronic,

First. Use the fourth treatment as for an acute attack, but continue twenty minutes. Repeat four or five times, or until an impression is made, which will certainly be marked by soreness, then treat for soreness. (See note.)

Second. No. 3 electrode between the shoulders on the spine in C, the sponge-cup stroking forward over one tonsil then the other in A, ten minutes each.

WHOOPING-COUGH.

No. 1 electrode base of brain in A, No. 4 electrode across stomach in C, fifteen minutes; then change No. 1 to throat in B, No. 4 or 6 across the liver, stomach, and spleen in D, ten minutes. Repeat at intervals.

PAROTIS—MUMPS.

First. No. 1 electrode base of the brain in B, No. 2 electrode or sponge-cup on swelling in D, ten minutes. If soreness, change No. 1 electrode to A and sponge-cup to C, five minutes more. If the soreness becomes extreme, apply sponge-cup to the swelling