in A, No. 3 electrode on the breast-bone in B, five minutes.

Second. General treatment. (See directions.)

Third. If metastasis occur to the brain, No. 2 electrode base of brain in A, No. 4 electrode across the lower part of back in D, ten minutes. Repeat.

Fourth. If to the mammary glands in the female, place No. 2 electrode on the opposite side of the breast from the swelling in C, the sponge-cup on the swelling in E, ten minutes. General treatment also. (See directions.)

Fifth. If to the testicles in the male, place the round electrode in a hot sitz-bath in D, the sponge-cup stroked down the spine from base of brain in A, ten minutes. Repeat three times daily.

Sixth. If gastric trouble supervenes, manipulate as for gastritis. (See directions.)

GOITRE.

First. No. 3 electrode base of brain or between the shoulders on spine in B, No. 2 electrode covered with a sponge upon the enlargement in D, ten to fifteen minutes. Alternate with C–D, treatments twice a day.

The size of the goitre will probably soon lessen, and thereupon soreness may occur; relieve with A–C current used in the same manner.

Second. No. 5 electrode from the nape of neck downward on the spine in B, sponge-cup moved about over the enlargement with a rocking or kneading motion in D, ten to fifteen minutes. Alternate with A–D current in the same manner.