INFLUENZA—COLD IN THE HEAD.

First. Place No. 1 electrode base of brain in A, No. 4 electrode across stomach in C, ten minutes.

Second. No. 5 electrode from base of brain, on the spine, downward in B, No. 6 electrode across the stomach, liver, and spleen in D, fifteen minutes.

Third. No. 1 electrode on throat in C, No. 2 electrode back of neck in E, ten minutes.

Manipulate as for nasal catarrh. (See directions.)

PHTHISIS—CONSUMPTION.

In all pulmonary diseases observe the general rule, viz., *never place the electrodes over the lungs or kidney;* confine the manipulations to the median line, front or back, or above or below the position of the lungs or the kidneys.

First. Place No. 5 electrode (bent to lie close) along the front of the throat, extending downward over the breast-bone in A, No. 4 electrode on the lower part of spine in F. B–D current may be used in the same manner, when the patient can endure it without pain.

Second. Manipulate as for bronchitis. Use the "running-off" and general treatments. (See directions.)

For the cough, place No. 6 electrode from the end of breast-bone downward in C, No. 5 electrode between the shoulders downward in E, ten minutes or longer. No. 1 electrode across the front of throat in A, No. 3 electrode on the back of neck from the hair downward in C, ten or fifteen minutes.