ing the patient on No. 7 electrode in D, the sponge-cup (used as directed in second treatment for pleurisy) stroked down and round both sides in A, fifteen or twenty minutes.

INDIGESTION.

Take daily treatments after meals.

First. No. 4 electrode across the stomach in D, No. 5 electrode between the kidneys in B, twenty-five minutes. Alternate with D–E and D–G in the same manner.

Second. No. 5 electrode on the middle of the spine in A, No. 6 electrode across the liver, stomach, and spleen, with a rocking motion,* in D, or the sponge-cup stroked around from both sides to the stomach in D, fifteen or twenty minutes. This treatment should be used in all cases when, after eating, an accumulation of gas causes an oppressive feeling, or a sensation of heaviness in the head or eyes. If accompanied by vertigo or dizziness, apply

Third. No. 6 electrode across the upper abdomen in E, No. 2 electrode at the base of brain in C, five minutes.

DYSPEPSIA.

This most common disease is frequently accompanied by urinary trouble. Treat, therefore, so as to

---

* The rocking motion so frequently directed for paralytic, paretic, tumorous conditions, etc., is produced in the following manner: The operator adjusts the electrode to the desired location, then places the right hand upon it, and, with a firm, gentle motion, removes the pressure of the hand gradually from the wrist to the finger-tips, and from the finger-tips to the wrist, continuously and alternately, during the allotted time of treatment.