the passage, the negative in sponge-cup at its summit, moved slowly over the abdomen in D, twenty or thirty minutes, or No. 6 electrode placed across the abdomen in D. If the cramps are very severe, treat as follows:

Second. Apply hot compress over the abdomen, on which place No. 7 electrode in D, No. 8 electrode on the back, long point between the kidneys in B, ten to thirty minutes or more, very strong.

Third. No. 3 electrode from the neck downward on the spine in C, No. 4 electrode across the stomach in E, fifteen to twenty minutes.

Fourth. No. 5 electrode on the middle part of spine in B, No. 6 electrode over the upper abdomen in G, fifteen minutes, then over the lower abdomen ten minutes longer.

Fifth. No. 5 electrode from the base of brain, on the spine, in D, No. 6 electrode from stomach down over the navel in H, fifteen to thirty minutes.

Sixth. As the patient convalesces, administer F–G as a mild stimulant and use general treatments, frequent applications of thirty or forty minutes.

DIARRHŒA AND DYSENTERY.

First. No. 8 electrode on the abdomen in D, No. 3 or 4 electrode on the lower part of spine in B, fifteen to thirty minutes.

Second. No. 5 electrode on the lower part of spine in A, No. 6 electrode from the groin upward on one side of the abdomen in G, ten minutes, then on the other side ten minutes longer.