DROPSY.

Dropsy is a frequent sequel of kidney affections, as well as the result of many other diseases, and requires a most thorough diagnosis to ascertain the cause, which should be manipulated in conjunction with the following special treatments:

First. Place No. 7 electrode upon a chair or stool, and seat the patient upon it in G, with a large sponge-cup in A; stroke down the spine from the base of brain to the coccyx repeatedly, occasionally giving the sponge-cup an outward movement below the kidneys toward the hips; continue for fifteen or twenty minutes.

Second. No. 7 electrode upon a chair or stool in D, No. 6 electrode over the liver, stomach, and spleen in B, fifteen or twenty minutes.

Third. No. 7 electrode on a chair or stool in A, No. 8 electrode under the feet in D, or the foot-bath may be used, fifteen minutes.

Fourth. No. 8 electrode on the back, the point between the kidneys in H, No. 7 electrode over the abdomen, with a rocking motion, in B, fifteen minutes.

Fifth. For œdema of the legs, place No. 7 electrode under the feet in E, No. 5 electrode bent to encircle the leg in C, and by both of the operator's hands drawn gently downward over the legs alternately, five or seven minutes each.

Sixth. The vapor-bath, especially if there is swelling of the face, or if this is not convenient, No. 5 electrode from the base of brain down the spine in