In the married, the vaginal instrument is used. Here, as in all other internal procedures, it must be remembered that the mucous secretion acts in a dispersive manner on electric currents; the reversed currents are used in many of these applications.

First. Have hot and cold water at hand; and everything else in good order. Place the battery on a stand, near the right of the couch; attach the vaginal electrode to A, and let it lie in hot water, by way of preparation; before introducing, rub it with fine soap or glycerine, also lubricate the index finger, throw a sheet over the patient, who should raise and flex the left knee, keeping the hands down; examine and correct the displacement; then pass the vaginal electrode alongside of the finger to the posterior cul-de-sac, or, if there be metritis, to the os; as indeed may be done in other cases; withdraw the finger, holding the instrument inside in undisturbed position; place a support firmly under the electrode; lower the knee; remove the hand; have ready the sponge-cup, attached to the conductor but not to the battery, with its handle inside the outer sleeve of your right arm, that it may not touch the patient's limbs; place the palm on the outside of the cup, and carry it up to the left groin; place the sponge well over the ovary; connect to the battery in D; make the current as strong as can be borne, then quickly return the cylinder with the left hand, which will cause active contraction; continue five to seven minutes.

Second. When, after this time, the contraction has nearly ceased, lift the pin from the post, and so dis-