rocking motion, in E, No. 4 electrode small of back in C, fifteen minutes; then change to B–H current in the same manner but without the rocking motion.

Sixth. No. 4 electrode above the external prominence, slowly rocked, in A, the cylinder electrode enclosed in a sponge placed at rectum, or vagina, or use the vaginal instrument with D fifteen minutes; then remove D to the feet, allowing A to rest on the tumor ten minutes longer.

OVARIAN TUMOR.

Diagnose thoroughly; if displacement of the uterus is discovered, give it immediate attention. (See directions for displacement.)

First. Place No. 6 electrode across the external prominence in B, the sponge-cup moved with a rolling or kneading motion around the outer edge of the tumor externally in D, fifteen to thirty minutes.

Second. No. 8 electrode, the short point over the supra-pubic region in H, No. 6 electrode across the lower part of back in C, twenty-five to thirty-five minutes. The current from the SILVER HELIX may be used in the same manner.

Third. No. 5 electrode from the coccyx on the spine upward in B, No. 6 electrode across the abdomen over the tumor in D, ten minutes; change to No. 4 electrode a little above the centre of the external prominence, slowly rocked by the operator, in A, the cylinder electrode enclosed in a sponge placed at the entrance to vagina in D, fifteen minutes longer.

Fourth. If the tumor is attached to the ligament,