In other cases, organic or mechanical obstruction may exist above the ulcer, and must be discovered and corrected, as in varicose ulcer. Aside from such indications, use,

First. Select a suitable electrode in size and shape, to place above the sore in B, the sponge-cup stroked around and over it in D, fifteen or twenty minutes; C–D or A–D fifteen minutes to complete the treatment.

Second. No. 3 electrode immediately above the sore in G, cylinder electrode, enclosed in a soft sponge wet in a weak solution carbolic acid, placed over it, in H, twenty minutes; change to A–D ten minutes longer.

Third. Vapor-bath.

Internal ulcers, if accessible, should be treated always with the negative attached to the internal instrument, and the positive located externally above.

Fistula. Use first C–D, then B–D and C–E currents; alternate with G–H and A–D currents. After each treatment, if the fistula is small and the surface well dried, a piece of adhesive plaster may be made to adhere to the vaginal aspect, supported by a cotton-glycerine pessary, thus expediting cicatrization.

Administer daily applications.

Carbuncles. Treat as for phlegmonous erysipelas. The core, or slough of deep fascia, is to be liberated by the wire-brush cautery, as soon as possible (see directions erysipelas). In boils there is also a slough or core, formed, however, of the superficial fascia. The core of the felon is formed of the theca of the finger, or of the palmar fascia.