Felon. Prepare a bath for the finger of saltpetre and turpentine.

First. No. 3 electrode around the arm, just above the wrist, in A; place the finger in the prepared solution with the cylinder electrode connected with D, quite strong, twenty-five or thirty minutes.

Second. No. 3 electrode on the arm in G, No. 1 electrode placed over a fine soft sponge, wet in the solution recommended, held gently on the affected part in H, twenty minutes; change to C–E current. Bandage the finger and keep it thoroughly wet with the solution. If it is necessary to lance the felon, the sensation may be deadened by A–C current applied above and below it during the operation; immediately after apply A–D. The patient seldom loses a nail, and the tendons of the finger need not become indurated or inflexible after this treatment.

PUSTULA MALIGNA.

A species of malignant boil. It has been induced by the reception of an animal infection into the system; is gangrenous, and usually fatal, because it evades detection until the entire system has become inoculated. Immediate and frequent use of nitrate of silver to cauterize the pustule, and the application of a sponge-cup saturated in a powerful solution of carbonate of soda, in D, sponge-cup at base of the brain in A, twenty-five minutes. If a limb or finger is affected, plunge it in the solution with cylinder-electrode in D, sponge-cup or an electrode placed above in A to prevent the poison, if possible, from spreading through the blood. Poul-