with a too early cessation of the lochia after parturition, is followed by puerperal fever.

Menstruation, although thus classified, is not a secretion, but is simply a periodical flow of somewhat altered blood, induced by (physiological) ovarian congestion, incident to monthly ovulation.

The principal pathological deviations are amenorrhœa (non-appearance, suppression, or retention), dysmenorrhœa (painful menstruation), or menorrhagia (excessive menstruation).

The diagnosis of these affections belongs properly to the gynæcologists.

Perspiration, when strongly odoriferous, indicates vicarious excretions through the perspiratory glands, and arises from insufficient action of the kidneys and bowels. A characteristic odor of the perspiration is peculiar to measles, scarlatina, small-pox, Bright's disease, typhus, gout, etc.

Motor System.—Inability to rise from a recumbent position may be due to debility, paralysis, inflammation, or injuries. Inability to lie down, due generally to dyspnœa (difficult breathing). A desire to lie upon the abdomen denotes colic; upon the back with knees drawn up, peritonitis; upon the right side, enlargement of heart or liver; prone or semi-prone position, aneurism of aorta.

Muscular debility arises from total want of exercise, acute disease, exhaustion, prostration, paralysis.

Spasms are tonic, clonic, and choreic. Tonic spasm is fixed rigidity, as in tetanus; clonic, successive muscular contraction at short intervals, as in ordinary convulsions; choreic, a jerking, irregular, muscular movement, beyond the power of the will, as in chorea.

Subsultus tendinum is a jerking of the tendons at the wrist, and is common in low fevers, indicating approaching dissolution.

Hemiplegia is paralysis of one side of the body; paraplegia is paralysis of both lower extremities, and are terms applied to loss of power or loss of sensibility, or both.

Sensory System.—Here the important symptom is pain, and this may be acute, sharp, cutting, shooting, darting, lancinating,