is characterized by soreness, disposition to hawk and expectorate, with hoarseness or partial aphonia.

APHASIA.—*D.*, Loss of speech. Admitting the observations of Trousseau, and the subsequent speculations of Broca, that the seat of the faculty of language is "in the anterior frontal convolutions of the left hemisphere of the cerebrum," then softening or inflammation of, or injury to that part of the brain, would account for this disease.

LARYNGISMUS STRIDULUS.—*D.*, Closure of the glottis; *S.*, stridulous or shrill, whistling respiration. A disease of infancy, and occurs during dentition.

CROUP.—*D.*, Inflammation of the trachea; *V.*, inflammatory, pseudo-membranous; *S.*, a hoarse, barking cough, oppressed and audible respiration, spasm of the glottis and muscular coat of the trachea, and in the latter variety formation of false membrane in the larynx and trachea.

PLEURODYNIA.—*D.*, Intercostal rheumatism; *S.*, dull, heavy pain on one or both sides, increased by deep breathing or coughing, or motion of arms and trunk, without fever; *C.*, rheumatic diathesis.

INTERCOSTAL NEURALGIA.—*S.*, Sharp, lancinating, intermitting or periodical pains between sixth, ninth, and tenth ribs; *C.*, neuralgic diathesis.

THORACIC MYALGIA.—*D.*, Pain in the superficial muscles of the chest; *C.*, pressure, constrained position, ill nourishment, overwork.

PHTHISIS PULMONALIS.—*D.*, Tuberculous consumption of the lungs; this is by far the most important, because the most common and most fatal disease of the respiratory organs; *V.*, acute, chronic, and latent; *S.*, may be the sequelæ to any of the diseases already described, or may begin as a primary disease, with a slight hacking cough, or with a hemorrhage, both slowly increasing in severity (hemorrhage not always present), pains in the chest, pallor, accelerated pulse, increased temperature, chills followed by fever, emaciation, arrest of menstruation in females, night-sweats, diarrhœa; spirits of patient generally exalted and hopeful even to the last. Perhaps