all the symptoms of renal congestion and uræmic poison. The progress of the disease may be slow, and may be months, or even years, in reaching a climax, but the tendency is always towards a fatal result; *C.*, predisposing causes are intemperance, gout, constitutional syphilis, exposure to cold, scarlet fever, diseases of the kidneys, bladder, and urethra.

Lithiasis.—*Syn.*, Gravel; *D.*, calcareous deposits in the kidneys or bladder; *S.*, pain in the bladder and in the back, itching in the glans penis, retraction of testicle, sudden stopping of the flow of urine.

Diabetes Insipidus.—*Syn.*, Polyuria; *D.*, excessive urination; *S.*, excessive discharge of almost colorless urine (two to six gallons a day), intense thirst (withholding fluids does not lessen the excessive urination), skin hard and dry; debility and emaciation if long continued.

Diabetes Mellitus.—*Syn.*, Glycosuria; *D.*, excessive urination, urine containing sugar; *S.*, beginning with all the symptoms of diabetes insipidus, the disease gradually simulates pulmonary consumption, and in that form the case terminates.

Hydronephrosis.—*D.*, Dropsy of the kidneys; *S.*, a soft, lobular, fluctuating, painless tumefaction in the hypochondriac, umbilical, and iliac regions, generally on one side; when extensive may result in uræmia, or bursting of the sac into the abdominal cavity.

Cancer of the Kidney.—*S.*, A tumor in the abdomen (sometimes enormously large), with copious discharges of bloody urine, is almost certain to be cancerous.

Tubercle of the Kidney.—*S.*, Recognized only after tubercles have softened by the abundant discharge of purulent urine, which, in the absence of tumor and bloody urine, distinguishes it from cancer.

Cystitis.—*D.*, Inflammation of the bladder; *V.*, acute and chronic, primary and secondary.

Acute Cystitis.—*S.*, Chills and fever, pain in the bladder, frequent desire to urinate, burning pain in the urethra, and tenesmus. The bladder felt as a small round tumor, painful