the brain is the exciting cause. In the hemorrhagic form, the rupture of a blood-vessel in the cerebrum, cerebellum, or ventricles, and the formation of "blood-clot." Age is a predisposing cause, so also are excesses of all kinds, especially in florid, short-necked, obese people.

PARALYSIS.—*V.*, According to cause, 1, cerebral; 2, spinal; 3, reflex; 4, toxæmic; 5, hysterical. According to parts affected, facial or other local paralysis, hemiplegia, paraplegia, general.

FACIAL PARALYSIS.—*C.*, Affection of the motor nerve of the face, and is usually due to inflammation (rheumatic) of the sheath of the nerve, near the stylo-mastoid foramen; *S.*, one side of the face is expressionless, and the eye on that side remains open. Pressure on, or disease of any nerve, may cause paralysis to the part it communicates with, until pressure or disease is removed; from this cause we may have loss of sight, of hearing, of taste, of smell.

HEMIPLEGIA.—*D.*, Paralysis, or loss of motion and sensation of one side of the body; *C.*, usually blood-clot, tumor, or softening of the brain.

PARAPLEGIA.—*D.*, Paralysis of the lower extremities; *C.*, injury or disease of the spinal cord; *S.*, pain in the back, numbness of the feet, and a gradual (sometimes sudden) loss of motion and sensation in the lower extremities. If the disease of the spinal cord is progressive, so also will be the paralysis, and may finally involve the bladder and rectum.

HYSTERICAL PARALYSIS.—This is one of the many remarkable forms of nervous phenomena manifested in that unexplained and unexplainable disease, hysteria, and may simulate any form of paralysis, and can only be differentiated by the history of the patient.

REFLEX PARALYSIS.—*D.*, Paralysis without any apparent brain or spinal lesion, and may be caused by nervous dysentery, diarrhœa, uterine irritation, teething, external injuries, diphtheria, scarlatina, disease of the kidneys.

DIPHTHERITIC PARALYSIS.—*D.*, The muscles used in swallowing, speaking, and sometimes of the upper and lower ex-