tremities, and sense of sight may become partially paralyzed after the termination of a severe attack of diphtheria. *C.*, doubtful.

Syphilitic Paralysis.—*D.*, Obscure paralysis, with no apoplectic symptoms in syphilitic constitutions. But of undoubted existence.

Lead Palsy.—*C.*, By exposure to lead-poisoning; *S.*, commences with the muscles of the forearm; blue line along the edge of the gums; pain attends the beginning of and recovery from the paralysis.

Paralysis Agitans.—*D.*, Shaking palsy; *S.*, a constant involuntary shaking of the head, hands, and arms, and more or less of the whole body.

Wasting Palsy.—*D.*, Muscular atrophy; *S.*, the muscles of one limb or the whole body lose their power and waste away.

Locomotor Ataxia.—*Syn.*, Duchenne's disease, "Ataxie locomotrice progressive." *S.*, A gradual loss of muscular coordination, preceded by rheumatic pains, sometimes strabismus, and incontinence of urine; gradually the gait becomes unsteady and awkward, feet become numb, and walking is insecure, usually resulting in complete general paralysis.

Paralysis of the Insane.—*D.*, Difficulty of speech, general tremor, gradual loss of motion and sensation characterize it.

Epilepsy.—*D.*, Periodical convulsions; *S.*, a creeping, blowing sensation like a current of air or a stream of water, commencing in the hand or foot and creeping towards the body, is called the "epileptic aura." With a scream the patient falls down in violent convulsions; foaming at the mouth, grinding of the teeth, rolling of the eyeballs, sometimes vomiting, or involuntary discharges from bladder or rectum, are common symptoms. *C.*, Hereditary transmission, intemperance, excessive venery, self-abuse, blows on the head, and fright are common causes.

Catalepsy.—*S.*, Unconsciousness, fixed rigidity of all the voluntary muscles.