the blood-vessels; symptomatic when occurring in connection with certain diseases, as bleeding of the nose in typhoid fever, bleeding of the lungs in phthisis, vomiting of blood in cancer of the stomach, bleeding from the bowels in hemorrhoids, etc.; critical, as at the close of yellow fever or remittent fever; vicarious when it follows as a substitute the suppression of a normal or habitual discharge, as in suppressed menstruation or sudden arrest of bleeding hemorrhoids, and manifests itself generally in the form of epistaxis, hæmatemesis, and hæmoptysis.

Hæmoptysis.—*D.*, Spitting of blood; when from the air-tubes, it is attended with difficult breathing; *S.*, the blood is coughed up, is florid, and sometimes frothy and mixed with expectoration.

Hæmatemesis.—*D.*, Vomiting of blood, when from the stomach; *S.*, there is nausea, the blood is vomited up, is dark and mixed with food.

Hæmaturia.—*D.*, Bloody urine may be from the kidneys or bladder. If thoroughly mixed with urine, it is probably from the kidneys, and may be due to venal congestion, or inflammation, or cancer; when the urine flows nearly pure and the blood follows, it is from the bladder or urethra.

Intestinal Hemorrhage.—*C.*, May be bleeding hemorrhoids, in which case the blood is bright red; in other hemorrhages, as from typhoid, yellow, or remittent fever, or aortic aneurism, congestion of the liver or abdominal cancer, it is dark and mixed with fæces.

Uterine Hemorrhages.—*C.*, may be excessive menstruation, placenta prævia, abortion, uterine cancer, ulceration of the mouth or neck of the uterus, tumors.

Dropsy.—*V.*, General and œdema; *D.*, tumefaction arising from the effusion or infiltration of serum into cellular or connective tissue; *C.*, one of three cardinal pathological conditions:

1. Obstruction to the venous circulation.
2. Arrest of excretion and absorption.
3. Excess of water in the blood.