physician doubtful of his diagnosis in calling it either one or the other.

Mumps.—*S.*, Contagious inflammation of the parotid gland, which swells and becomes hot, painful, and tender to the touch; one or both may be affected, either simultaneously or after an interval of years. It is doubtless contagious, but not dangerous, unless when by metastasis it is transferred to the breast, testicles, or brain, which may be avoided by guarding against taking cold.

Whooping-Cough.—*S.*, Six days of incubation, after exposure, are followed by symptoms of a cold, with slight fever and a cough, which soon shows its specific character by being paroxysmal and whooping. It is a self-limited disease, but seldom ends in less than six weeks, and sometimes lingers for months.

Diphtheria.—*S.*, A few days of indisposition, slight sore throat, swelling of glands behind the jaw, fever, headache, furred tongue, constipation, difficulty of swallowing, palate, tonsils, and fauces swollen, red or purple. Soon a whitish or yellowish-white membrane appears over one or both of the tonsils. All the symptoms of pseudo-membranous croup may here follow, or it may assume an ulcerated form and become "putrid sore throat," or a malignant form with all the symptoms aggravated, and followed by prostration, coma, and death.

Glanders.—This is a disease peculiar to the horse, but may undoubtedly be communicated to man; *S.*, manifests itself in from two to seven days after exposure, by fever, inflammation of the nostrils, pains in the joints, patches of red skin that may become gangrenous, pustules on the face and limbs, inflammation of nostrils becomes ulcerous and even gangrenous, discharging a muco-purulent fluid; throat and lungs may become involved, the face becomes œdematous; diarrhœa, delirium, coma, and death.

Influenza.—*Syn.*, Epidemic catarrh. Common as catarrhal affections are, there is an occasional epidemic form of this disease in which the symptoms of a "bad cold" are all aggravated.