and affect a whole community to an extent that is hardly accounted for by ordinary climatic conditions. It is rarely fatal except among very old people.

Intermittent Fever.—*Syn.*, Ague, chills and fever; *S.*, First, or cold stage.—After a short period of languor and yawning, a cold, creeping, shivering sensation ushers in the chill, which may last from ten minutes to three hours.

Second, or hot stage.—A sense of warmth gradually returns, which resolves itself into a hot or dry fever, more or less violent, with accompanying headache, often vomiting; mouth dry and tongue furred. This stage may last from one to sixteen hours, and is followed by the third, or sweating stage, which is somewhat gradual in its approach; face first becomes moist, soon the whole body, the fever subsides, headache lessens, and, with a profuse perspiration, the patient is likely to sink into a quiet, peaceful slumber, and all goes well until the intermission is passed and the first stage returns, which may be in one, two, three, four, five, six, seven, or eight days. *C.*, it may reasonably be called an autumnal fever, as comparatively few cases occur at any other time of the year, and is the result only of malarial influences, near sea-level, and near marshes, shallow lakes, and sluggish streams.

Remittent, or Bilious Fever.—*S.*, First general indisposition, slight headache, nausea, furred tongue, followed by a chill lasting for perhaps an hour; after this comes fever, skin hot, dry, harsh, pulse accelerated, face flushed, head aching and throbbing, violent pain in back and limbs, nausea and vomiting of bilious matter common, bowels costive and stools colored with bile, and thirst great. There is an abatement of these symptoms, at least in severity, in from eight to twenty hours; after from six to twenty-four hours of comparative comfort the chill again comes on, and all the conditions are more aggravated than at first, and then follows a succession of periodical paroxysms that must vary with the history of the case; if uncomplicated, it may terminate favorably in a week; but when, as sometimes happens, inflammations, such as of the brain, lungs, stomach, liver, spleen, or bowels, intervene, the