POISONS AND THEIR ANTIDOTES.

POISONS.	ANTIDOTES.
Acids.	The Alkalies. Common soap (soft or hard), in solution, is an efficient remedy, and has the advantage of being always at hand. It should be followed by copious draughts of tepid water or flaxseed tea. For *nitric* and *oxalic* acids, the carbonates of magnesia and lime (chalk and water) are the best antidotes. When sulphuric acid has been taken, the use of much water is improper.
Alkalies and their salts.	The Vegetable Acids. Common vinegar being always at hand, is most frequently used. The fixed oils, as castor, flaxseed, almond, and olive oils, form soap with the alkalies, and thus also destroy their caustic effect. They should be given in *large quantity.*
Baryta and its salts. Lime.	*Epsom* or *Glauber's Salt*, in solution, or diluted sulphuric acid. The fixed oils also have the same effect as with the alkalies proper, when these bases are uncombined.
Iodine. Iodide of Potassium.	Starch, wheat flour, or arrow-root, in large quantities, well mixed with water. Let the patient drink *freely* of starch, or wheat flour and water, and afterwards of a strong mixture of vinegar and water (any acid in which there is oxygen); when this is done, the whole process of saving the patient's life is accomplished, and it will only be necessary to evacuate the bowels.
Cyanide of Potassium and Prussic Acid.	Chlorine water; solution of chlorinated soda; aqua ammonia; cold affusion.
Antimony and its salts.	Astringent infusions, as of galls, oak bark, Peruvian bark, or green tea very strong.
Arsenic and its compounds.	Any oil or fat (sweet oil, butter, milk); magnesia in large quantity; hydrated oxide of iron (recently made), in tablespoonful doses, every 5 or 10 minutes.
Bismuth and its compounds.	Albumen. Copious draughts of milk, with sweet mucilaginous drinks.
Copper and its compounds.	Albumen, as milk or white of egg in solution, should be freely administered. Ferrocyanuret of potassium (freely). *Vinegar must be avoided.*
Gold, salts of.	Sulphate of iron, with a free use of mucilaginous drinks.
Iron, salts of.	Carbonate of soda, with mucilaginous drinks.
Lead, salts of.	Albumen; sulphate of magnesia (Epsom salts); sulphate of soda (Glauber's salt); diluted sulphuric acid; lemonade; opium.
Mercury, salts of.	Albumen, as white of egg, milk, or wheat flour beaten up with water; to be followed by an emetic.
Silver, salts of.	Common salt, freely, in solution.