POISONS.	ANTIDOTES.
Tin, salts of.	Albumen; white of egg; milk; or flour.
Zinc, salts of.	Albumen, or carbonate of soda, with copious draughts of warm water, and especially milk.
Phosphorus.	Magnesia, with water and copious draughts of mucilaginous drinks.
Gases.	Ammonia, cautiously inhaled, is recommended for chlorine. Asphyxia, produced by noxious gases, must be treated by copious cold affusions, especially to the head; blood-letting; artificial respiration; and stimulants carefully administered.
Creasote.	Albumen, or white of egg; milk; or wheat flour.
Alcohol or spirituous liquors.	A powerful emetic should be given, followed by copious draughts of warm water. Congestion of the brain, and other symptoms, are to be treated on general principles.
Opium and other narcotics.	The chief reliance is to be placed on the most active emetics (as mustard, alum, tartar emetic, or sulphate of zinc) and the stomach-pump. Emetics are preferable to the stomach-pump, when the narcotic has been taken in substance. The patient should be kept in motion, and cold water dashed on the head and shoulders. Should the above means fail, the electro-magnetic battery, or if that cannot be obtained, artificial respiration must be resorted to.
Strychnia.	Emetics should be freely given. Kermes mineral; tannic acid; tincture of iodine; chlorine water; ether or chloroform, by inhalation. If these means fail, Dr. Marshall Hall's ready method may be tried.
Arnica.	Vinegar.
Aconite.	Stimulants, externally and internally.
Veratrum Viride.	Laudanum; alcohol.

TO MAKE HYDRATED OXIDE OF IRON.

"Take of Solution of Tersulphate of Iron, a pint; Water of Ammonia, Water, each, a sufficient quantity. To the solution of Tersulphate of Iron, previously mixed with three pints of Water, add Water of Ammonia with constant stirring, until in slight excess. Then pour the whole on a wet muslin strainer, wash the precipitate with water, pressing the strainer forcibly with the hands until no more liquid passes. Lastly, mix the precipitate with sufficient Water to bring the mixture to the measure of a pint and a half, and transfer it to a wide-mouthed bottle, which must be well stopped."—U. S. P.