


# Resistance Mechanisms



Monitoring Approaches: Serial biopsies and liquid biopsy track resistance evolution

# Primary (Intrinsic) Resistance

## Pre-existing Resistance Factors



### Key Characteristics

- ▶ Pre-existing in tumor cells before treatment
- ▶ Germline or somatic mutations present at diagnosis
- ▶ Tumor heterogeneity with resistant subclones
- ▶ Immediate treatment failure or minimal response


### Mechanisms

- ▶ Target gene mutations preventing drug binding
- ▶ Low or absent target protein expression
- ▶ High expression of drug efflux transporters
- ▶ Pre-existing bypass pathway activation


### Clinical Example

**EGFR-mutant NSCLC with T790M:** Approximately 60% of EGFR-mutant non-small cell lung cancers that develop resistance to first-generation EGFR TKIs (erlotinib, gefitinib) harbor the T790M gatekeeper mutation, which increases ATP affinity and prevents drug binding.

# Acquired Resistance



## Adaptive Mechanisms During Therapy



### Key Characteristics

- ▶ Develops during treatment exposure
- ▶ Initial response followed by progression
- ▶ Clonal selection under drug pressure
- ▶ Time to resistance: months to years

### Common Mechanisms

- ▶ On-target: Secondary mutations in target gene
- ▶ Off-target: Parallel pathway activation
- ▶ Drug metabolism: Increased efflux, decreased uptake
- ▶ Histologic transformation: SCLC conversion

### Clinical Example


**Osimertinib Resistance in EGFR T790M+ NSCLC:** C797S mutation emerges in approximately 15-30% of patients treated with osimertinib (third-generation EGFR TKI), preventing covalent drug binding. MET amplification accounts for another 15-20% of resistance mechanisms.

# Bypass Pathway Activation


## Normal Signaling



## Drug Block



## Bypass Activation



## Key Characteristics

- ▶ Alternative pathways maintain signaling output
- ▶ Signal redundancy overcomes single-target inhibition
- ▶ Can be pre-existing or acquired during treatment
- ▶ Often involves parallel RTK activation

## Common Bypass Routes


- ▶ MET amplification bypassing EGFR inhibition
- ▶ HER2/HER3 activation maintaining PI3K signaling
- ▶ IGF-1R upregulation in hormone receptor+ breast cancer
- ▶ PIK3CA mutations activating downstream signaling

## Clinical Example


**MET Amplification in EGFR-mutant NSCLC:** MET gene amplification occurs in 5-15% of patients with acquired resistance to EGFR TKIs. MET activation maintains downstream PI3K/AKT and MAPK signaling despite EGFR inhibition. Combination of EGFR and MET inhibitors shows clinical benefit in this subset.

# Combination Therapy Strategies


## Vertical Blockade



## Horizontal Blockade



## Targeted + Immune



### Benefits of Combination Therapy

- Delay or prevent resistance emergence
- Synergistic efficacy
- Target multiple resistance mechanisms
- Address tumor heterogeneity
- Deeper responses

## Combination Approaches

- ▶ Vertical: Target multiple nodes in same pathway
- ▶ Horizontal: Inhibit parallel signaling pathways
- ▶ Targeted + Chemo: Overcome different resistance types
- ▶ Targeted + Immune: Direct + immune-mediated killing

## Clinical Considerations

- ▶ Increased toxicity requiring dose adjustments
- ▶ Cost and access considerations
- ▶ Biomarker-driven selection of combinations
- ▶ Sequential vs. concurrent administration

## Clinical Example



**Dabrafenib + Trametinib in BRAF V600E Melanoma:** Combined BRAF and MEK inhibition (vertical blockade) significantly improves progression-free survival compared to BRAF inhibitor alone (11 vs 7 months median PFS). The combination delays resistance by preventing feedback reactivation of the MAPK pathway and reduces paradoxical MAPK activation.

# Resistance Mechanisms: Summary & Clinical Impact

## Primary Resistance

- ▶ **Timing:** Present before treatment
- ▶ **Frequency:** 20-30% of patients
- ▶ **Strategy:** Upfront biomarker testing, alternative therapies

## Acquired Resistance

- ▶ **Timing:** Develops during therapy
- ▶ **Frequency:** Nearly universal (6-24 months)
- ▶ **Strategy:** Resistance profiling, sequential therapies

## Bypass Pathways

- ▶ **Mechanism:** Alternative signaling activation
- ▶ **Frequency:** 15-25% of resistance cases
- ▶ **Strategy:** Horizontal combination therapy

## Combination Strategy

- ▶ **Goal:** Prevent/delay resistance
- ▶ **Approaches:** Vertical, horizontal, multi-modal
- ▶ **Challenge:** Toxicity management

## Clinical Decision Framework

### Initial Assessment

- Comprehensive genomic profiling
- Tumor heterogeneity evaluation
  - Baseline resistance screening

### Monitoring

- Serial liquid biopsy (ctDNA)
- Radiologic response tracking
- Resistance biomarker surveillance

### Progression Strategy

- Re-biopsy for mechanism
- Mechanism-directed therapy
- Clinical trial consideration

**Key Takeaway:** Understanding resistance mechanisms guides rational therapy selection, optimal sequencing strategies, and novel combination approaches to improve patient outcomes in targeted cancer therapy.