

1. (a) We begin by noting that:

$$0 + 0 = 0 \in S + I$$

$$1 + 0 = 1 \in S + I$$

Let $s_x + a_x, s_y + a_y \in S + I$. We note that $-s_x \in S$ and $-1 \cdot a_x \in I$. Thus:

$$-s_x + (-a_x) = -(s_x + a_x) \in S + I$$

For addition, we get that:

$$s_x + a_x + s_y + a_y = (s_x + s_y) + (a_x + a_y) \in S + I$$

For multiplication, we get that:

$$(s_x + a_x) \cdot (s_y + a_y) = (s_x s_y) + (s_y a_x + s_x a_y + a_x a_y) \in S + I$$

Thus, we conclude that $S + I$ is a subring.

- (b) We begin by noting that $0 \in I$ and $0 \in S$, so $0 \in S \cap I$. Let $a, b \in S \cap I$. We note that $a + b \in S$ and $a + b \in I$. Thus, $a + b \in S \cap I$. Lastly, let $s \in S$. We note that $as \in S$ and $as \in I$, so $as \in S \cap I$. This proves $S \cap I$ is an ideal of S .
- (c) We first note that $S \subseteq S + I$. Thus, the natural projection map $\pi : S \rightarrow (S + I)/I$ where $s \mapsto [s]$ is a ring homomorphism.

We then note that for all $x \in (S + I)/I$, there exists a $s + a$ such that $s \in S$ and $a \in I$ where $x = [s + a] = [s] + [a] = [s]$ because $I \mid a$, which implies $[a] = [0]$. Thus, $x = \pi(s)$. Hence, π is surjective, so it implies $\text{im}(\pi) = (S + I)/(I)$.

For $x \in S \cap I$, $\pi(x) = [x] = [0]$ as $I \mid x$, so $x \in \ker(\pi)$. Meanwhile, for $x \in \ker(\pi)$, we note that $[x] = [0]$, which implies $I \mid x$ or $x \in I$. Thus, we get $x \in S \cap I$ and that $\ker(\pi) = S \cap I$.

By the First Isomorphism Theorem, we get that $S/(S \cap I) \cong (S + I)/I$.

2. (a) We note that $[0] \in J'$, so $0 \in J$. For $a, b \in J$, we get that $[a], [b] \in J'$, so $[a + b] \in J'$ and $a + b \in J$. For $r \in R$, note that $[r] \in R/I$, so $[ra] \in J'$ thus $ra \in J$. This proves J is an ideal of R .
- (b) We note that $0 \in J$, so $[0] \in J/I$. For $[a], [b] \in J/I$, we note that $a + b \in J$, so $[a + b] = [a] + [b] \in J/I$. For $[r] \in R/I$, we note that $ra \in J$, so $[ra] = [r] \cdot [a] \in J/I$. This proves J/I is an ideal of R/I .
- (c) We note that the natural projection map of $\pi : R \mapsto R/I$ where $r \mapsto [r]_I$ is a ring homomorphism. Meanwhile, the natural projection map of $\hat{\pi} : R/I \rightarrow (R/I)/(J/I)$ where $[r]_I \mapsto [[r]_I]_{J/I}$ is a ring homomorphism. Hence, if we denote $\varphi = \hat{\pi} \circ \pi$, we get that $\varphi : R \mapsto (R/I)/(J/I)$ where $r \mapsto [[r]]$ is a ring homomorphism.

For all $x \in (R/I)/(J/I)$, there exists an $[r] \in R/I$ where $x = [[r]]$ and consequently, an $r \in R$ where $[[r]] = \varphi(r)$. This proves φ is surjective, so $\text{im}(\varphi) = (R/I)/(J/I)$.

For $x \in J$, $\varphi(x) = [[x]]$. We also note that $[x] \in J/I$, so $[[x]] = [[0]]$ thus $x \in \ker(\varphi)$. Meanwhile, for $x \in \ker(\varphi)$, $\varphi(x) = [[0]]$, so $[x] \in J/I$, which further implies $x \in J$. Thus, $\ker(\varphi) = J$.

By the First Isomorphism Theorem, $R/J \cong (R/I)/(J/I)$.

3. (a) A proper ideal I is prime ideal if and only if $ab \in I$ then $a \in I$ or $b \in I$

(b) Assume (r) is a prime ideal. By contradiction, we assume r is not prime. Then there exists $a, b \in R$ where $r \mid ab$ but $r \nmid a$ and $r \nmid b$. We note that $ab \in (r)$. This implies either $a \in (r)$ or $b \in (r)$. If we assume $a \in (r)$, then there exists a $q \in R$ where $rq = a$, but that would mean $r \mid a$, a contradiction. Thus, if (r) is a prime ideal, then r is prime.

Assume r is prime. By contradiction, we assume (r) is not a prime ideal, so there exists an $ab \in (r)$ where $a, b \notin (r)$. Since $ab \in (r)$, there exists a $q \in R$ where $ab = rq$, so $r \mid ab$. This implies either $r \mid a$ or $r \mid b$. We assume $r \mid a$, so there exists a $q' \in R$ where $q'r = a$. However, this implies $a \in (r)$, a contradiction. Thus, if r is prime, then (r) is a prime ideal.

(c) Assume I is a prime ideal of R . By contradiction, we assume R/I is not an integral domain, so there exists $[a], [b] \neq [0]$ and $[ab] = [0]$. Since $[ab] = [0]$, $I \mid ab$. This implies either $I \mid a$ or $I \mid b$, so either $[a] = [0]$ or $[b] = [0]$, a contradiction. Thus, if I is a prime ideal, R/I is an integral domain.

Assume R/I is an integral domain. By contradiction, I is not a prime ideal of R . Thus, there exists $a, b \notin I$ but $I \mid ab$. This implies $[ab] = [0]$. However, R/I is an integral domain, so either $[a] = [0]$ or $[b] = [0]$. But that implies either $I \mid a$ or $I \mid b$, a contradiction. Thus, if R/I is an integral domain, then I is a prime ideal of R .

(d) We note that \mathbb{Z} is a PID. Thus, for all prime ideals I of \mathbb{Z} , there exists an $x \in \mathbb{Z}$ where $I = (x)$. From b), x must be prime. By Euclid's Lemma, all prime numbers are prime, so their principal ideals are also prime ideals. However, 0 satisfies the definition of prime because \mathbb{Z} is an integral domain, so if $0 \mid ab$ then either a or b must be zero, so (0) is also a prime ideal. Thus, all prime ideals of \mathbb{Z} are principal ideals of prime numbers and 0 .

(e) From 1b), we note that $S \cap I$ is an ideal. For $a, b \in S$, if $ab \in S \cap I$, then $ab \in I$, which implies either $a \in I$ or $b \in I$. In other words, we get that either $a \in S \cap I$ or $b \in S \cap I$. Thus, $S \cap I$ is a prime ideal.

4. (a) If $[x]^2 = [x]$ for $0 \leq x \leq 2024$, then it implies $[x^2 - x] = [0]$ or $2025 \mid x(x-1)$. We then note that $2025 = 81 \cdot 25$ and that $\gcd(81, 25) = 1$. Thus, we can apply Theorem 7.11, where $m = 81$ and $n = 25$ and note that:

$$\begin{aligned} [x(x-1)]_{2025} &\mapsto [x(x-1)]_{81} \times [x(x-1)]_{25} \\ [0] &\mapsto [0]_{81} \times [0]_{25} \end{aligned}$$

We also note that this map is a ring homomorphism, so since $[x^2 - x] = [0]$, it implies that $[x(x-1)] = [0]_{81}$ and $[x(x-1)] = [0]_{25}$. In other words, we get that $81 \mid x(x-1)$ and $25 \mid x(x-1)$. We then note that 25 is a prime power of 5^2 . By Euclid's Lemma, either $5 \mid x$ or $5 \mid x-1$. Since $\gcd(x-1, x) = 1$, only one of the factors could be divisible by 5 and will be the one also divisible by 25. A similar argument can be applied that only one of the factors is divisible by 81. Thus, we get that either $25 \mid x$ or $25 \mid x-1$ and $81 \mid x$ or $81 \mid x-1$. This gives us 4 possible combinations.

Case 1 If $25 \mid x$ and $81 \mid x$, since 81 and 25 are co-prime, we get that $2025 \mid x$. The only x that satisfies this is if $x = 0$.

Case 2 If $25 \mid x-1$ and $81 \mid x-1$, since 81 and 25 are co-prime, we get that $2025 \mid x-1$. The only x that satisfies this is if $x-1 = 0$ or $x = 1$.

Case 3 If $25 \mid x-1$ and $81 \mid x$, it implies there exist $a, b \in \mathbb{Z}$ where $25a = x-1$ and $81b = x$. Thus:

$$\begin{aligned} 25a &= 81b - 1 \\ 1 &= 81b + 25(-a) \end{aligned}$$

We apply the Division Algorithm strategy back in Claim 2.7 to compute that $b = 21$, so $x = 21 \cdot 81 = 1701$.

Case 4 If $25 \mid x$ and $81 \mid x-1$, it implies there exist $a, b \in \mathbb{Z}$ where $25a = x$ and $81b = x-1$. Thus:

$$\begin{aligned} 81b &= 25a - 1 \\ 1 &= 25a + 81(-b) \end{aligned}$$

We apply the same strategy to compute that $a = 13$ thus $x = 13 \cdot 25 = 325$

Hence, there are 4 idempotent elements in $\mathbb{Z}/2025\mathbb{Z}$.

- (b) We note that $0^2 = 0$ and $1^2 = 1$, so $0, 1 \in S$. For $a, b \in S$, we note that:

$$\begin{aligned} (a+b)^2 &= a^2 + 2ab + b^2 \\ &= a^2 = 2 \cdot 1 \cdot ab + b^2 \\ &= a^2 + b^2 \\ &= a + b \end{aligned}$$

Thus, $a + b \in S$. Meanwhile::

$$\begin{aligned} (ab)^2 &= a^2b^2 \\ &= ab \end{aligned}$$

Thus, $ab \in S$. Lastly, we note that:

$$\begin{aligned} a + a &= 2a \\ &= 2 \cdot 1 \cdot a \\ &= 0 \end{aligned}$$

Thus, we note that $-a = a$ and since $a \in S$, we get that $-a \in S$. We proved S is a subring of R .

- (c) We first note that $(0) = \{r0 : r \in R\} = \{0\}$. Meanwhile, we note that the map $\varphi : R \rightarrow R$ where $r \mapsto r$ is a ring homomorphism. Meanwhile, the $\text{im}(\varphi) = R$ and that $\ker(\varphi) = \{0\} = (0)$. By the First Isomorphism Theorem, we get that $R/(0) \cong R$.

For $(e) + (1 - e)$, we note that for all $r \in R$ that

$$er + (1 - e)r = 1r = r$$

Hence, $r \in (e) + (1 - e)$ and $(e) + (1 - e) = R$. This allows us to apply Theorem 8.24 to get that $R/((e)(1 - e)) \cong R/(e) \times R/(1 - e)$. We then note that for any $a, b \in R$, we get that

$$(1 - e)a \cdot (e)b = (e - e^2)ab = 0ab = 0$$

This implies that any finite sum in the form of $\sum (e)a_i(1 - e)b_i$ is a sum of finitely many zeros, which sums to zero. Hence, $(1 - e)(e) = \{0\} = (0)$ and we get that $R \cong R/(0) \cong R/(e) \times R/(1 - e)$ or $R \cong R/(e) \times R/(1 - e)$ as desired.

- (d) e