



Recruitment rate stochasticity at the design stage of a clinical trial

Supervision by Malgorzata Roos

Pilar Pastor



Why recruitment rates?

- Timely recruitment vital to the success of a clinical trial
- Inadequate number of subjects → lack of power
- Recruitment period too long → competing treatments
- Recruitment of patients varies at each stage
- Accrual = Cumulative Recruitment
- [Carter \(2004\)](#)




Target Population





Target Population

Eligibility







Enrollment





Randomization






Statistical Analysis





Patient Attrition



Uncertainty

- **Aleatory**: randomness inherent and unpredictable
- **Epistemic**: arises from limited knowledge about parameters




Figure: Visualization of two types of uncertainty (Yang and Li, 2023)



Models for Counts

Methods	Counts	Expectation	Variance	Aleatory	Epistemic
Expectation	$C(t) = \lambda t$	λt	0	No	No
Poisson	$C(t) \sim \text{Po}(\lambda t)$	λt	λt	Yes	No
Negative Binomial	$C(t) \sim \text{Po}(\cdot t); \Lambda \sim G(\alpha, \beta)$	$\frac{\alpha}{\beta}$	$\frac{\alpha(\beta+1)}{\beta^2}$	Yes	Yes

Table: Aleatory and epistemic uncertainty in accrual shown by different models for counts.



Shrunk Slide 1



Shrunk slide 2



Results



Conclusion



Next steps



References

- Carter, R. E. (2004). Application of stochastic processes to participant recruitment in clinical trials. *Controlled clinical trials*, 25(5):429–436.
- Yang, C.-I. and Li, Y.-P. (2023). Explainable uncertainty quantifications for deep learning-based molecular property prediction. *Journal of Cheminformatics*, 15(1):13.