



Recruitment rate stochasticity at the design stage of a clinical trial

Supervision by Malgorzata Roos

Pilar Pastor



Why recruitment rates?

- Timely recruitment vital to the success of a clinical trial
- Inadequate number of subjects → lack of power
- Recruitment period too long → competing treatments
- Recruitment of patients varies at each stage
- Accrual = Cumulative Recruitment
- [Carter \(2004\)](#)




Target Population



Target Population





Eligibility







Enrollment





Randomization






Statistical Analysis



Patient Attrition



Uncertainty

- **Aleatory**: randomness inherent and unpredictable
- **Epistemic**: arises from limited knowledge about parameters




Figure: Visualization of two types of uncertainty (Yang and Li, 2023)



Models for Counts

Methods	Counts	Expectation	Variance	Aleatory	Epistemic
Expectation	$C(t) = \lambda t$	λt	0	No	No
Poisson	$C(t) \sim \text{Po}(\lambda t)$	λt	λt	Yes	No
Negative Binomial	$C(t) \sim \text{Po}(\cdot t); \Lambda \sim G(\alpha, \beta)$	$\frac{\alpha}{\beta}$	$\frac{\alpha(\beta+1)}{\beta^2}$	Yes	Yes

Table: Aleatory and epistemic uncertainty in accrual shown by different models for counts.



Study

- Time $t = 550$ days
- Recruitment Rate $\lambda = \frac{\text{Counts}}{\text{Time}} = 0.591$



Study

- Time $t = 550$ days
- Recruitment Rate $\lambda = \frac{\text{Counts}}{\text{Time}} = 0.591$
- Models for Counts:
 - Expectation: $EC(t) = \lambda t = 0.591 \cdot 550 = 325$
 - Poisson: $C(t) \sim \text{Po}(\lambda t)$
 - Negative Binomial: $C(t) \sim \text{Po}(\tilde{t}); \Lambda \sim G(\alpha, \beta)$



Study

- Time $t = 550$ days
- Recruitment Rate $\lambda = \frac{\text{Counts}}{\text{Time}} = 0.591$
- Models for Counts:
 - **Expectation:** $EC(t) = \lambda t = 0.591 \cdot 550 = 325$
 - **Poisson:** $C(t) \sim \text{Po}(\lambda t)$
 - Negative Binomial: $C(t) \sim \text{Po}(\Lambda t); \Lambda \sim G(\alpha, \beta)$




Accrual at time point t

- **Expectation:** $EC(t) = \underbrace{EC + \dots + C}_{t \text{ times}} = tEC = \lambda t$
- **Poisson:** $\underbrace{Po(\lambda) + \dots + Po(\lambda)}_{t \text{ times}} = Po(\lambda t)$



Accrual of 1 study


Accrual of 1 study






Accrual of 2 studies

Accrual of 2 studies




Accrual of 100 studies






Theoretical PMF






Theoretical CDF



Uncertainty bands





Negative binomial derived from Poisson-Gamma model

Recruitment in one unit of time ($t = 1$)

....



Study

- Time $t = 550$ days
- Recruitment Rate $\lambda = \frac{\text{Counts}}{\text{Time}} = 0.591$
- Models for Counts:
 - Expectation: $EC(t) = \lambda t = 0.591 \cdot 550 = 325$
 - **Poisson:** $C(t) \sim \text{Po}(\lambda t)$
 - **Negative Binomial:** $C(t) \sim \text{Po}(\Lambda t); \Lambda \sim G(\alpha, \beta)$



Comparison between Poisson and Negative Binomial - PMF



Comparison between Poisson and Negative Binomial - CDF



Shrunk Slide 1



Results



Conclusion



Next steps



References

- Carter, R. E. (2004). Application of stochastic processes to participant recruitment in clinical trials. *Controlled clinical trials*, 25(5):429–436.
- Liu, J., Jiang, Y., Wu, C., Simon, S., Mayo, M. S., Raghavan, R., and Gajewski, B. J. (2023). *accrual: Bayesian Accrual Prediction*. R package version 1.4.
- Spiegelhalter, D., Pearson, M., and Short, I. (2011). Visualizing uncertainty about the future. *Science*, 333(6048):1393–1400.
- Yang, C.-I. and Li, Y.-P. (2023). Explainable uncertainty quantifications for deep learning-based molecular property prediction. *Journal of Cheminformatics*, 15(1):13.