

– Supplementary material –

Production of neutral species in Titan’s ionosphere through dissociative recombination of ions

Sylvain Plessis

Univ Paris-Sud, Laboratoire de Chimie Physique,
UMR 8000, Orsay, F-91405

Nathalie Carrasco

Laboratoire Atmosphères, Milieux, Observations Spatiales,
Université de Versailles Saint-Quentin, F-78280, France

Michel Dobrijevic

Université de Bordeaux, Observatoire Aquitain des Sciences de l’Univers,
CNRS, UMR 5408, Laboratoire d’Astrophysique de Bordeaux,
2 rue de l’Observatoire, BP 89, F-33271 Floirac Cedex, France

and

Pascal Pernot

Univ Paris-Sud, Laboratoire de Chimie Physique, UMR 8000, Orsay, F-91405
CNRS, Orsay, F-91405
`pascal.pernot@u-psud.fr`

December 14, 2011

1 The Database

Models of ion-neutral chemistry generally include DR processes through the reaction rate constant

$$\alpha(T_e) = \alpha_0 \times (T_e/T_0)^{-\beta} \quad (1)$$

where α_0 is the rate constant at a reference electron temperature T_0 (typically 300 K), and T_e is the electron temperature of interest. When more than one product channel is accessible, the partial rate for channel i is $\alpha_i(T_e) = b_i \times \alpha(T_e)$, where $\{b_i, i = 1, N\}$ are the *branching ratios* of the reaction. As the global DR rate $\alpha(T_e)$ and the branching ratios $\{b_i\}$ are typically derived from different experiments, they are treated separately.

The following tables report data for ions identified in Titan’s ionosphere. **At the present stage, this is a demonstration database for the use of Nested Dirichlet representations of uncertain branching ratios, and it should not be considered or used as a reference database without caution.**

1.1 Rate constant parameters

The procedure for data representation as probability density functions is explained in the main paper.

1.2 Branching ratios

Information on branching ratios is typically sparser than for reaction rates, and it is exceptional to have to consider conflicting data. The following set of considerations is used in order to define the structures of the (Nested) Dirichlet distributions.

Maximum number of fragments. In absence of experimentally characterized products, one should consider all the exoergic channels and state a total lack of knowledge on the corresponding branching ratios (Dirichlet distribution). It is not possible to make further hypotheses wrt. the relative stability of the products. From Ref. [1], we know for instance that the measured branching ratios have no definite correlation with the exoergicity of the pathways. An issue in building a list of exoergic pathways is the number of fragments that can be accepted in each pathway. For instance, on the basis of their previous results, for $C_3H_7^+$ Ehlerding *et al.*[2] stop at a three body breakup pattern, even though some four body channels are opened ($C_3H_3 + H_2 + 2 H$, $C_2H_2 + CH_3 + 2 H$). Other authors consider also four body breakup patterns in their analysis: CD_3CDO^+ , [3] CD_3CND^+ , [4] and CH_2CHCNH^+ . [5] In the present treatment, we favored an exhaustive treatment and we enabled four body breakups when possible.

H₂ vs. 2H. An empirical rule appears throughout the database of branching ratios: the loss of 2 H atoms is more – often much more – probable than the loss of an H₂ molecule. The only measured exception is NH_4^+ . This rule is used to reduce uncertainty by nesting both pathways using a Dirichlet distribution.

Heavy fragments as a basis for nesting. A corollary of the previous rule is that we are often induced to nest together pathways involving the same heavy fragment (e.g. $X + 2H$ and $X + H_2$). For hydrocarbons, and in absence of experimental values, we adopted this as a general guideline to ensure a balanced treatment amongst those heavy fragments. This rule becomes ambiguous for N-bearing molecules and was restricted to hydrocarbon ions.

Spin states. Some species in the photochemical model have specified spin states. This is mostly the case for CH_2 and N. When the spin states of DR products have not been measured, we nest them inside a uniform Dirichlet subtree.

An overall view of the database content is provided in Fig. 1




Figure 1: Representation of the fragmentation patterns for all ions treated in the present dataset. The line width is proportional to the production rate of the corresponding neutral, normalized to the rate constant. The colors depict the number of heavy atoms in the original ion.

Table 1: DR database

	reaction	$\alpha_0(cm^3.s^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
$H^+ + e^-$	$\rightarrow H$	1.6×10^{-6} [this work]	2 [this work]	$0.5 - 1.5$ [this work]	-	-
$H_2^+ + e^-$	$\rightarrow 2H$	1.6×10^{-6} [1]	2 [this work]	0.43 [1]	1 [this work]	-
$H_3^+ + e^-$	$\rightarrow H_3$	7.2×10^{-8} [6-8]	1.15 [6-8]	$0.48 - 1.0$ [1, 9-11]	-	$0.00 - 0.09$ [1]
	$\rightarrow H_2 + H$					$0.25 - 0.40$ [1, 10]
	$\rightarrow 3H$					$0.51 - 0.75$ [1, 10]
$C^+ + e^-$	$\rightarrow C + h\nu$	4.67×10^{-12} [12, 13]	1.5 [13]	0.6 [13]	0.5 [Woodall et al. [13]]	-
$CH^+ + e^-$	$\rightarrow C + H$	$0.97 - 3.1 \times 10^{-7}$ [1, 13-15]		$0.37 - 0.89$ [1, 13-15]	-	-
$CH_2^+ + e^-$	$\rightarrow CH + H$	$2.5 - 8.7 \times 10^{-7}$ [14, 16]	-	$0.50 - 0.60$ [14, 16]	-	0.25 ± 0.04 [16]
	$\rightarrow C + H_2$					0.12 ± 0.02 [16]
	$\rightarrow C + 2H$					0.63 ± 0.06 [16]
$CH_3^+ + e^-$	$\rightarrow \begin{cases} {}^1CH_2 + H \\ {}^3CH_2 + H \end{cases}$	$3.56 - 12.0 \times 10^{-7}$ [1, 14, 15]	-	0.51 [1, 14, 15]	0.04 [1, 14, 15]	0.40 ± 0.1 [17]
	$\rightarrow CH + H_2$					0.14 ± 0.1 [17]
	$\rightarrow CH + 2H$					0.16 ± 0.15 [17]
	$\rightarrow C + H_2 + H$					0.30 ± 0.08 [17]
$CH_4^+ + e^-$	$\rightarrow \begin{cases} CH_3 + H \\ \begin{cases} {}^1CH_2 + H_2 \\ {}^3CH_2 + H_2 \end{cases} \\ \begin{cases} {}^1CH_2 + 2H \\ {}^3CH_2 + 2H \end{cases} \\ \begin{cases} CH + H_2 + H \\ CH + 3H \end{cases} \\ C + 2H_2 \end{cases}$	$3.5 - 12.8 \times 10^{-7}$ [1, 14, 15]	-	0.51 [1, 14, 15]	0.04 [1, 14, 15]	$0 - 1$ [this work]

Table 1: DR database *cont'd*

reaction	$\alpha_0(cm^3.s^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
$CH_3^+ + e^-$					
$\rightarrow CH_4 + H$	$2.8 - 14 \times 10^{-7}$ [1, 14, 15, 18-22]	-	$0.3 - 1$ ($T_e \leq 300$ K); 1.5 ($T_e \geq 300$ K) [1, 14, 15, 18, 21, 22]	1 ($T_e \geq 300$ K) [this work]	0.049 ± 0.013 [21]; 0.95 ± 0.05 [23]
$\rightarrow CH_3 + H_2$ (a)					0.048 ± 0.002 [21]; (a) + (b) ≤ 0.08 [23]
$\rightarrow CH_3 + 2H$ (b)					0.698 ± 0.008 [21]; (a) + (b) ≤ 0.08 [23]
$\rightarrow \begin{cases} {}^1CH_2 + H_2 + H \\ {}^3CH_2 + H_2 + H \end{cases}$					0.172 ± 0.016 [21]; ≤ 0.01 [23]
$\rightarrow CH + 2H_2$					0.033 ± 0.011 [21]; ≤ 0.01 [23]
$C_2H^+ + e^-$					
$\rightarrow C_2 + H$	2.7×10^{-7} [1, 24]	1.26 [24]	0.76 [1, 15, 24]	0.33 [1, 15, 24]	0.43 ± 0.03 [24]
$\rightarrow CH + C$					0.39 ± 0.04 [24]
$\rightarrow 2C + H$					0.18 ± 0.04 [24]
$C_2H_2^+ + e^-$					
$\rightarrow C_2H + H$	$2.7 - 10 \times 10^{-7}$ [1, 15]	-	$0.5 - 1$ [1, 15]	-	0.50 ± 0.06 [25]
$\rightarrow C_2 + 2H$					0.30 ± 0.05 [25]
$\rightarrow C_2 + H_2$					0.02 ± 0.03 [25]
$\rightarrow \begin{cases} {}^1CH_2 + C \\ {}^3CH_2 + C \end{cases}$					0.05 ± 0.01 [25]
$\rightarrow 2CH$					0.13 ± 0.01 [25]
$C_2H_3^+ + e^-$					
$\rightarrow C_2H_2 + H$	$4.5 - 16 \times 10^{-7}$ [1, 15, 26, 27]	-	0.84 ($T_e \leq 800$ K) [1, 15, 26, 27]; $1.38 - 1.5$ ($T_e \geq 800$ K) [26, 27]	0.3 ($T < 800$ K) [1, 15, 26, 27]	0.29 ± 0.04 [26]
$\rightarrow C_2H + H_2$					0.06 ± 0.03 [26]
$\rightarrow C_2H + 2H$					0.584 ± 0.06 for $\sum = 1$ (0.59) [26]
$\rightarrow C_2 + H + H_2$					0.03 ± 0.01 [26]
$\rightarrow CH_3 + C$					0.006 ± 0.002 [26]
$\rightarrow \begin{cases} {}^1CH_2 + CH \\ {}^3CH_2 + CH \end{cases}$					0.03 ± 0.01 [26]

Table 1: DR database *cont'd*

	reaction	$\alpha_0(cm^3 \cdot s^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
$C_2H_4^+ + e^-$	$\rightarrow C_2H_3 + H$	5.6×10^{-7} [24]	1.11 [24]	0.76 [24, 27]	1 [this work]	0.11 ± 0.07 [24]
	$\rightarrow C_2H_2 + H_2$					0.06 ± 0.03 [24]
	$\rightarrow C_2H_2 + 2H$					0.66 ± 0.06 [24]
	$\rightarrow C_2H + H_2 + H$					0.10 ± 0.04 [24]
	$\rightarrow CH_4 + C$					0.01 ± 0.01 [24]
	$\rightarrow CH_3 + CH$					0.02 ± 0.02 [24]
	$\rightarrow \begin{cases} {}^1CH_2 + {}^1CH_2 \\ {}^1CH_2 + {}^3CH_2 \\ {}^3CH_2 + {}^3CH_2 \end{cases}$					0.04 ± 0.02 [24]
$C_2H_5^+ + e^-$	$\rightarrow C_2H_4 + H$	8.35×10^{-7} [1, 19, 20, 22]	1.30 [1, 19, 20, 22]	0.8 ($T_e \leq 300$ K) [22]; 1.2 ($T_e \geq 300$ K) [22]	1 [this work]	0.12 ± 0.03 [28]
	$\rightarrow C_2H_3 + 2H$					0.27 ± 0.04 [28]
	$\rightarrow C_2H_2 + 3H$					0.13 ± 0.03 [28]
	$\rightarrow C_2H_2 + H_2 + H$					0.30 ± 0.03 for $\sum = 1$ (0.29) [28]
	$\rightarrow \begin{cases} CH_3 + {}^1CH_2 \\ CH_3 + {}^3CH_2 \end{cases}$					0.17 ± 0.01 [28]
	$\rightarrow CH_4 + CH$					0.01 ± 0.01 [28]
	$\rightarrow C_3$					0.000 ± 0.001 [27, 29]
	$\rightarrow C_2 + C$					1.000 ± 0.002 [27, 29]
	$\rightarrow 3C$					0.000 ± 0.001 [27, 29]
	$\rightarrow C_3 + H$					0.662 ± 0.015 [1, 15, 30, 31]
$C_3H^+ + e^-$	$\rightarrow C_2H + C$	$0.5 - 3 \times 10^{-6}$ [this work]	-	0.5 - 1.5 [this work]	-	0.338 ± 0.017 [1, 15, 30, 31]
	$\rightarrow C_2 + CH$					0.00 ± 0.01 [1, 15, 30, 31]
	$\rightarrow C_3$					

Table 1: DR database *cont'd*

	reaction	$\alpha_0(cm^3.s^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
$C_3H_2^+ + e^-$	$\rightarrow \left\{ \begin{array}{l} C_3 + H_2 \\ C_3 + 2H \\ C_3H + H \end{array} \right\}$	$0.5 - 3 \times 10^{-6}$ [this work]	-	$0.5 - 1.5$ [this work]	-	0.875 ± 0.017 [1, 15, 30, 31]
	$\rightarrow \left\{ \begin{array}{l} C_2H + CH \\ C_2H + C + H \end{array} \right\}$					0.125 ± 0.021 [1, 15, 30, 31]
$c - C_3H_3^+ + e^-$	$\rightarrow \left\{ \begin{array}{l} C_3H_2 + H \\ C_3H + H_2 \\ C_3H + 2H \\ C_3 + H + H_2 \end{array} \right\}$	8.00×10^{-7} [32]	1.0125 [32]	0.0 [1, 15, 27]	$\sigma_n = 0.7$ [1, 15, 27]	0.907 ± 0.011 [1, 15, 30, 31]
	$\rightarrow \left\{ \begin{array}{l} C_2H_2 + CH \\ C_2H_2 + C + H \end{array} \right\}$					
	$\rightarrow \left\{ \begin{array}{l} C_2H + {}^1CH_2 \\ C_2H + {}^3CH_2 \\ C_2 + CH_3 \end{array} \right\}$					0.093 ± 0.005 [1, 15, 30, 31]
	$\rightarrow \left\{ \begin{array}{l} C_3H_2 + H \\ C_3H + H_2 \\ C_3H + 2H \\ C_3 + H + H_2 \\ C_2H_2 + CH \\ C_2H_2 + C + H \end{array} \right\}$	1.15×10^{-7} [1, 15, 27, 32]	1.17 [32]	0.0 [1, 15, 27]	$\sigma_n = 0.2$ [1, 15, 27]	0.907 ± 0.011 [1, 15, 30, 31]
$1 - C_3H_3^+ + e^-$	$\rightarrow \left\{ \begin{array}{l} C_2H + {}^1CH_2 \\ C_2H + {}^3CH_2 \\ C_2 + CH_3 \end{array} \right\}$					0.093 ± 0.005 [1, 15, 30, 31]
	$\rightarrow C_3H_3 + H$					
$C_3H_4^+ + e^-$	$\rightarrow C_3H_3 + H$	2.95×10^{-6} [33]	1.034 [33]	0.67 [33]	0.03 [33]	0.87 ± 0.04 [1, 15, 30, 31, 33]
	$\rightarrow C_3H_2 + H_2$					≤ 0.02 [1, 15, 30, 31, 33]
	$\rightarrow C_3H_2 + 2H$					≤ 0.05 [1, 15, 30, 31, 33]
	$\rightarrow C_2H_3 + CH$					0.01 ± 0.01 [1, 15, 30, 31, 33]

Table 1: DR database *cont'd*

reaction	$\alpha_0(cm^3.s^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
$\rightarrow \left\{ \begin{array}{l} C_2H_2 + {}^1CH_2 \\ C_2H_2 + {}^3CH_2 \end{array} \right\}$ $\rightarrow C_2H + CH_3$					0.06 ± 0.02 [1, 15, 30, 31, 33] 0.01 ± 0.01 [1, 15, 30, 31, 33]
$C_3H_5^+ + e^-$ $\rightarrow \left\{ \begin{array}{l} C_3H_4 + H \\ \left\{ \begin{array}{l} C_3H_3 + H_2 \\ C_3H_3 + 2H \end{array} \right\} \\ C_3H_2 + H_2 + H \\ C_3H + 2H_2 \\ C_2H_5 + C \\ C_2H_4 + CH \\ \left\{ \begin{array}{l} C_2H_3 + {}^1CH_2 \\ C_2H_3 + {}^3CH_2 \end{array} \right\} \\ \left\{ \begin{array}{l} C_2H_2 + {}^1CH_2 + H \\ C_2H_2 + {}^3CH_2 + H \end{array} \right\} \\ C_2H_2 + CH + H_2 \\ C_2H_2 + CH_3 \\ C_2H + CH_3 + H \\ C_2H + CH_4 \end{array} \right\}$	$0.5 - 3 \times 10^{-6}$ [this work]	-	$0.5 - 1.5$ [this work]	-	0.867 ± 0.004 [1, 15, 30, 31]
$C_3H_6^+ + e^-$ $\rightarrow \left\{ \begin{array}{l} C_3H_5 + H \\ C_3H_4 + 2H \\ C_3H_3 + H_2 + H \\ \left\{ \begin{array}{l} C_3H_2 + 2H_2 \\ C_3H_2 + H_2 + 2H \end{array} \right\} \end{array} \right\}$	$0.5 - 3 \times 10^{-6}$ [this work]	-	$0.5 - 1.5$ [this work]	-	0.693 ± 0.016 [1, 15, 30, 31]

Table 1: DR database *cont'd*

	reaction	$\alpha_0(cm^3.s^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
	$\left\{ \begin{array}{l} C_2H_5 + CH \\ \left\{ \begin{array}{l} C_2H_4 + {}^1CH_2 \\ C_2H_4 + {}^3CH_2 \end{array} \right\} \\ C_2H_4 + CH + H \\ C_2H_4 + C + H_2 \\ \left\{ \begin{array}{l} C_2H_3 + CH + H_2 \\ C_2H_3 + CH_3 \end{array} \right\} \\ \left\{ \begin{array}{l} C_2H_2 + CH_4 \\ \left\{ \begin{array}{l} C_2H_2 + {}^1CH_2 + H_2 \\ C_2H_2 + {}^3CH_2 + H_2 \end{array} \right\} \\ C_2H_2 + CH_3 + H \\ C_2H + CH_4 + H \\ C_2H + CH_3 + H_2 \end{array} \right\} \\ \rightarrow C + CH_2 + CH_4 \end{array} \right\} \rightarrow$					0.307 ± 0.013 [1, 15, 30, 31]
						0.00 ± 0.01 [1, 15, 30, 31]
$C_3H_7^+ + e^-$	$\begin{array}{l} \rightarrow C_3H_6 + H \\ \rightarrow C_3H_5 + H_2 \\ \rightarrow C_3H_5 + 2H \\ \rightarrow C_3H_4 + H_2 + H \\ \rightarrow C_2H_4 + CH_3 \\ \rightarrow C_2H_3 + CH_4 \\ \rightarrow C_2H_3 + CH_3 + H \\ \rightarrow C_2H_2 + CH_4 + H \\ \rightarrow C_2H_2 + CH_3 + H_2 \end{array}$	$0.5 - 3 \times 10^{-6}$ [2, 19, 34]	-	0.68 [2]	0.015 [2]	0.13 ± 0.05 [34] 0.12 ± 0.05 [34] 0.22 ± 0.08 [34] 0.09 ± 0.02 [34] 0.03 ± 0.02 [34] 0.02 ± 0.02 [34] 0.15 ± 0.04 [34] 0.03 ± 0.03 [34] 0.21 ± 0.04 [34]
$C_3H_8^+ + e^-$	$\begin{array}{l} \rightarrow \left\{ \begin{array}{l} C_3H_7 + H \\ \left\{ \begin{array}{l} C_3H_6 + H_2 \\ C_3H_6 + 2H \end{array} \right\} \\ C_3H_5 + H_2 + H \\ \left\{ \begin{array}{l} C_3H_4 + 2H_2 \\ C_3H_4 + H_2 + 2H \end{array} \right\} \end{array} \right\} \rightarrow$	$0.5 - 3 \times 10^{-6}$ [this work]	-	$0.5 - 1.5$ [this work]	-	0.679 ± 0.029 [1, 15, 30, 31]

Table 1: DR database *cont'd*

reaction	$\alpha_0(cm^3.s^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
$\rightarrow \left\{ \begin{array}{l} C_3H_2 + C \\ C_3H + CH \\ \left\{ \begin{array}{l} C_3 + ^1CH_2 \\ C_3 + ^3CH_2 \end{array} \right\} \\ C_2H_2 + C_2 \\ 2C_2H \end{array} \right\}$					0.040 ± 0.016 [1, 31, 35]
$\rightarrow \left\{ \begin{array}{l} C_2H_2 + C_2 \\ 2C_2H \end{array} \right\}$					0.177 ± 0.015 [1, 31, 35]
$C_4H_3^+ + e^-$	6.2×10^{-7} [1, 15, 36]	2 [this work]	$0.5 - 1.5$ [this work]	-	0.759 ± 0.005 for $\sum = 1$ (0.760) [1, 31, 35]
$\rightarrow \left\{ \begin{array}{l} C_4H_2 + H \\ \left\{ \begin{array}{l} C_4H + H_2 \\ C_4H + 2H \end{array} \right\} \\ C_4 + H + H_2 \\ C_3H_3 + C \\ C_3H_2 + CH \\ \left\{ \begin{array}{l} C_3H + ^1CH_2 \\ C_3H + ^3CH_2 \end{array} \right\} \\ C_3 + CH_3 \\ C_2H_3 + C_2 \\ C_2H_2 + C_2H \end{array} \right\}$					0.063 ± 0.004 [1, 31, 35]
$\rightarrow \left\{ \begin{array}{l} C_4H_3 + H \\ \left\{ \begin{array}{l} C_4H_2 + H_2 \\ C_4H_2 + 2H \end{array} \right\} \\ C_4H + H_2 + H \\ C_3H_4 + C \\ C_3H_3 + CH \\ \left\{ \begin{array}{l} C_3H_2 + ^1CH_2 \\ C_3H_2 + ^3CH_2 \end{array} \right\} \\ C_3H_2 + C + H_2 \\ C_3H + CH_3 \\ \left\{ \begin{array}{l} C_3 + CH_4 \\ C_3 + CH_3 + H \end{array} \right\} \end{array} \right\}$	$0.5 - 3 \times 10^{-6}$ [this work]	-	$0.5 - 1.5$ [this work]	-	0.766 ± 0.027 [1, 31, 35]
$C_4H_4^+ + e^-$					0.063 ± 0.02 [1, 31, 35]

Table 1: DR database *cont'd*

	reaction	$\alpha_0 (cm^3 s^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
	$\left\{ \begin{array}{l} C_2H_4 + C_2 \\ C_2H_3 + C_3H \\ \left\{ \begin{array}{l} 2 C_3H_2 \\ C_2H_2 + C_3H + H \\ C_2H_2 + C_2 + H_2 \end{array} \right\} \\ 2 C_3H + H_2 \end{array} \right\} \rightarrow$					0.171 \pm 0.021 [1, 31, 35]
$C_4H_5^+ + e^-$	$\left\{ \begin{array}{l} C_4H_4 + H \\ \left\{ \begin{array}{l} C_4H_3 + H_2 \\ C_4H_3 + 2 H \end{array} \right\} \\ C_4H_2 + H_2 + H \\ C_4H + 2 H_2 \\ C_3H_5 + C \\ C_3H_4 + CH \\ \left\{ \begin{array}{l} C_3H_3 + {}^1CH_2 \\ C_3H_3 + {}^3CH_2 \end{array} \right\} \\ C_3H_2 + CH_3 \\ C_3H + CH_4 \\ C_2H_5 + C_2 \\ C_2H_4 + C_2H \\ C_2H_3 + C_2H_2 \\ \left\{ \begin{array}{l} 2 C_2H_2 + H \\ C_2H_2 + C_2H + H_2 \end{array} \right\} \end{array} \right\} \rightarrow$	8.2×10^{-7} [37]	1.30 [37]	0.5 – 1.5 [this work]	-	0.460 \pm 0.054 [1, 31, 35]
						0.093 \pm 0.021 [1, 31, 35]
						0.447 \pm 0.053 [1, 31, 35]
$C_4H_6^+ + e^-$	$\left\{ \begin{array}{l} C_4H_5 + H \\ \left\{ \begin{array}{l} C_4H_4 + H_2 \\ C_4H_4 + 2 H \end{array} \right\} \\ C_4H_3 + H_2 + H \\ \left\{ \begin{array}{l} C_4H_2 + 2 H_2 \\ C_4H_2 + H_2 + 2 H \end{array} \right\} \end{array} \right\} \rightarrow$	$0.5 - 3 \times 10^{-6}$ [this work]	-	0.5 – 1.5 [this work]	-	0.589 \pm 0.035 [1, 31, 35]

Table 1: DR database *cont'd*

reaction	$\alpha_0 (cm^3 s^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
$\left\{ \begin{array}{l} C_3H_6 + C \\ C_3H_5 + CH \\ \left\{ \begin{array}{l} C_3H_4 + {}^1CH_2 \\ C_3H_4 + {}^3CH_2 \end{array} \right\} \\ C_3H_4 + C + H_2 \\ \left\{ \begin{array}{l} C_3H_3 + CH_3 \\ C_3H_3 + CH + H_2 \end{array} \right\} \\ \left\{ \begin{array}{l} C_3H_2 + CH_4 \\ C_3H_2 + CH_3 + H \\ \left\{ \begin{array}{l} C_3H_2 + {}^1CH_2 + H_2 \\ C_3H_2 + {}^3CH_2 + H_2 \end{array} \right\} \\ \left\{ \begin{array}{l} C_3H + CH_4 + H \\ C_3H + CH_3 + H_2 \end{array} \right\} \end{array} \right\}$	\rightarrow				$0.090 \pm 0.015 [1, 31, 35]$
$\left\{ \begin{array}{l} C_2H_6 + C_2 \\ C_2H_5 + C_2H \\ \left\{ \begin{array}{l} C_2H_4 + C_2H + H \\ C_2H_4 + C_2H_2 \\ C_2H_4 + C_2 + H_2 \end{array} \right\} \\ \left\{ \begin{array}{l} C_2H_3 + C_2H_2 + H \\ C_2H_3 + C_2H + H_2 \end{array} \right\} \\ C_2H_2 + CH + CH_3 \\ C_2H_2 + C + CH_4 \end{array} \right\}$	\rightarrow				$0.321 \pm 0.034 [1, 31, 35]$
$\left\{ \begin{array}{l} C_4H_6 + H \\ \left\{ \begin{array}{l} C_4H_5 + H_2 \\ C_4H_5 + 2H \end{array} \right\} \\ C_4H_4 + H_2 + H \\ C_4H_3 + 2H_2 \end{array} \right\}$	\rightarrow	$0.5 - 3 \times 10^{-6}$ [this work]	$0.5 - 1.5$ [this work]	-	$0.198 \pm 0.044 [1, 31, 35]$

 $C_4H_7^+ + e^-$

Table 1: DR database *cont'd*

reaction	$\alpha_0 (cm^3 s^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
$\left\{ \begin{array}{l} C_3H_7 + C \\ C_3H_6 + CH \\ \left\{ \begin{array}{l} C_3H_5 + {}^1CH_2 \\ C_3H_5 + {}^3CH_2 \end{array} \right\} \\ \left\{ \begin{array}{l} C_3H_4 + CH_3 \\ \left\{ \begin{array}{l} C_3H_4 + {}^1CH_2 + H \\ C_3H_4 + {}^3CH_2 + H \end{array} \right\} \\ C_3H_4 + CH + H_2 \end{array} \right\} \\ \left\{ \begin{array}{l} C_3H_3 + CH_4 \\ C_3H_3 + CH_3 + H \end{array} \right\} \\ \left\{ \begin{array}{l} C_3H_3 + {}^1CH_2 + H_2 \\ C_3H_3 + {}^3CH_2 + H_2 \end{array} \right\} \\ \left\{ \begin{array}{l} C_3H_2 + CH_4 + H \\ C_3H_2 + CH_3 + H_2 \end{array} \right\} \\ C_3H + CH_4 + H_2 \end{array} \right\}$	\rightarrow				0.655 ± 0.059 [1, 31, 35]
$\left\{ \begin{array}{l} C_2H_6 + C_2H \\ C_2H_5 + C_2H_2 \\ \left\{ \begin{array}{l} C_2H_4 + C_2H_2 + H \\ C_2H_4 + C_2H + H_2 \end{array} \right\} \\ \left\{ \begin{array}{l} 2C_2H_3 + H \\ C_2H_3 + C_2H_2 + H_2 \end{array} \right\} \\ C_2H_2 + {}^1CH_2 + CH_3 \\ C_2H_2 + {}^3CH_2 + CH_3 \end{array} \right\}$	\rightarrow				0.147 ± 0.034 [1, 31, 35]
$\left\{ \begin{array}{l} C_4H_7 + H \\ \left\{ \begin{array}{l} C_4H_6 + H_2 \\ C_4H_6 + 2H \end{array} \right\} \\ \left\{ \begin{array}{l} C_4H_5 + H_2 + H \\ C_4H_5 + 3H \end{array} \right\} \\ \left\{ \begin{array}{l} C_4H_4 + 2H_2 \\ C_4H_4 + H_2 + 2H \end{array} \right\} \end{array} \right\}$	\rightarrow				0.00 ± 0.01 [1, 31, 35]
$C_4H_8^+ + e^-$	\rightarrow	$0.5 - 3 \times 10^{-6}$ [this work]	$0.5 - 1.5$ [this work]	-	0.305 ± 0.031 [1, 31, 35]

Table 1: DR database *cont'd*

	reaction	$\alpha_0(\text{cm}^3 \cdot \text{s}^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
	$\left\{ \begin{array}{l} \left\{ \begin{array}{l} \text{C}_2\text{H}_4 + \text{CH} + \text{CH}_3 \\ \text{C}_2\text{H}_4 + \text{C} + \text{CH}_4 \end{array} \right\} \\ \text{C}_2\text{H}_3 + \text{CH} + \text{CH}_4 \\ \left\{ \begin{array}{l} \text{C}_2\text{H}_2 + 2\text{CH}_3 \\ \left\{ \begin{array}{l} \text{C}_2\text{H}_2 + {}^1\text{CH}_2 + \text{CH}_4 \\ \text{C}_2\text{H}_2 + {}^3\text{CH}_2 + \text{CH}_4 \end{array} \right\} \\ \text{C}_2\text{H} + \text{CH}_4 + \text{CH}_3 \\ \text{C}_2 + 2\text{CH}_4 \end{array} \right\}$					0.000 \pm 0.010 [1, 31, 35]
$\text{C}_4\text{H}_9^+ + \text{e}^-$	$\left\{ \begin{array}{l} \left\{ \begin{array}{l} \text{C}_4\text{H}_8 + \text{H} \\ \left\{ \begin{array}{l} \text{C}_4\text{H}_7 + \text{H}_2 \\ \text{C}_4\text{H}_7 + 2\text{H} \end{array} \right\} \\ \text{C}_4\text{H}_6 + \text{H}_2 + \text{H} \\ \left\{ \begin{array}{l} \text{C}_4\text{H}_5 + 2\text{H}_2 \\ \text{C}_4\text{H}_5 + \text{H}_2 + 2\text{H} \end{array} \right\} \\ \text{C}_3\text{H}_8 + \text{CH} \\ \left\{ \begin{array}{l} \text{C}_3\text{H}_7 + {}^1\text{CH}_2 \\ \text{C}_3\text{H}_7 + {}^3\text{CH}_2 \end{array} \right\} \\ \left\{ \begin{array}{l} \text{C}_3\text{H}_6 + \text{CH}_3 \\ \left\{ \begin{array}{l} \text{C}_3\text{H}_6 + {}^1\text{CH}_2 + \text{H} \\ \text{C}_3\text{H}_6 + {}^3\text{CH}_2 + \text{H} \end{array} \right\} \\ \text{C}_3\text{H}_6 + \text{CH} + \text{H}_2 \\ \left\{ \begin{array}{l} \text{C}_3\text{H}_5 + \text{CH}_4 \\ \text{C}_3\text{H}_5 + \text{CH}_3 + \text{H} \end{array} \right\} \\ \left\{ \begin{array}{l} \text{C}_3\text{H}_5 + {}^1\text{CH}_2 + \text{H}_2 \\ \text{C}_3\text{H}_5 + {}^3\text{CH}_2 + \text{H}_2 \end{array} \right\} \\ \left\{ \begin{array}{l} \text{C}_3\text{H}_4 + \text{CH}_4 + \text{H} \\ \text{C}_3\text{H}_4 + \text{CH}_3 + \text{H}_2 \end{array} \right\} \\ \text{C}_3\text{H}_3 + \text{CH}_4 + \text{H}_2 \end{array} \right\}$	8.3×10^{-7} [19]	1.30 [19]	0.5 – 1.5 [this work]	-	0.574 \pm 0.041 for $\sum = 1$ (0.575) [1, 31, 35, 38]
						0.411 \pm 0.038 [1, 31, 35, 38]

Table 1: DR database *cont'd*

	reaction	$\alpha_0(cm^3.s^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
	$\rightarrow \left\{ \begin{array}{l} \left\{ \begin{array}{l} C_2H_6 + C_2H_3 \\ C_2H_6 + C_2H_2 + H \\ C_2H_6 + C_2H + H_2 \end{array} \right\} \\ \left\{ \begin{array}{l} C_2H_5 + C_2H_4 \\ C_2H_5 + C_2H_3 + H \\ C_2H_5 + C_2H_2 + H_2 \end{array} \right\} \\ 2 C_2H_4 + H \end{array} \right\}$					0.015 \pm 0.026 [1, 31, 35, 38]
	$\rightarrow \left\{ \begin{array}{l} C_2H_4 + CH + CH_4 \\ \left\{ \begin{array}{l} C_2H_4 + {}^1CH_2 + CH_3 \\ C_2H_4 + {}^3CH_2 + CH_3 \end{array} \right\} \\ C_2H_3 + 2CH_3 \\ \left\{ \begin{array}{l} C_2H_3 + {}^1CH_2 + CH_4 \\ C_2H_3 + {}^3CH_2 + CH_4 \end{array} \right\} \\ C_2H_2 + CH_3 + CH_4 \end{array} \right\}$					0.000 \pm 0.010 [1, 31, 35, 38]
$C_3H_3^+ + e^-$	\rightarrow unknown	9×10^{-7} [39]	1.30 [39]	0.5 – 1.5[work]	-	-
$C_3H_9^+ + e^-$	\rightarrow unknown	9.5×10^{-7} [37]	1.32 [37]	0.5 – 1.5[work]	-	-
$C_6H_4^+ + e^-$	\rightarrow unknown	1.10×10^{-6} [37]	1.30 [37]	0.5 – 1.5[work]	-	-
$C_6H_5^+ + e^-$	\rightarrow unknown	1.10×10^{-6} [37]	1.30 [37]	0.5 – 1.5[work]	-	-
$C_6H_6^+ + e^-$	\rightarrow unknown	1.0×10^{-6} [39]	1.33 [39]	0.5 – 1.5[work]	-	-
$C_6H_7^+ + e^-$	\rightarrow unknown	2.4×10^{-6} [22]	1.37 [22]	0.5 – 1.5 [this work] ($T \leq T_1$); 1.3 ($T \geq T_1$) [22]; $T_1 \leq 300$ K	1 [this work]	-
$C_7H_5^+ + e^-$	\rightarrow unknown	7.0×10^{-7} [39]	1.29 [39]	0.5 – 1.5[work]	-	-

Table 1: DR database *cont'd*

	reaction	$\alpha_0(cm^3.s^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
$N^+ + e^-$	$\rightarrow N + h\nu$	4.0×10^{-12} [12, 13]	1.50 [12, 13]	0.58 [12, 13]	0.5 [12, 13]	-
$NH^+ + e^-$	$\rightarrow N + H$	4.33×10^{-8} [1, 13, 15]	2 [this work]	0.5 – 1.5 [this work]	-	-
$NH_2^+ + e^-$	$\rightarrow N + H_2$	$0.5 - 10 \times 10^{-7}$ [this work]	-	0.5 – 1.5 [this work]	-	0.04 \pm 0.03 [40, 41]
	$\rightarrow NH + H$					0.38 \pm 0.06 [40, 41]
	$\rightarrow \left\{ \begin{array}{l} (^4S)N + 2H \\ (^2D)N + 2H \\ (^2P)N + 2H \end{array} \right\}$					0.58 ± 0.09 $\left\{ \begin{array}{l} 0.53 \pm 0.04 \\ 0.45 \pm 0.05 \end{array} \right\}$ [40, 41] 0.02 \pm 0.02
$NH_3^+ + e^-$	$\rightarrow \left\{ \begin{array}{l} NH_2 + H \\ \left\{ \begin{array}{l} NH + H_2 \\ NH + 2H \end{array} \right\} \\ \left\{ \begin{array}{l} N + H_2 + H \\ N + 3H \end{array} \right\} \end{array} \right\}$	3.1×10^{-7} [13]	2 [this work]	0.5 – 1.5 [this work]	-	0-1 [this work]
$NH_4^+ + e^-$	$\rightarrow NH_3 + H$	$9 - 15 \times 10^{-7}$ [1, 15, 42-44]	-	0.75 [1, 15, 43, 44]	0.2 [1, 15, 43, 44]	0.85 \pm 0.04 [27, 40, 44, 45]
	$\rightarrow NH_2 + H_2$					0.2 \pm 0.02 [27, 40, 44, 45]
	$\rightarrow NH_2 + 2H$					0.13 \pm 0.01 [27, 40, 44, 45]
	$\rightarrow NH + H + H_2$					0.00 \pm 0.01 [27, 40, 44, 45]
	$\rightarrow N + 2H_2$					0.00 \pm 0.01 [27, 40, 44, 45]
$N_2^+ + e^-$	$\rightarrow (^4S)N + (^2D)N$	2.2×10^{-7} [46-53]	1.25 [46-53]	0.35 [46, 47, 49, 50, 53]	0.15 [46, 47, 49, 50, 53]	0.42 \pm 0.10 [53, 54]
	$\rightarrow (^4S)N + (^2P)N$					0.09 \pm 0.09 [53, 54]
	$\rightarrow (^2D)N + (^2D)N$					0.49 \pm 0.10 [53, 54]
$N_2H^+ + e^-$	$\rightarrow N_2 + H$	2×10^{-7} [1, 15, 18, 55-60]	1.5 [1, 15, 18, 55-60]	0.5 [55, 58-60]	0.5 [55, 58-60]	0.95-1 [18, 23, 61]

Table 1: DR database *cont'd*

	reaction	$\alpha_0(cm^3.s^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
	$\rightarrow NH + N$					0-0.05 [18, 23, 61]
$CN^+ + e^-$	$\rightarrow (^4S)N + (^3P)C$	3.4×10^{-7} [62]	1.16 [62]	0.55 [27, 62]	1 [this work]	≤ 0.018 [62]
	$\rightarrow (^4S)N + (^1D)C$					0.038 ± 0.028 [62]
	$\rightarrow \left\{ \begin{array}{l} (^2D)N + (^3P)C \\ (^4S)N + (^1S)C \end{array} \right\}$					0.142 ± 0.014 [62]
	$\rightarrow \left\{ \begin{array}{l} (^2P)N + (^3P)C \\ (^2D)N + (^1D)C \end{array} \right\}$					0.561 ± 0.026 [62]
	$\rightarrow \left\{ \begin{array}{l} (^2P)N + (^1D)C \\ (^2D)N + (^1S)C \end{array} \right\}$					0.255 ± 0.014 [62]
	$\rightarrow (^2P)N + (^1S)C$					≤ 0.014 [62]
	$\rightarrow CN + H$	2.5×10^{-6} [63]	2 [this work]	1 [63]	0.02 [63, 64]	-
	$\rightarrow CN + H$	1.3×10^{-7} [63, 64]	1.7 [63, 64]	0.98 [63, 64]	0.02 [63, 64]	-
$HCN^+ + e^-$	$\rightarrow HCN + H$	3.2×10^{-7} [1, 65-67]	1.25 [1, 65-67]	$0.65 - 1.38$ [65, 67]	-	$0.26-0.41$ [65, 68, 69]
	$\rightarrow HNC + H$					$0.29-0.39$ [65, 68, 69]
	$\rightarrow CN + H_2$					0 ± 0.017 [65]
	$\rightarrow CN + 2H$					0.325 ± 0.032 [65]
$CH_2NH_2^+ + e^-$	$\rightarrow \left\{ \begin{array}{l} CH_3N + H \\ ^1CH_2 + NH_2 \\ ^3CH_2 + NH_2 \end{array} \right\}$	$[0.5 - 3] \times 10^{-6}$ [this work]	-	$0.5 - 1.5$ [this work]	-	$[0 - 1]$ [this work]
	$\rightarrow \left\{ \begin{array}{l} HCN + H_2 + H \\ HNC + H_2 + H \\ CN + 2H_2 \end{array} \right\}$					
$CH_3NH_3^+ + e^-$	\rightarrow unknown	14.0×10^{-7} [1]	2 [this work]	$0.5 - 1.5$ [this work]	-	-

Table 1: DR database *cont'd*

	reaction	$\alpha_0 (cm^3 s^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
$CH_3CNH^+ + e^-$	$\left\{ \begin{array}{l} CH_3CN + H \\ \left\{ \begin{array}{l} C_2H_2N + H_2 \\ C_2H_2N + 2H \end{array} \right\} \\ C_2HN + H_2 + H \end{array} \right\}$	$3.3 \times 10^{-7} [1, 15, 51, 66, 67]$	$1.15 [66, 67]$	$1.03 [67]$	$1 [this\ work]$	$0.65 \pm 0.03, a \in [0.25, 1],$ $b \in [0.025] [4]$
	$\left\{ \begin{array}{l} C_2N + 2H_2 \\ \left\{ \begin{array}{l} C_2H_4 + N \\ C_2H_2 + NH_2 \\ C_2H_2 + NH + H \\ C_2H_2 + N + H_2 \\ C_2H + NH_3 \\ C_2H + NH + H_2 \end{array} \right\} \end{array} \right\}$					
	$\left\{ \begin{array}{l} \left\{ \begin{array}{l} HCN + CH_3 \\ HNC + CH_3 \end{array} \right\} \\ \left\{ \begin{array}{l} HCN + {}^1CH_2 + H \\ HCN + {}^3CH_2 + H \\ HNC + {}^1CH_2 + H \\ HNC + {}^3CH_2 + H \end{array} \right\} \\ HCN + CH + H_2 \\ HNC + CH + H_2 \end{array} \right\}$					
	$\left\{ \begin{array}{l} CN + CH_3 + H \\ CH_3N + CH \\ \left\{ \begin{array}{l} CH_2N + {}^1CH_2 \\ CH_2N + {}^3CH_2 \end{array} \right\} \\ CN + CH_4 \\ \left\{ \begin{array}{l} CN + {}^1CH_2 + H_2 \\ CN + {}^3CH_2 + H_2 \end{array} \right\} \end{array} \right\}$					
	$\rightarrow 2CN$			$0.5 - [this\ work]$	$1.5 [this\ work]$	$- [70]$
	$\rightarrow CN + C_2$			$0.5 - [this\ work]$	$1.5 [this\ work]$	$- [13]$
	$\rightarrow C_3N + H$			$0.5 - [this\ work]$	$1.5 [this\ work]$	$0.44 \pm 0.04 [71]$
$C_2N_2^+ + e^-$		$0.5 - 3 \times 10^{-6} [this\ work]$	-			
$C_3N^+ + e^-$		$0.5 - 3 \times 10^{-6} [this\ work]$	-			
$CHCCN^+ + e^-$		$0.5 - 3 \times 10^{-6} [this\ work]$	-			

Table 1: DR database *cont'd*

reaction	$\alpha_0(cm^3.s^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
$\rightarrow \begin{cases} C_2H + CN \\ C_2 + H + CN \\ \left\{ \begin{array}{l} HCN + C_2 \\ HNC + C_2 \end{array} \right\} \\ H + C + C_2N \\ \left\{ \begin{array}{l} C_2N + CH \\ N + C_3H \end{array} \right\} \\ HC_2N + C \\ NH + C_3 \end{cases}$					0.48 ± 0.05 [71] 0.04 ± 0.02 [71] 0.02 ± 0.01 [71] 0.02 ± 0.01 [71] 0.00 ± 0.01 [71]
$CHCCNH^+ + e^- \rightarrow \begin{cases} \left\{ \begin{array}{l} HC_3N + H \\ C_3NH + H \end{array} \right\} \\ \left\{ \begin{array}{l} C_2NCH + H \\ HC_2NC + H \end{array} \right\} \\ C_3N + H_2 \\ \left\{ \begin{array}{l} HCN + C_2H \\ HNC + C_2H \end{array} \right\} \\ CN + C_2H_2 \end{cases}$	$0.5 - 3 \times 10^{-6}$ [this work]	-	$0.5 -$ work	1.5 [this work]	0.52 ± 0.05 , $a \geq b$ [71, 72] 0.48 ± 0.05 [71, 72]
$CH_2CHCNH^+ + e^- \rightarrow \begin{cases} C_3H_3N + H \\ \left\{ \begin{array}{l} C_3H_2N + H_2 \\ C_3H_2N + 2H \end{array} \right\} \\ C_3HN + H + H_2 \\ \left\{ \begin{array}{l} C_3N + 2H_2 \\ C_3N + 2H + H_2 \end{array} \right\} \end{cases}$	1.76×10^{-6} [5]	1.17 [5]	0.80 [5]	0.025 [5]	0.50 ± 0.04 [5]

Table 1: DR database *cont'd*

reaction	$\alpha_0 (cm^3 s^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
$\left\{ \begin{array}{l} \text{CNH}_3 + \text{C}_2\text{H} \\ \text{CNH}_2 + \text{C}_2\text{H}_2 \\ \left\{ \begin{array}{l} \text{HCN} + \text{C}_2\text{H}_3 \\ \text{HNC} + \text{C}_2\text{H}_3 \end{array} \right\} \\ \left\{ \begin{array}{l} \text{HCN} + \text{C}_2\text{H}_2 + \text{H} \\ \text{HNC} + \text{C}_2\text{H}_2 + \text{H} \end{array} \right\} \\ \left\{ \begin{array}{l} \text{HCN} + \text{C}_2\text{H} + \text{H}_2 \\ \text{HNC} + \text{C}_2\text{H} + \text{H}_2 \end{array} \right\} \\ \text{CN} + \text{C}_2\text{H}_4 \\ \text{CN} + \text{C}_2\text{H}_2 + \text{H}_2 \\ \text{C}_2\text{H}_3\text{N} + \text{CH} \\ \left\{ \begin{array}{l} \text{C}_2\text{H}_2\text{N} + {}^1\text{CH}_2 \\ \text{C}_2\text{H}_2\text{N} + {}^3\text{CH}_2 \end{array} \right\} \\ \text{C}_2\text{HN} + \text{CH}_3 \\ \text{C}_2\text{N} + \text{CH}_4 \\ \text{C}_2\text{N} + \text{CH}_3 + \text{H} \\ \text{NH}_3 + \text{C}_3\text{H} \\ \text{NH}_2 + \text{C}_3\text{H}_2 \\ \text{NH} + \text{C}_3\text{H}_3 \\ \text{N} + \text{C}_3\text{H}_4 \end{array} \right\} \rightarrow$					0.49 ± 0.04 [5]
$\left\{ \begin{array}{l} \text{C}_2\text{H}_5\text{CNH}^+ + \text{e}^- \\ \rightarrow \text{unknown} \end{array} \right\}$	4.6 × 10 ⁻⁷ [66, 67]	1.15 [66, 67]	0.81 [67]	1 [this work]	-
$\left\{ \begin{array}{l} \text{H}_2\text{O}^+ + \text{e}^- \\ \rightarrow \text{OH} + \text{H} \\ \rightarrow \text{O} + \text{H}_2 \\ \rightarrow \text{O} + 2\text{H} \end{array} \right\}$	3.34 × 10 ⁻⁷ [1, 15, 73, 74]	1.30 [1, 15, 73, 74]	0.77 [1, 15]	0.05 [1, 15]	0.25 ± 0.1 [17, 73–75] 0.10 ± 0.06 [17, 73–75] 0.65 ± 0.1 [17, 73–75]
$\left\{ \begin{array}{l} \text{H}_3\text{O}^+ + \text{e}^- \\ \rightarrow \text{OH} + 2\text{H} \\ \rightarrow \text{H}_2\text{O} + \text{H} \\ \rightarrow \text{OH} + \text{H}_2 \\ \rightarrow \text{O} + \text{H}_2 + \text{H} \end{array} \right\}$	8 × 10 ⁻⁷ [1, 20, 76, 77]	1.5 [1, 20, 76, 77]	0.9 [76, 78]	0.12 [76, 78]	0.60 ± 0.15 [17, 77–80] 0.25 ± 0.05 [17, 77–80] 0.14 ± 0.05 [17, 77–80] 0.01 ± 0.05 [17, 77–80]

Table 1: DR database *cont'd*

reaction	$\alpha_0(cm^3.s^{-1})$	$F_{\alpha_0} = 1 + \frac{u(\alpha_0)}{\alpha_0}$	β	$\frac{u(\beta)}{\beta}$	b
$CO^+ + e^- \rightarrow C + O$	2.0×10^{-7} [51, 81, 82]	1.5 [51, 81, 82]	0.55 [81]	1 [this work]	-
$HCO^+ + e^- \rightarrow CO + H$	2.1×10^{-7} [1, 13, 15, 18, 20, 55, 59, 83-85]	1.41 [1, 13, 15, 18, 20, 55, 59, 83-85]	0.69 [1, 13, 15, 27, 55, 59]	0.45 [1, 13, 15, 27, 55, 59]	0.92 ± 0.03 [45, 59]
$\rightarrow C + OH$					0.07 ± 0.02 [45, 59]
$\rightarrow O + CH$					0.01 ± 0.01 [45, 59]
$CH_2OH^+ + e^- \rightarrow H_2O + CH$	7×10^{-7} [86]	2 [this work]	0.78 [86]	1 [this work]	0.022 ± 0.022 [86]
$\rightarrow CH_2 + OH$					0.055 ± 0.055 [86]
$\left\{ \begin{array}{l} HCOH + H \\ HCO + H_2 \\ HCO + 2H \\ CO + H_2 + H \\ CO + 3H \end{array} \right\} \rightarrow$					0.923 ± 0.1 [86]
$NO^+ + e^- \rightarrow (^4S)N + (^3P)O$	4×10^{-7} [1, 13, 17, 48, 87-89]	1.50 [1, 13, 17, 48, 87-89]	0.8 [1, 13, 89]	0.5 [1, 13, 89]	0.85 ± 0.06 [17, 90]
$\rightarrow \left\{ \begin{array}{l} (^4S)N + (^1D)O \\ (^2D)N + (^3D)O \end{array} \right\}$					0.15 ± 0.06 [17]
$OH^+ + e^- \rightarrow O + H$	3.75×10^{-8} [1, 13, 15, 91]	2 [this work]	0.5 [1, 13, 15, 91]	1 [this work]	-





Figure 2: A normalization to unity or changing only the maximal value has no consequences over the sampled space, as this modification is always smaller than the measurements’ uncertainties. Black dots: C_4H^+ DR branching ratios normalized to unity; blue dots: C_4H^+ DR branching ratios with a recalculated maximal value; red dots: $C_4H_2^+$ DR branching ratios normalized to unity; green dots: $C_4H_2^+$ DR branching ratios with a recalculated maximal value; orange dots: $C_4H_3^+$ DR branching ratios normalized to unity; violet dots: $C_4H_3^+$ DR branching ratios with a recalculated maximal value.

Notes

- Sometimes the sum of the branching ratios found in the literature does not sum to one. In this case, it has been chosen to modify the maximal value. This change is specified when necessary, the reported measured value is given in parenthesis. This choice rather than renormalizing the whole set is actually of no consequence. Indeed, the differences implied by either a renormalization or the change in the maximal value are by the measurements’ uncertainties. (see Fig. 2 for a three dimensions example).
- $HCNH^+$: The available data are for the ratio and the sum of the two species HNC and HCN. The ratio is characterized by Ishii et al. [69] ($0.77 \leq \frac{[HCN]}{[HNC]} \leq 1.32$) and the sum by Semaniak et al. [65] ($[HCN] + [HNC] = 0.675 \pm 0.016$). The values given in the database have been figured out from a Monte Carlo propagation with the former equations, they are in agreement with Hickman et al. [68] calculations, no further spreading of the branching ratios was necessary.
- $l,c-C_3H_3^+$: the measurements of the $C_3H_3^+$ dissociative recombination rate and branching ratios are difficult to translate into a database. McLain et al. [32] give two different global rate constants for the l and c species, but the branching ratios only for the global form $C_3H_3^+$ of the species. Angelova et al. [30] give the branching ratios as follows:
We find that the most probable channel for the recombination of $C_3H_3^+$ is the C_3 channel accounting for 90.7% of the total. This channel corresponds to the loss of a single hydrogen atom or to the loss of a H_2 molecule from the cyclic ion. The linear ion, if recombining, can also lose two separate H atoms. The $C_2 + C$ channel is not negligible, and accounts either for a unique C–C bound break in the linear ion, or for ring opening in the cyclic ion. In this case, two C–C bounds must be broken, or considerable rearrangement must occur prior to splitting.

Florescu-Mitchell and Mitchell [1] translate into actual channels with



We translated it with each isomer giving all the channels, even though this is not what is actually happening.

- There is an actual controversy over CH_5^+ , indeed Molek et al. [23] using a Flowing Afterglow Electron Ionizer (FAEI) found totally different results than the previous storage ring (SR) experiment [21]:

		Molek et al. [23] (FAEI)	Semaniak et al. [21] (SR CRYRING)
$\text{CH}_5^+ + \text{e}^-$	$\rightarrow \text{CH}_4 + \text{H}$	$95 \pm 5\%$	$4.9 \pm 1.3\%$
	$\rightarrow \text{CH}_3 + \text{H}_2$		$4.8 \pm 0.2\%$
	$\rightarrow \text{CH}_3 + 2\text{H}$	$\leq 8\%$	$69.8 \pm 0.8\%$
	$\rightarrow \text{CH}_2 + \text{H} + \text{H}_2$	$\leq 1\%$	$17.2 \pm 1.6\%$
	$\rightarrow \text{CH} + \text{H}_2 + \text{H}_2$	$\leq 1\%$	$3.3 \pm 1.1\%$

They explain the discrepancies by the fact that CH_4 have a stated trend to form CH_3 by DR, and thus their produced CH_4 is consumed by the electrons. The CRYRING values for the branching ratios are taken at a null center-of-mass energy.

The SR scheme was implemented in the present model.

- C_2H^+ , C_2H_2^+ and C_2H_3^+ : Florescu-Mitchell and Mitchell [1], Stockholm University [15] take their rate constant and temperature dependencies from Mul and McGowan [92]. It is to be noted that the α_0 value reported in Stockholm University [15] is the 100 K rate constant value in Mul and McGowan [92] and that the proposed parameters in Florescu-Mitchell and Mitchell [1] do not yield the 100 K value of Mul and McGowan [92].
- CH_4^+ , NH_3^+ : the branching ratios are the combinatorics with a limitation to a three-body breakup. Indeed, three-body breakups are exceptionnal at those collision energies and four-body breakups have been observed at high temperatures. Novotný et al. [93] give thresholds of 3.3 and 5.3 eV for the four-body dissociation channels of the DR of D_3O^+ , which corresponds to electron temperatures of 38,295 K and 61,504 K. These temperatures are largely too high to be considered for this database. No thermodynamics considerations are done, thus a total uncertainty is stated.
- C_2H_5^+ , $\text{CH}_3\text{CNH}_3^+$, CHCCN^+ , CHCCNH^+ : the neutral products and branches are taken as analogous to the deuterated species. Geppert et al. [28] measured the branching ratios for the DR of C_2D_5^+ , Vigren et al. [4] for the DR of CD_3CND^+ , Geppert et al. [71] for DCCCN^+ and DCCCND^+ , which complete Osamura et al. [72] calculation for the DR of CHCCNH^+ . Larsson et al. [34] discuss the comparison of C_3H_7^+ and C_3D_7^+ dissociative recombination branching ratios. The values of the branching ratios are stated as totally uncertain.
- H_3^+ : The values of the rate constant at 300 K for the dissociative recombination of H_3^+ was a very challenging issue, as the measurements would be discrepant up to a factor of 10^5 . In Larsson et al. [8] paper, they give a review of the " H_3^+ -saga", and the most probable value appears to be the calculated value of Kokouline and Greene [6, 7] of $7.2 \times 10^{-8} \text{ cm}^{-3}.\text{s}^{-1}$ at a 15% precision.
- C_3H_7^+ : The value of Ehlerding et al. [2] is considered as certainly too high by Larsson et al. [34], a unique procedure of two-step normalization being the potential cause of this bias. Nonetheless, considering this value with its uncertainty ($1.6 \times 10^{-6} \pm 0.35$ (1σ level) in $\text{cm}^{-3}.\text{s}^{-1}$) in regard of Lehfaoui et al. [19] previous value of $8.7 \times 10^{-7} \text{ cm}^{-3}.\text{s}^{-1}$, given with a 30% uncertainty (no equivalence in σ given). Thus with the hypothesis that the uncertainty of Lehfaoui et al. [19] is at a 2σ level, we found an interval of [4.785–29.5] at a 3σ level in $10^{-7} \text{ cm}^{-3}.\text{s}^{-1}$ units for the elicitation, which justify the use of the interval [5,30] in $10^{-7} \text{ cm}^{-3}.\text{s}^{-1}$ as the maximum interval in case of no information.
- C_6H_6^+ is the benzene isomer only.
- N^+ and C^+ : Nahar and Pradhan [12] calculated the rate constant over a range of temperature, we performed a fit, wich we found to be corresponding to Woodall et al. [13] values from the same fit. The RMS errors we found on our fit are lower than the uncertainties reported in Woodall et al. [13], thus we report here the biggest uncertainty values.
- For unknown α_0 of small molecules - typically small nitriles - it was found more reasonable to establish an upper limit to $10^{-6} \text{ cm}^{-3}.\text{s}^{-1}$ rather than the high value of $3 \times 10^{-6} \text{ cm}^{-3}.\text{s}^{-1}$, suitable for heavy hydrocarbons (see Adams et al. [27], Adams and Babcock [60], Tab. 1 and Fig. 8 resp.), and a lower bound of $5 \times 10^{-8} \text{ cm}^{-3}.\text{s}^{-1}$
- CNC^+ , NH_2^+ , C_2N_2^+ : the trend for the rate constant at 300 K to increase has been observed by Adams et al. [27], Adams and Babcock [60] for hydrocarbons, but as far as small molecules are concerned, the rate constants at 300 K are either smaller or in the 10^{-7} order of magnitude in $\text{cm}^{-3}.\text{s}^{-1}$. We thus chose an interval of $0.5 - 10 \times 10^{-7}$ as the interval of maximum amplitude for small nitriles for which the rate constant at 300 K is unknown, see above point.

- N_2H^+ : Amano [55] provide measurements at 110, 210 and 273 K. A bayesian fit of the form $\alpha(T_e) = \alpha_0 \cdot \left(\frac{T_e}{300}\right)^{-n}$ provides $\alpha_0 = 6.6 \pm 0.4 \times 10^{-7} \text{ cm}^{-3} \cdot \text{s}^{-1}$ and $n = 0.75 \pm 1$ (within the [0.5,1.5] bounds). Adams and Babcock [60], Tab. 2 perform the same calculations and give a similar results ($7.1 \times 10^{-7} \text{ cm}^{-3} \cdot \text{s}^{-1}$). Yet those measurements are made for vibrationnally excited molecules, and as they are very discrepant with other measures, giving a value around $2 \times 10^{-7} \text{ cm}^{-3} \cdot \text{s}^{-1}$, they were not taken into account for this elicitation.
- Geppert et al. [57, 59] give from CRYRING results a branching ratio of *circa* 2/3 for the $\text{NH} + \text{H}$ channels. However, Molek et al. [23, 61] suspect the parasitic reaction $^{15}\text{N}^{14}\text{N} + \text{e}^- \rightarrow ^{15}\text{N} + ^{14}\text{N}$ and a detection of ^{15}N to be the reason for a measurement of so high a value for the NH channel.
- $\text{HCN}^+/\text{HNC}^+$: The notation CHN^+ in Sheehan et al. [64] denotes an unknown mixture of the two isomers HCN^+ and HNC^+ . Sheehan et al. [64] measured a rate constant of the form $\alpha(T_e) = 3.9 \times 10^{-7} \left(\frac{T_e}{300}\right)^{-0.96}$ for CHN^+ DR, and a rate constant of the form $\alpha(T_e) = 1.82 \times 10^{-7} \left(\frac{T_e}{300}\right)^{-0.96}$ for the HNC^+ isomer DR. Talbi et al. [63] calculated rate constants as $\alpha_{\text{HCN}^+}(T_e) = 2.5 \times 10^{-6} \left(\frac{T_e}{300}\right)^{-1}$ and $\alpha_{\text{HNC}^+}(T_e) = 9 \times 10^{-8} \left(\frac{T_e}{300}\right)^{-1}$. Talbi et al. [63] and Sheehan et al. [64] have a factor 2 between the measured and calculated values of the DR of HNC^+ , thus we elicited HCN^+ with the calculated value and a 100% uncertainty.
- C_2N_2^+ , C_2HN_2^+ : Woodall et al. [13] give estimations of the rate constant and branching ratios, we thus considered the given channels and stated a total uncertainties over the values of the rate constants, the branching ratios and the possible isomers (HNC and HCN for C_2HN_2^+ DR).
- CN^+ : the different isomers of the carbon are given, even though the differentiation is not made in the model. We kept the different isomers because of the nested pattern which do not allow to put together the channels with common nitrogen.
- H_3O^+ : Williams et al. [94] provide branching ratios that are in contradiction with the other measurements: they were not taken into account.
- CH_2OH^+ : we have access only to Hamberg et al. [86] abstract, and no thermodynamics data have been found. Thus for the conservation of the C-O bound, it is the combinatorics up to all the possibilities. If the $\text{CO} + 3 \text{ H}$ channel is not thermodynamically possible, it should not be of great consequences: in Vuitton et al. [95] mass spectrum assignation, CH_2OH^+ does not appear, and in a steady state of our model, its density is about $0.25 \pm 0.2 \text{ cm}^{-3}$ at 1100 km (neutral species densities taken from ?).

References

- [1] A.I. Florescu-Mitchell and J.B.A. Mitchell. Dissociative recombination. *Physics Reports*, 430:277–374, 2006.
- [2] A. Ehlerding, S.T. Arnold, A.A. Viggiano, S. Kalhori, J. Semaniak, A.M. Derkatch, S. Rosen, M. af Ugglas, and M. Larsson. Rates and products of the dissociative recombination of C_3H_7^+ in low-energy electron collisions. *Journal of Physical Chemistry A*, 107(13):2179–2184, 2003. ISSN 1089-5639. URL http://pubs3.acs.org/acs/journals/doilookup?in_doi=10.1021/jp027729+.
- [3] Erik Vigren, Magdalena Kaminska, Mathias Hamberg, Vitali Zhaunerchyk, Richard D. Thomas, Jacek Semaniak, Mathias Danielsson, Mats Larsson, and Wolf D. Geppert. Dissociative recombination of the deuterated acetaldehyde ion, CD_3CDO^+ : product branching fractions, absolute cross sections and thermal rate coefficient. *Physical Chemistry Chemical Physics*, 9(22):2856–2861, 2007. URL <http://dx.doi.org/10.1039/b700649g>.
- [4] Erik Vigren, Magdalena Kaminska, Mathias Hamberg, Vitali Zhaunerchyk, Richard D. Thomas, Mathias Danielsson, Jacek Semaniak, Patrik U. Andersson, Mats Larsson, and Wolf D. Geppert. Dissociative recombination of fully deuterated protonated acetonitrile, CD_3CND^+ : product branching fractions, absolute cross section and thermal rate coefficient. *Physical Chemistry Chemical Physics*, 10(27):4014–4019, 2008. URL <http://dx.doi.org/10.1039/b801566j>.
- [5] E. Vigren, M. Hamberg, V. Zhaunerchyk, M. Kaminska, R. D. Thomas, M. Larsson, T. J. Millar, C. Walsh, , and W. D. Geppert. The dissociative recombination of protonated acrylonitrile, $\text{CH}_2\text{CHCNH}^+$, with implications for the nitrile chemistry in dark molecular clouds and the upper atmosphere of Titan. *The Astrophysical Journal*, 695(1):317–324, 2009. URL <http://stacks.iop.org/0004-637X/695/317>.

- [6] Viatcheslav Kokoouline and Chris H. Greene. Unified theoretical treatment of dissociative recombination of D_{3h} triatomic ions: Application to H_3^+ and D_3^+ . *Phys. Rev. A*, 68(1):012703–, July 2003. URL <http://link.aps.org/abstract/PRA/v68/e012703>.
- [7] Viatcheslav Kokoouline and Chris H. Greene. Theory of dissociative recombination of D_{3h} triatomic ions applied to H_3^+ . *Phys. Rev. Lett.*, 90(13):133201–, April 2003. URL <http://link.aps.org/abstract/PRL/v90/e133201>.
- [8] M. Larsson, B.J. McCall, and A.E. Orel. The dissociative recombination of H_3^+ - a saga coming to an end? *Chemical Physics Letters*, 462(4-6):145 – 151, 2008. ISSN 0009-2614. doi: DOI:10.1016/j.cplett.2008.06.069. URL <http://www.sciencedirect.com/science/article/B6TFN-4SVKST5-1/2/6abd65314f9f7b37b196684e1b84816c>.
- [9] M. T. Leu, Manfred A. Biondi, and R. Johnsen. Measurements of recombination of electrons with H_3^+ and H_5^+ ions. *Phys. Rev. A*, 8(1):413–419, July 1973. URL <http://link.aps.org/abstract/PRA/v8/p413>.
- [10] B. J. McCall, A. J. Huneycutt, R. J. Saykally, N. Djuric, G. H. Dunn, J. Semaniak, O. Novotny, A. Al-Khalili, A. Ehlerding, F. Hellberg, S. Kalhori, A. Neau, R. D. Thomas, A. Paal, F. Österdahl, and M. Larsson. Dissociative recombination of rotationally cold H_3^+ . *Phys. Rev. A*, 70(5):052716, Nov 2004. doi: 10.1103/PhysRevA.70.052716.
- [11] J. Glosík, R. Plašil, A. Pysanenko, O. Novotný, P. Hlavenka, P. Macko, and G. Bánó. Recombination studies in a He-Ar- H_2 plasma. *Journal of Physics: Conference Series*, 4:104–110, 2005. URL <http://stacks.iop.org/1742-6596/4/104>.
- [12] Sultana N. Nahar and Anil K. Pradhan. Electron-ion recombination rate coefficients, photoionization cross sections, and ionization fractions for astrophysically abundant elements. I. Carbon and Nitrogen. *The Astrophysical Journal Supplement Series*, 111:339–355, July 1997.
- [13] J. Woodall, M. Agúndez, A. J. Markwick-Kemper, and T. J. Millar. The UMIST database for astrochemistry 2006. *Astron. Astrophys.*, In Press.
- [14] C. H. Sheehan and J. P. St.-Maurice. Dissociative recombination of the methane family ions: rate coefficients and implications. *Advances in Space Research*, 33(2):216–220, 2004. URL <http://www.sciencedirect.com/science/article/B6V3S-4B84VVV-B/2/a792ccafaa1be2a9e0df7307025adddb>.
- [15] Sweden Stockholm University. Dissociative recombination database of Molecular Physics group, Department of Physics, Stockholm University. URL <http://mol.physto.se/DRdatabase/>.
- [16] A. Larson, A. Le Padellec, J. Semaniak, C. Stromholm, M. Larsson, S. Rosen, R. Peverall, H. Danared, N. Djuric, G.H. Dunn, and S. Datz. Branching fractions in dissociative recombination of CH_2^+ . *The Astrophysical Journal*, 505(1):459–465, 1998. URL <http://www.journals.uchicago.edu/doi/abs/10.1086/306164>.
- [17] L. Vejby-Christensen, L.H. Andersen, O. Heber, D. Kella, H.B. Pedersen, H.T. Schmidt, and D. Zajfman. Complete branching ratios for the dissociative recombination of H_2O^+ , H_3O^+ , and CH_3^+ . *The Astrophysical Journal*, 483(1):531–540, 1997. URL <http://www.journals.uchicago.edu/doi/abs/10.1086/304242>.
- [18] N. G. Adams, D. Smith, and Erich Alge. Measurements of dissociative recombination coefficients of H_3^+ , HCO^+ , N_2H^+ , and CH_5^+ at 95 and 300 K using the FALP apparatus. *J. Chem. Phys.*, 81(4):1778–1784, August 1984. URL <http://link.aip.org/link/?JCP/81/1778/1>.
- [19] L. Lehfaoui, C. Rebrion-Rowe, S. Laube, J. B. A. Mitchell, and B. R. Rowe. The dissociative recombination of hydrocarbon ions. I. Light alkanes. *J. Chem. Phys.*, 106(13):5406–5412, April 1997. URL <http://link.aip.org/link/?JCP/106/5406/1>.
- [20] Theodosia Gougousi, Michael F. Golde, and Rainer Johnsen. Electron-ion recombination rate coefficient measurements in a flowing afterglow plasma. *Chemical Physics Letters*, 265(3-5):399–403, February 1997. URL <http://www.sciencedirect.com/science/article/B6TFN-3S9DMHK-M/2/d3d0832b91a8f1ee0914e9ca1335db0f>.
- [21] J. Semaniak, Å. Larson, A. Le Padellec, C. Strömholm, M. Larsson, S. Rosén, R. Peverall, H. Danared, N. Djurić, G. H. Dunn, and S. Datz. Dissociative recombination and excitation of CH_5^+ : absolute cross sections and branching fractions. *The Astrophysical Journal*, 498:886–895, 1998.

- [22] J.L. McLain, V. Poterya, C.D. Molek, L.M. Babcock, and N.G. Adams. Flowing afterglow studies of the temperature dependencies for dissociative recombination of O_2^+ , CH_5^+ , C_2H_5^+ , and C_6H_7^+ with electrons. *J. Phys. Chem. A*, 108(32):6704–6708, 2004. URL http://pubs3.acs.org/acs/journals/doilookup?in_doi=10.1021/jp0402151.
- [23] Christopher D. Molek, Viktoriya Poterya, Nigel G. Adams, and Jason L. McLain. Development of a novel technique for quantitatively determining the products of electron-ion dissociative recombination. *International Journal of Mass Spectrometry*, 285(1-2):1 – 11, 2009. ISSN 1387-3806. doi: DOI:10.1016/j.ijms.2009.02.031. URL <http://www.sciencedirect.com/science/article/B6VND-4VYXMTY-1/2/e62674e26b30ce780b70bea5fe4ce917>.
- [24] Anneli Ehlerding, Fredrik Hellberg, Richard Thomas, Shirzad Kalhori, Albert A. Viggiano, Susan T. Arnold, Mats Larsson, and Magnus af Ugglas. Dissociative recombination of C_2H^+ and C_2H_4^+ : absolute cross sections and product branching ratios. *Phys. Chem. Chem. Phys.*, 6:949–954, 2004. URL <http://www.rsc.org/Publishing/Journals/CP/article.asp?doi=b314882c>.
- [25] A. M. Derkatch, A. Al-Khalili, L. Vikor, A. Neau, W. Shi, H. Danared, M. af Ugglas, and M. Larsson. Branching ratios in dissociative recombination of the C_2H_2^+ molecular ion. *Journal of Physics B: Atomic, Molecular and Optical Physics*, 32(14):3391, 1999. URL <http://stacks.iop.org/0953-4075/32/3391>.
- [26] S. Kalhori, A.A. Viggiano, S.T. Arnold, S. Rosén, J. Semaniak, A.M. Derkatch, M. af Ugglas, and M. Larsson. Dissociative recombination of C_2H_3^+ . *Astron. Astrophys.*, 391:1159–1165, 2002.
- [27] Nigel G. Adams, Viktoriya Poterya, and Lucia M. Babcock. Electron molecular ion recombination: Product excitation and fragmentation. *Mass Spectrometry Reviews*, 25(5):798–828, 2006. URL <http://dx.doi.org/10.1002/mas.20084>.
- [28] W. Geppert, A. Ehlerding, F. Hellberg, S. Kalhori, R. D. Thomas, O. Novotny, S. T. Arnold, T. M. Miller, A. A. Viggiano, and M. Larsson. First observation of four-body breakup in electron recombination: C_2D_5^+ . *Phys. Rev. Lett.*, 93(15):153201–, October 2004. URL <http://link.aps.org/abstract/PRL/v93/e153201>.
- [29] O. Heber, K. Seiersen, H. Bluhme, A. Svendsen, L. H. Andersen, and L. Maunoury. Dissociative recombination of small carbon cluster cations. *Physical Review A (Atomic, Molecular, and Optical Physics)*, 73(2):022712, 2006. doi: 10.1103/PhysRevA.73.022712. URL <http://link.aps.org/abstract/PRA/v73/e022712>.
- [30] G. Angelova, O. Novotny, J. B. A. Mitchell, C. Rebrion-Rowe, J. L. Le Garrec, H. Bluhme, A. Svendsen, and L. H. Andersen. Branching ratios for the dissociative recombination of hydrocarbon ions: III. The cases of C_3H_n^+ ($n=1-8$). *International Journal of Mass Spectrometry*, 235(1):7 – 13, 2004. ISSN 1387-3806. doi: DOI:10.1016/j.ijms.2004.03.002. URL <http://www.sciencedirect.com/science/article/B6VND-4CBVVTN-2/2/5688e1c7b26b2265da8f0bf5021f46f4>.
- [31] J B A Mitchell, G Angelova, C Rebrion-Rowe, O Novotny, J L Le Garrec, H Bluhme, K Seiersen, A Svendsen, and L H Andersen. Branching ratios for the dissociative recombination of hydrocarbon ions. *Journal of Physics: Conference Series*, 4:198–204, 2005. URL <http://stacks.iop.org/1742-6596/4/198>.
- [32] Jason L. McLain, Viktoriya Poterya, Christopher D. Molek, Douglas M. Jackson, Lucia M. Babcock, and Nigel G. Adams. C_3H_3^+ isomers: Temperature dependencies of production in the H_3^+ reaction with allene and loss by dissociative recombination with electrons. *The Journal of Physical Chemistry A*, 109(23): 5119–5123, 2005. doi: 10.1021/jp0444159. URL <http://pubs.acs.org/doi/abs/10.1021/jp0444159>.
- [33] W. D. Geppert, R. Thomas, A. Ehlerding, F. Hellberg, F. Österdahl, M. af Ugglas, and M. Larsson. Dissociative recombination of C_3H_4^+ : preferential formation of the C_3H_3 radical. *International Journal of Mass Spectrometry*, 237(1):25–32, September 2004. URL <http://www.sciencedirect.com/science/article/B6VND-4D09MWJ-1/2/805064fbc0ebcb0cf3a578fc6be7f14>.
- [34] M. Larsson, A. Ehlerding, W. D. Geppert, F. Hellberg, S. Kalhori, R. D. Thomas, N. Djuric, F. Österdahl, G. Angelova, J. Semaniak, O. Novotny, S. T. Arnold, and A. A. Viggiano. Rate constants and branching ratios for the dissociative recombination of C_3D_7^+ and C_4D_9^+ . *The Journal of Chemical Physics*, 122(15):156101, 2005. doi: 10.1063/1.1872839. URL <http://link.aip.org/link/?JCP/122/156101/1>.

- [35] G. Angelova, O. Novotny, J. B. A. Mitchell, C. Rebrion-Rowe, J. L. Le Garrec, H. Bluhme, K. Seiersen, and L. H. Andersen. Branching ratios for the dissociative recombination of hydrocarbon ions: II. The cases of $C_4H_n^+$ ($n=1-9$). *International Journal of Mass Spectrometry*, 232(2):195 – 203, 2004. ISSN 1387-3806. doi: DOI:10.1016/j.ijms.2003.12.013. URL <http://www.sciencedirect.com/science/article/B6VND-4BRBCW9-1/2/40fb39b7c380a971dca0f5c4a67c2bbb>.
- [36] Nigel G. Adams and David Smith. Measurements of the dissociative recombination coefficients for several polyatomic ion species at 300 K. *Chemical Physics Letters*, 144(1):11 – 14, 1988. ISSN 0009-2614. doi: DOI:10.1016/0009-2614(88)87081-7. URL <http://www.sciencedirect.com/science/article/B6TFN-451FS1G-3/2/5443bdec7aa8a296444a66291317f492>.
- [37] C. Rebrion-Rowe, L. Lehfaoui, B. R. Rowe, and J. B. A. Mitchell. The dissociative recombination of hydrocarbon ions. II. Alkene and alkyne derived species. *J. Chem. Phys.*, 108(17):7185–7189, May 1998. URL <http://link.aip.org/link/?JCP/108/7185/1>.
- [38] J. B. A. Mitchell, C. Rebrion-Rowe, J. L. Le Garrec, G. Angelova, H. Bluhme, K. Seiersen, and L. H. Andersen. Branching ratios for the dissociative recombination of hydrocarbon ions. I: The cases of $C_4H_9^+$ and $C_4H_5^+$. *International Journal of Mass Spectrometry*, 227(2):273–279, June 2003. URL <http://www.sciencedirect.com/science/article/B6VND-48GPCSR-1/2/7b0520235e727216f28a2c66e3907248>.
- [39] H. Abouelaziz, J. C. Gomet, D. Pasquero, B. R. Rowe, and J. B. A. Mitchell. Measurements of $C_3H_3^+$, $C_5H_3^+$, $C_6H_6^+$, $C_7H_5^+$, and $C_{10}H_8^+$ dissociative recombination rate coefficients. *J. Chem. Phys.*, 99(1):237–243, July 1993. URL <http://link.aip.org/link/?JCP/99/237/1>.
- [40] L. Viktor, A. Al-Khalili, H. Danared, N. Djurić, G.-H. Dunn, M. Larsson, A. Le Padellec, S. Rosén, and M. Af Ugglas. Branching fractions in the dissociative recombination of NH_4^+ and NH_2^+ molecular ions. *Astron. Astrophys.*, 344:1027–1033, April 1999.
- [41] R. D. Thomas, F. Hellberg, A. Neau, S. Rosén, M. Larsson, C. R. Vane, M. E. Bannister, S. Datz, A. Petrigiani, and W. J. van der Zande. Three-body fragmentation dynamics of amidogen and methylene radicals via dissociative recombination. *Phys. Rev. A*, 71(3):032711–16, March 2005. URL <http://link.aps.org/abstract/PRA/v71/e032711>.
- [42] Chou-Mou Huang, Manfred A. Biondi, and Rainer Johnsen. Recombination of electrons with $NH_4^+ \cdot (NH_3)_n$ -series ions. *Phys. Rev. A*, 14(3):984–, September 1976. URL <http://link.aps.org/abstract/PRA/v14/p984>.
- [43] R. D. DuBois, J. B. Jeffries, and G. H. Dunn. Dissociative recombination cross sections for NH_4^+ ions and electrons. *Phys. Rev. A*, 17(4):1314–, April 1978. URL <http://link.aps.org/abstract/PRA/v17/p1314>.
- [44] J. Öjekull, P. U. Andersson, M. B. Någård, J. B. C. Pettersson, A. M. Derkach, A. Neau, S. Rosén, R. Thomas, M. Larsson, F. Österdahl, J. Semaniak, H. Danared, A. Källberg, M. af Ugglas, and N. Marković. Dissociative recombination of NH_4^+ and ND_4^+ ions: Storage ring experiments and ab initio molecular dynamics. *J. Chem. Phys.*, 120(16):7391–7399, April 2004. URL <http://link.aip.org/link/?JCP/120/7391/1>.
- [45] N. G. Adams, C. R. Herd, M. Geoghegan, D. Smith, A. Canosa, J. C. Gomet, B. R. Rowe, J. L. Queffelec, and M. Morlais. Laser induced fluorescence and vacuum ultraviolet spectroscopic studies of H-atom production in the dissociative recombination of some protonated ions. *The Journal of Chemical Physics*, 94(7):4852–4857, 1991. doi: 10.1063/1.460570. URL <http://link.aip.org/link/?JCP/94/4852/1>.
- [46] W. H. Kasner. Study of the temperature dependence of electron-ion recombination in Nitrogen. *Phys. Rev.*, 164(1):194–200, Dec 1967. doi: 10.1103/PhysRev.164.194.
- [47] F. J. Mehr and Manfred A. Biondi. Electron temperature dependence of recombination of O_2^+ and N_2^+ ions with electrons. *Phys. Rev.*, 181(1):264–271, May 1969. doi: 10.1103/PhysRev.181.264.
- [48] M R Mahdavi, J B Hasted, and M M Nakshbandi. Electron-ion recombination measurements in the flowing afterglow. *Journal of Physics B: Atomic and Molecular Physics*, 4(12):1726–1737, 1971. URL <http://stacks.iop.org/0022-3700/4/1726>.
- [49] A J Cunningham and R M Hobson. Dissociative recombination at elevated temperatures. IV. N_2^+ dominated afterglows. *Journal of Physics B: Atomic and Molecular Physics*, 5(12):2328–2331, 1972. URL <http://stacks.iop.org/0022-3700/5/2328>.

- [50] Edward C. Zipf. The dissociative recombination of vibrationally excited N ions. *Geophys. Res. Lett.*, 7: 645–648, 1980. doi: 10.1029/GL007i009p00645.
- [51] M Geoghegan, N G Adams, and D Smith. Determination of the electron-ion dissociative recombination coefficients for several molecular ions at 300 K. *Journal of Physics B: Atomic, Molecular and Optical Physics*, 24(10):2589–2599, 1991. URL <http://stacks.iop.org/0953-4075/24/2589>.
- [52] A. Canosa, J. C. Gomet, B. R. Rowe, and J. L. Queffelec. Flowing afterglow langmuir probe measurement of the $N_2^+(v=0)$ dissociative recombination rate coefficient. *The Journal of Chemical Physics*, 94(11): 7159–7163, 1991. doi: 10.1063/1.460199. URL <http://link.aip.org/link/?JCP/94/7159/1>.
- [53] J. R. Peterson, A. Le Padellec, H. Danared, G. H. Dunn, M. Larsson, A. Larson, R. Peverall, C. Strömholm, S. Rosén, M. af Ugglas, and W. J. van der Zande. Dissociative recombination and excitation of N_2^+ : Cross sections and product branching ratios. *The Journal of Chemical Physics*, 108 (5):1978–1988, 1998. doi: 10.1063/1.475577. URL <http://link.aip.org/link/?JCP/108/1978/1>.
- [54] D. Kella, P. J. Johnson, H. B. Pedersen, L. Vejby-Christensen, and L. H. Andersen. Branching ratios for dissociative recombination of $^{15}N^{14}N^+$. *Phys. Rev. Lett.*, 77(12):2432–2435, Sep 1996. doi: 10.1103/PhysRevLett.77.2432.
- [55] T. Amano. The dissociative recombination rate coefficients of H_3^+ , HN_2^+ , and HCO^+ . *J. Chem. Phys.*, 92:6492–6501, 1990.
- [56] Richard E. Rosati, Rainer Johnsen, and Michael F. Golde. Yield of electronically excited N_2 molecules from the dissociative recombination of N_2H^+ with e^- . *J. Chem. Phys.*, 120(17):8025–8030, May 2004. URL <http://link.aip.org/link/?JCP/120/8025/1>.
- [57] W. D. Geppert, R. Thomas, J. Semaniak, A. Ehlerding, T. J. Millar, F. Österdahl, M. af Ugglas, N. Djurić, A. Paál, and M. Larsson. Dissociative recombination of N_2H^+ : Evidence for fracture of the N-N bond. *The Astrophysical Journal*, 609(1):459–464, 2004. URL <http://www.journals.uchicago.edu/doi/abs/10.1086/420733>.
- [58] V. Poterya, J.L. McLain, N.G. Adams, and L.M. Babcock. Mechanisms of electron-ion recombination of N_2H^+/N_2D^+ and HCO^+/DCO^+ ions: Temperature dependence and isotopic effect. *J. Phys. Chem. A*, 109(32):7181–7186, 2005. URL http://pubs3.acs.org/acs/journals/doilookup?in_doi=10.1021/jp051945b.
- [59] W. D. Geppert, R. Thomas, A. Ehlerding, F. Hellberg, F. Österdahl, M. Hamberg, J. Semaniak, V. Zhaunerchyk, M. Kaminska, A. Källberg, A. Paál, and M. Larsson. Dissociative recombination branching ratios and their influence on interstellar clouds. *J. Phys.: Conf. Ser.*, 4:26–31, 2005.
- [60] Nigel G. Adams and Lucia M. Babcock. Molecular ion recombination in trapped and flowing plasmas: methods, recent results, new goals, open questions. *Journal of Physics: Conference Series*, page 38, 2005. URL <http://stacks.iop.org/1742-6596/4/38>.
- [61] C. D. Molek, J. L. McLain, V. Poterya, and N. G. Adams. A remeasurement of the products for electron recombination of N_2H^+ using a new technique: No significant $NH + N$ production. *The Journal of Physical Chemistry A*, 111(29):6760–6765, July 2007. ISSN 1089-5639. URL <http://dx.doi.org/10.1021/jp068965d>.
- [62] A. Le Padellec, J. B. A. Mitchell, A. Al-Khalili, H. Danared, A. Källberg, Å. Larson, S. Rosén, M. af Ugglas, L. Viktor, and M. Larsson. Storage ring measurements of the dissociative recombination and excitation of the cyanogen ion $CN^+(X^1\Sigma^+ \text{ and } a^3\Pi, \nu = 0)$. *The Journal of Chemical Physics*, 110(2): 890–901, 1999. doi: 10.1063/1.478056. URL <http://link.aip.org/link/?JCP/110/890/1>.
- [63] D Talbi, A Le Padellec, and J B A Mitchell. Quantum chemical calculations for the dissociative recombination of HCN^+ and HNC^+ . *Journal of Physics B: Atomic, Molecular and Optical Physics*, (18):3631, 2000. URL <http://stacks.iop.org/0953-4075/33/3631>.
- [64] C Sheehan, C Le Padellec, C Lennard, D Talbi, and J B A Mitchell. Merged beam measurement of the dissociative recombination of HCN^+ and HNC^+ . *Journal of Physics B: Atomic, Molecular and Optical Physics*, 32(14):3347–3360, 1999. URL <http://stacks.iop.org/0953-4075/32/3347>.
- [65] J. Semaniak, B. F. Minaev, A. M. Derkatch, F. Hellberg, A. Neau, S. Rosén, R. Thomas, M. Larsson, H. Danared, A. Paál, and M. af Ugglas. Dissociative recombination of $HCNH^+$: Absolute cross-sections and branching ratios. *The Astrophysical Journal Supplement Series*, 135(2):275–283, 2001. URL <http://www.journals.uchicago.edu/doi/abs/10.1086/321797>.

- [66] J.L. McLain, C.D. Molek, D. Osborne Jr., and N.G. Adams. Flowing afterglow studies of the electron recombination of protonated cyanides $(RCN)H^+$ and their proton-bound dimer ions $(RCN)_2H^+$ where R is H, CH_3 , and CH_3CH_2 . *International Journal of Mass Spectrometry*, 282(3):85 – 90, 2009. ISSN 1387-3806. doi: DOI:10.1016/j.ijms.2009.02.004. URL <http://www.sciencedirect.com/science/article/B6VND-4VM445F-3/2/ab43e578ccdf6a67a7feeb258a66fb>.
- [67] J.L. McLain and N.G. Adams. Flowing afterglow studies of temperature dependencies for electron dissociative recombination of $HCNH^+$, CH_3CNH^+ and $CH_3CH_2CNH^+$ and their symmetrical proton-bound dimers. *Planetary and Space Science*, 57(13):1642 – 1647, 2009. ISSN 0032-0633. doi: DOI:10.1016/j.pss.2009.09.005. URL <http://www.sciencedirect.com/science/article/B6V6T-4X6MSXW-2/2/09c4d6fe7753bdaae0c1a654ada5d6ee>. Surfaces and Atmospheres of the Outer Planets, Their Satellites and Ring Systems: Part V.
- [68] A. P. Hickman, R. D. Miles, C. Hayden, and D. Talbi. Dissociative recombination of $e + HCNH^+$: Diabatic potential curves and dynamics calculations. *A&A*, 438(1):31–37, jul 2005. doi: 10.1051/0004-6361:20052658. URL <http://dx.doi.org/10.1051/0004-6361:20052658>.
- [69] Keisaku Ishii, Asami Tajima, Tetsuya Taketsugu, and Koichi Yamashita. Theoretical elucidation of the unusually high $[HNC]/[HCN]$ abundance ratio in interstellar space: Two-dimensional and two-state quantum wave packet dynamics study on the branching ratio of the dissociative recombination reaction $HCNH^+ + e^- \rightarrow HNC/HCN + H$. *The Astrophysical Journal*, 636(2):927–931, 2006. URL <http://www.journals.uchicago.edu/doi/abs/10.1086/498343>.
- [70] Marilyn E. Jacox and Warren E. Thompson. Infrared spectroscopy and photochemistry of $NCCN^+$ and $CNCN^+$ trapped in solid neon. *The Journal of Chemical Physics*, 126(5):054308, 2007. doi: 10.1063/1.2433945. URL <http://link.aip.org/link/?JCP/126/054308/1>.
- [71] W. D. Geppert, A. Ehlerding, F. Hellberg, J. Semaniak, F. Osterdahl, M. Kaminska, A. Al-Khalili, V. Zhaunerchyk, R. Thomas, M. af Ugglas, A. Kallberg, A. Simonsson, , and M. Larsson. Dissociative recombination of nitrile ions: $DCCCN^+$ and $DCCCNH^+$. *The Astrophysical Journal*, 613(2):1302–1309, 2004. URL <http://stacks.iop.org/0004-637X/613/1302>.
- [72] Yoshihiro Osamura, Kaori Fukuzawa, R. Terzieva, , and Eric Herbst. A molecular orbital study of the $HC_3NH^+ + e^-$ dissociative recombination and its role in the production of cyanoacetylene isomers in interstellar clouds. *The Astrophysical Journal*, 519(2):697–704, 1999. URL <http://stacks.iop.org/0004-637X/519/697>.
- [73] M. J. Jensen, R. C. Bilodeau, O. Heber, H. B. Pedersen, C. P. Safvan, X. Urbain, D. Zajfman, and L. H. Andersen. Dissociative recombination and excitation of H_2O^+ and HDO^+ . *Phys. Rev. A*, 60(4):2970–2976, October 1999. URL <http://link.aps.org/abstract/PRA/v60/p2970>.
- [74] S. Rosén, A. Derkach, J. Semaniak, A. Neau, A. Al-Khalili, A. Le Padellec, L. Viktor, R. Thomas, H. Danared, M. af Ugglas, and M. Larsson. Recombination of simple molecular ions studied in storage ring: dissociative recombination of H_2O^+ . *Faraday Discussions*, 115:295–302, 2000. URL <http://dx.doi.org/10.1039/a909314a>.
- [75] B. R. Rowe, F. Vallee, J. L. Queffelec, J. C. Gomet, and M. Morlais. The yield of oxygen and hydrogen atoms through dissociative recombination of H_2O^+ ions with electrons. *The Journal of Chemical Physics*, 88(2):845–850, 1988. doi: 10.1063/1.454164. URL <http://link.aip.org/link/?JCP/88/845/1>.
- [76] R. A. Heppner, F. L. Walls, W. T. Armstrong, and G. H. Dunn. Cross-section measurements for electron- H_3O^+ recombination. *Phys. Rev. A*, 13(3):1000–1011, Mar 1976. doi: 10.1103/PhysRevA.13.1000.
- [77] M.J. Jensen, R.C. Bilodeau, C.P. Safvan, K. Seiersen, L.H. Andersen, H.B. Pedersen, and O. Heber. Dissociative recombination of H_3O^+ , HD_2O^+ , and D_3O^+ . *The Astrophysical Journal*, 543(2):764–774, 2000. URL <http://www.journals.uchicago.edu/doi/abs/10.1086/317137>.
- [78] A. Neau, A. Al Khalili, S. Rosén, A. Le Padellec, A. M. Derkach, W. Shi, L. Viktor, M. Larsson, J. Semaniak, R. Thomas, M. B. Någård, K. Andersson, H. Danared, and M. af Ugglas. Dissociative recombination of D_3O^+ and H_3O^+ : Absolute cross sections and branching ratios. *The Journal of Chemical Physics*, 113(5):1762–1770, 2000. doi: 10.1063/1.481979. URL <http://link.aip.org/link/?JCP/113/1762/1>.
- [79] L. H. Andersen, O. Heber, D. Kella, H. B. Pedersen, L. Vejby-Christensen, and D. Zajfman. Production of water molecules from dissociative recombination of H_3O^+ with electrons. *Phys. Rev. Lett.*, 77(24):4891–, December 1996. URL <http://link.aps.org/abstract/PRL/v77/p4891>.

- [80] Megumi Kayanuma, Tetsuya Taketsugu, and Keisaku Ishii. Ab initio surface hopping simulation on dissociative recombination of H_3O^+ . *Chemical Physics Letters*, 418(4-6):511–518, February 2006. URL <http://www.sciencedirect.com/science/article/B6TFN-4HP6GCY-4/2/4b472ede977ab2c81ee491be14cd3dd6>.
- [81] S. Rosén, R. Peverall, M. Larsson, A. Le Padellec, J. Semaniak, Å. Larson, C. Strömholm, W. J. van der Zande, H. Danared, and G. H. Dunn. Absolute cross sections and final-state distributions for dissociative recombination and excitation of $\text{CO}^+(\text{v}=0)$ using an ion storage ring. *Phys. Rev. A*, 57(6):4462–, June 1998. URL <http://link.aps.org/abstract/PRA/v57/p4462>.
- [82] S. Laubé, L. Lehfaoui, B. R. Rowe, and J. B. A. Mitchell. The dissociative recombination of CO^+ . *Journal of Physics B: Atomic, Molecular and Optical Physics*, 31(18):4181, 1998. URL <http://stacks.iop.org/0953-4075/31/4181>.
- [83] M. T. Leu, Manfred A. Biondi, and R. Johnsen. Measurements of recombination of electrons with HCO^+ ions. *Phys. Rev. A*, 8(1):420–422, Jul 1973. doi: 10.1103/PhysRevA.8.420.
- [84] B. Ganguli, M. A. Biondi, R. Johnsen, and J. L. Dulaney. Electron-temperature dependence of the recombination of HCO^+ ions with electrons. *Phys. Rev. A*, 37(7):2543–2547, Apr 1988. doi: 10.1103/PhysRevA.37.2543.
- [85] B. R. Rowe, J. C. Gomet, A. Canosa, C. Rebrion, and J. B. A. Mitchell. A further study of HCO^+ dissociative recombination. *The Journal of Chemical Physics*, 96(2):1105–1110, 1992. doi: 10.1063/1.462196. URL <http://link.aip.org/link/?JCP/96/1105/1>.
- [86] M. Hamberg, W. D. Geppert, R. D. Thomas, V. Zhaunerchyk, F. Österdahl, A. Ehlerding, M. Kamin-ska, J. Semaniak, M. af Ugglas, A. Källberg, A. Paal, A. Simonsson, and M. Larsson. Experimental determination of dissociative recombination reaction pathways and absolute reaction cross-sections of CH_2OH^+ , CD_2OD^+ and CD_2OD_2^+ . *Molecular Physics: An International Journal at the Interface Between Chemistry and Physics*, 105(5):899–906, 2007. ISSN 0026-8976. URL <http://www.informaworld.com/10.1080/00268970701206642>.
- [87] Charles S. Weller and Manfred A. Biondi. Recombination, attachment, and ambipolar diffusion of electrons in photo-ionized NO afterglows. *Phys. Rev.*, 172(1):198–206, Aug 1968. doi: 10.1103/PhysRev.172.198.
- [88] T Mostefaoui, S Laube, G Gautier, C Rebrion-Rowe, B R Rowe, and J B A Mitchell. The dissociative recombination of NO^+ : the influence of the vibrational excitation state. *Journal of Physics B: Atomic, Molecular and Optical Physics*, 32(22):5247–5256, 1999. URL <http://stacks.iop.org/0953-4075/32/5247>.
- [89] Clint H. Sheehan and J.-P. St.-Maurice. Dissociative recombination of N_2^+ , O_2^+ , and NO^+ : rate coefficients for ground state and vibrationally excited ions. *J. Geophys. Res.*, 109:–, March 2004. URL <http://dx.doi.org/10.1029/2003JA010132>.
- [90] Dieter Kley, George M. Lawrence, and Edward J. Stone. The yield of $\text{N}(^2D)$ atoms in the dissociative recombination of NO^+ . *The Journal of Chemical Physics*, 66(9):4157–4165, 1977. doi: 10.1063/1.434491. URL <http://link.aip.org/link/?JCP/66/4157/1>.
- [91] P M Mul, J W McGowan, P Defrance, and J B A Mitchell. Merged electron-ion beam experiments. V. Dissociative recombination of OH^+ , H_2O^+ , H_3O^+ and D_3O^+ . *Journal of Physics B: Atomic and Molecular Physics*, 16(16):3099–3107, 1983. URL <http://stacks.iop.org/0022-3700/16/3099>.
- [92] P. M. Mul and J. W. McGowan. Dissociative recombination of C_2^+ , C_2H^+ , C_2H_2^+ and C_2H_3^+ . *The Astrophysical Journal*, 237:749–751, may 1980. doi: 10.1086/157921.
- [93] Oldřich Novotný, Henrik Buhr, Julia Stützel, Mario B. Mendes, Max H. Berg, Dennis Bing, Michael Froese, Manfred Grieser, Oded Heber, Brandon Jordon-Thaden, Claude Krantz, Michael Lange, Michael Lestinsky, Steffen Novotny, Sebastian Menk, Dmitry A. Orlov, Annemieke Petrigani, Michael L. Rappaport, Andrey Shornikov, Dirk Schwalm, Daniel Zajfman, and Andreas Wolf. Fragmentation channels in dissociative electron recombination with hydronium and other astrophysically important species. *The Journal of Physical Chemistry A*, pages –, January 2010. ISSN 1089-5639. URL <http://dx.doi.org/10.1021/jp9104097>.
- [94] T. L. Williams, N. G. Adams, L. M. Babcock, C. R. Herd, and M. Geoghegan. Production and loss of the water-related species H_3O^+ , H_2O and OH in dense interstellar clouds. *Monthly Notices of the Royal Astronomical Society*, 282:413–420, September 1996.

- [95] V. Vuitton, R. V. Yelle, and V. G. Anicich. The nitrogen chemistry of Titan’s upper atmosphere revealed. *The Astrophysical Journal*, 647:L175–L178, 2006.