

# Integration of HEPData in INSPIRE

# Where are we ?

- HEPData records are harvested and uploaded into a demo site containing complete INSPIRE
- Data is displayed together with records (ugly)

# Where do we want to be ?

- HEPData record should be harvested and INSPIRE records updated once a day in the production environment
- Indexes should be established to enable search
- The data should be displayed in a pretty way
- There should be format for displaying standalone HEPData records

An additional HEPData tab is displayed for records having data attached



[Information](#) [References \(29\)](#) [Citations \(0\)](#) [Files](#) [Plots](#) [HEP Data](#)

/ Experiment-HEP CERN-PH-EP-2011-077

## Search for Heavy Long-Lived Charged Particles with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV

## Search for Heavy Long-Lived Charged Particles with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV

[Aad, Georges](#) (Freiburg U.) ; [Abbott, Brad](#) (Oklahoma U.) ; [Abdallah, Jalal](#) (Barcelona, IFAE) ; [Abdelalim, Ahmed Ali](#) (Geneva U.) ; [Abdesselam, Abdelouahab](#) (Oxford U.) ; [Abdinov, Ovsat](#) (Baku, Inst. Phys.) ; [Abi, Babak](#) (Oklahoma State U.) ; [Abolins, Maris](#) (Michigan State U.) ; [Abramowicz, Halina](#) (Tel Aviv U.) ; [Abreu, Henso](#) (Orsay, LAL) ; [Acerbi, Emilio](#) (INFN, Milan) ; [Acharya, Bobby Samir](#) (INFN, Udine) ; [Adams, David](#) (Brookhaven) ; [Addy, Tetteh](#) (Hampton U.) ; [Adelman, Jahred](#) (Yale U.) ; [Aderholz, Michael](#) (Munich, Max Planck Inst.) ; [Adomeit, Stefanie](#) (Munich U.) ; [Adragna, Paolo](#) (Queen Mary, U. of London) ; [Adye, Tim](#) (Rutherford) ; [Aefsky, Scott](#) (Brandeis U.) ; [Aguilar-Saavedra, Juan Antonio](#) (CAFPE, Granada) ; [Aharrouche, Mohamed](#) (Mainz U.) ; [Ahlen, Steven](#) (Boston U.) ; [Ahles, Florian](#) (Freiburg U.) ; [Ahmad, Ashfaq](#) (SUNY, Stony Brook) [...] [Show all 3023 authors](#)

**Keyword(s):** [SUSY](#) ; [ATLAS](#) ; [Long-lived particles](#)  
**Note:** \* Temporary entry \*

Record created 2011-06-23, last modified 2011-12-13 [Similar records](#)

# HEPData entries as separate records

```
001100000 001__ 1100000
001100000 245__ $$9HEPDATA$$a<br>Charged particle MEAN(PTOUT**2) distribution<br>Location: F 9,T 12
001100000 336__ $$tDATASET
001100000 520__ $$9HEPDATA
001100000 6531_ $$c0$$k$$v 
001100000 6531_ $$c1$$kSqrt(S)$$v52 TO 57 GeV
001100000 6531_ $$c0$$k$$v 
001100000 6531_ $$c1$$kSqrt(S)$$v52.0-57.0 GeV
001100000 710__ $$gAMY Collaboration
001100000 786__ $$q0$$rKEK-PREPRINT-89-149$$w283337
001100000 8564_ $$uhttp://inspirehepdev2.cern.ch/record/1100000/files/Data$$ydata extracted from the table
001100000 910__ $$dMEAN(PTOUT**2) IN GEV**2$$n0$$t 
001100000 910__ $$d(1/N) * D(MEAN(NAME=PTOUT**2)) IN GEV**2$$n1$$tRE : E+ E- --> CHARGED X
001100000 980__ $$aDATA
```

# The harvesting process

- List of all HEPData records is exposed
- Harvesting script reads the page once every day and checks which records have changed

# The backend

# The interface

<http://inspirehepdev2.cern.ch/record/902309/hepdata>

The image shows a screenshot of the HEPData interface with several red annotations and arrows pointing to specific elements:

- Data headers**: Points to the first column header, **ABS(YRAP) : 2.0-2.5**.
- The comment extracted from HEPData**: Points to the **Comments:** section, which contains the text: "Inclusive jet double-differential cross section as a function of jet transverse momentum in the [rapidity] range 2.0 to 2.5 using an anti-kT jet resolution parameter R of 0.5. Location: F 1".
- Expand the graphics**: Points to the vertical text **↑↑↑Expand plot↑↑↑** on the right side of the table.
- Expand the data**: Points to the horizontal text **↓↓↓Expand↓↓↓** at the bottom left of the table.
- Column headers**: Points to the main header text **PT IN GEVD2(SIG)/DPT/DYRAP IN PB/GEVNLO THEORY CORRECTION FACTOR**.

The interface content includes:

**Comments:**  
Inclusive jet double-differential cross section as a function of jet transverse momentum in the [rapidity] range 2.0 to 2.5 using an anti-kT jet resolution parameter R of 0.5.  
Location: F 1

**Table**

ABS(YRAP) : 2.0-2.5
$R : 0.5$
$PP \rightarrow JET X$
$\sqrt{S} : 7000.0 \text{ GeV}$
PT IN GEVD2(SIG)/DPT/DYRAP IN PB/GEVNLO THEORY CORRECTION FACTOR



**Comments:**

Inclusive jet double-differential cross section as a function of jet transverse momentum in the |rapidity| range 2.0 to 2.5 using an anti-kT jet resolution parameter R of 0.5.  
Location: F 1

**Table**

ABS(YRAP) : 2.0-2.5		
:	$R : 0.5$	
:	$PP \rightarrow JET X$	
:	$\sqrt{S} : 7000.0 \text{ GeV}$	
PT IN GEV	D2(SIG)/DPT/DYRAP IN PB/GEV	NLO THEORY CORRECTION FACTOR
†††Collapse†††		
19.4 (bin: 18. – 21.)	10600000 ± 0.7% (stat) +20.6%,-17.0% (sys)	1.28 ± 27.9%
22.4 (bin: 21. – 24.)	5550000 ± 1.0% (stat) +18.3%,-15.4% (sys)	1.23 ± 25.4%
25.9 (bin: 24. – 28.)	2770000 ± 1.3% (stat) +16.6%,-14.2% (sys)	1.19 ± 22.1%
29.9 (bin: 28. – 32.)	1410000 ± 1.8% (stat) +15.3%,-13.2% (sys)	1.15 ± 18.8%
34.3 (bin: 32. – 37.)	748000 ± 2.3% (stat) +14.3%,-12.5% (sys)	1.11 ± 15.8%
39.8 (bin: 37. – 43.)	351000 ± 1.3% (stat) +13.5%,-11.9% (sys)	1.09 ± 12.9%
45.8 (bin: 43. – 49.)	169000 ± 1.9% (stat) +13.0%,-11.6% (sys)	1.07 ± 10.6%
52.3 (bin: 49. – 56.)	90100 ± 2.4% (stat) +12.7%,-11.4% (sys)	1.06 ± 8.8%
59.7 (bin: 56. – 64.)	44000 ± 1.0% (stat) +12.6%,-11.3% (sys)	1.05 ± 7.2%
68.6 (bin: 64. – 74.)	21500 ± 1.3% (stat) +12.6%,-11.4% (sys)	1.04 ± 5.9%
78.7 (bin: 74. – 84.)	10300 ± 1.9% (stat) +12.9%,-11.6% (sys)	1.04 ± 4.8%
90.0 (bin: 84. – 97.)	5020 ± 0.6% (stat) +13.2%,-11.8% (sys)	1.03 ± 3.8%
104.8 (bin: 97. – 114.)	2190 ± 0.8% (stat) +13.6%,-12.2% (sys)	1.03 ± 3.0%
122.7 (bin: 114. – 133.)	896 ± 0.4% (stat) +14.3%,-12.8% (sys)	1.03 ± 2.3%
142.2 (bin: 133. – 153.)	384 ± 0.6% (stat) +15.2%,-13.5% (sys)	1.03 ± 1.7%
162.7 (bin: 153. – 174.)	171 ± 0.6% (stat) +16.2%,-14.2% (sys)	1.03 ± 1.3%
184.2 (bin: 174. – 196.)	78.7 ± 0.8% (stat) +17.3%,-15.1% (sys)	1.03 ± 1.0%
207.1 (bin: 196. – 220.)	36 ± 0.8% (stat) +18.7%,-16.1% (sys)	1.03 ± 0.7%
231.6 (bin: 220. – 245.)	16.5 ± 1.1% (stat) +20.5%,-17.4% (sys)	1.03 ± 0.4%
257.5 (bin: 245. – 272.)	7.26 ± 1.2% (stat) +22.5%,-18.9% (sys)	1.03 ± 0.2%
285.0 (bin: 272. – 300.)	3.29 ± 1.7% (stat) +25.0%,-20.6% (sys)	1.03 ± 0.00%
313.9 (bin: 300. – 330.)	1.4 ± 2.5% (stat) +27.9%,-22.5% (sys)	1.03 ± 0.2%
344.8 (bin: 330. – 362.)	0.582 ± 3.7% (stat) +31.3%,-24.7% (sys)	1.03 ± 0.4%
377.3 (bin: 362. – 395.)	0.221 ± 5.9% (stat) +35.8%,-27.4% (sys)	1.03 ± 0.7%
411.2 (bin: 395. – 430.)	0.078 ± 9.4% (stat) +40.5%,-30.0% (sys)	1.02 ± 1.0%
508.1 (bin: 430. – 737.)	0.00555 ± 11.4% (stat) +52.7%,-36.1% (sys)	1.02 ± 1.5%

††Expand plot††

Plot summarising all the data

**Comments:**


Inclusive jet double-differential cross section as a function of jet transverse momentum in the  $|\text{rapidity}|$  range 2.0 to 2.5 using an anti-kT jet resolution parameter  $R$  of 0.5.  
Location: F 1

**Table**


ABS(YRAP) : 2.0-2.5		
:	$R : 0.5$	
:	$: PP \rightarrow JET X$	
:	$\sqrt{S} : 7000.0 \text{ GeV}$	
PT IN GEV	D2(SIG)/DPT/DYRAP IN PB/GEV	NLO THEORY CORRECTION FACTOR
+++Collapse+++		
19.4 (bin: 18. – 21.)	$10600000 \pm 0.7\% \text{ (stat)} +20.6\%, -17.0\% \text{ (sys)}$	$1.28 \pm 27.9\%$
22.4 (bin: 21. – 24.)	$5550000 \pm 1.0\% \text{ (stat)} +18.3\%, -15.4\% \text{ (sys)}$	$1.23 \pm 25.4\%$
25.9 (bin: 24. – 28.)	$2770000 \pm 1.3\% \text{ (stat)} +16.6\%, -14.2\% \text{ (sys)}$	$1.19 \pm 22.1\%$
29.9 (bin: 28. – 32.)	$1410000 \pm 1.8\% \text{ (stat)} +15.3\%, -13.2\% \text{ (sys)}$	$1.15 \pm 18.8\%$
34.3 (bin: 32. – 37.)	$748000 \pm 2.3\% \text{ (stat)} +14.3\%, -12.5\% \text{ (sys)}$	$1.11 \pm 15.8\%$
39.8 (bin: 37. – 43.)	$351000 \pm 1.3\% \text{ (stat)} +13.5\%, -11.9\% \text{ (sys)}$	$1.09 \pm 12.9\%$
45.8 (bin: 43. – 49.)	$169000 \pm 1.9\% \text{ (stat)} +13.0\%, -11.6\% \text{ (sys)}$	$1.07 \pm 10.6\%$
52.3 (bin: 49. – 56.)	$90100 \pm 2.4\% \text{ (stat)} +12.7\%, -11.4\% \text{ (sys)}$	$1.06 \pm 8.8\%$
59.7 (bin: 56. – 64.)	$44000 \pm 1.0\% \text{ (stat)} +12.6\%, -11.3\% \text{ (sys)}$	$1.05 \pm 7.2\%$
68.6 (bin: 64. – 74.)	$21500 \pm 1.3\% \text{ (stat)} +12.6\%, -11.4\% \text{ (sys)}$	$1.04 \pm 5.9\%$
78.7 (bin: 74. – 84.)	$10300 \pm 1.9\% \text{ (stat)} +12.9\%, -11.6\% \text{ (sys)}$	$1.04 \pm 4.8\%$
90.0 (bin: 84. – 97.)	$5020 \pm 0.6\% \text{ (stat)} +13.2\%, -11.8\% \text{ (sys)}$	$1.03 \pm 3.8\%$
104.8 (bin: 97. – 114.)	$2190 \pm 0.8\% \text{ (stat)} +13.6\%, -12.2\% \text{ (sys)}$	$1.03 \pm 3.0\%$
122.7 (bin: 114. – 133.)	$896 \pm 0.4\% \text{ (stat)} +14.3\%, -12.8\% \text{ (sys)}$	$1.03 \pm 2.3\%$
142.2 (bin: 133. – 153.)	$384 \pm 0.6\% \text{ (stat)} +15.2\%, -13.5\% \text{ (sys)}$	$1.03 \pm 1.7\%$
162.7 (bin: 153. – 174.)	$171 \pm 0.6\% \text{ (stat)} +16.2\%, -14.2\% \text{ (sys)}$	$1.03 \pm 1.3\%$
184.2 (bin: 174. – 196.)	$78.7 \pm 0.8\% \text{ (stat)} +17.3\%, -15.1\% \text{ (sys)}$	$1.03 \pm 1.0\%$
207.1 (bin: 196. – 220.)	$36 \pm 0.8\% \text{ (stat)} +18.7\%, -16.1\% \text{ (sys)}$	$1.03 \pm 0.7\%$
231.6 (bin: 220. – 245.)	$16.5 \pm 1.1\% \text{ (stat)} +20.5\%, -17.4\% \text{ (sys)}$	$1.03 \pm 0.4\%$
257.5 (bin: 245. – 272.)	$7.26 \pm 1.2\% \text{ (stat)} +22.5\%, -18.9\% \text{ (sys)}$	$1.03 \pm 0.2\%$
285.0 (bin: 272. – 300.)	$3.29 \pm 1.7\% \text{ (stat)} +25.0\%, -20.6\% \text{ (sys)}$	$1.03 \pm 0.00\%$
313.9 (bin: 300. – 330.)	$1.4 \pm 2.5\% \text{ (stat)} +27.9\%, -22.5\% \text{ (sys)}$	$1.03 \pm 0.2\%$
344.8 (bin: 330. – 362.)	$0.582 \pm 3.7\% \text{ (stat)} +31.3\%, -24.7\% \text{ (sys)}$	$1.03 \pm 0.4\%$
377.3 (bin: 362. – 395.)	$0.221 \pm 5.9\% \text{ (stat)} +35.8\%, -27.4\% \text{ (sys)}$	$1.03 \pm 0.7\%$
411.2 (bin: 395. – 430.)	$0.078 \pm 9.4\% \text{ (stat)} +40.5\%, -30.0\% \text{ (sys)}$	$1.02 \pm 1.0\%$
508.1 (bin: 430. – 737.)	$0.00555 \pm 11.4\% \text{ (stat)} +52.7\%, -36.1\% \text{ (sys)}$	$1.02 \pm 1.5\%$

+++Collapse+++  
this is the place for a plot

ABS(YRAP) : 2.0-2.5




ABS(YRAP) : 2.0-2.5



Plots of separate data columns

Level of detail axis

# Improved display of data headers



- The structure of header is parsed and translated into LaTeX
- Symbols are replaced using a dictionary (that should be maintained on the HEPData side and harvested together with data)
- Javascript plugin renders formulas in the browser

# TODO

- Establish way of harvesting the dictionaries of symbols (for improved data headers display)
- Invent nicer way of breaking long lines of headers/titles
- Improve the general look&feel
- Create BibFormat for standalone HEPData records
- Establish indexes for HEPData search
- Invent way to display HEPData entries not connected to publications (virtual records?)