

Important information about REVLIMID

REVLIMID is a prescription medicine used to treat adults with multiple myeloma (MM), an advanced form of the blood cancer disease. It is administered through intravenous (IV) infusions. REVLIMID is not a chemotherapy drug, and it is administered differently than other anticancer drugs. REVLIMID should not be used in people who have certain symptoms of blood disease, such as low platelets or low red blood cells. REVLIMID should not be used in people who have certain symptoms of blood disease, such as low platelets or low red blood cells. REVLIMID should not be used in people who have certain symptoms of blood disease, such as low platelets or low red blood cells.

REVLIMID is not a blood thinner, risk of low blood counts and blood clots. REVLIMID may cause serious side effects, including:

- **Possible blood clots (venous thrombosis) or death of an unborn baby.** Venous thrombosis is a blood clot in a vein. Blood clots can lead to a heart attack or stroke. REVLIMID may cause blood clots in the lungs, which can lead to a heart attack or stroke. REVLIMID may cause blood clots in the lungs, which can lead to a heart attack or stroke.

REVLIMID is similar to the medicine hydroxyurea. REVLIMID is similar to the medicine hydroxyurea, which is used to treat certain types of blood cancer. REVLIMID may cause blood clots in the lungs, which can lead to a heart attack or stroke. REVLIMID may cause blood clots in the lungs, which can lead to a heart attack or stroke.

Female read not get pregnant. REVLIMID may cause serious side effects, including:

- **Female read not get pregnant.** REVLIMID may cause serious side effects, including:

Female who can become pregnant. REVLIMID may cause serious side effects, including:

- **Female who can become pregnant.** REVLIMID may cause serious side effects, including:

If you become pregnant while taking REVLIMID, stop taking it right away and call your healthcare provider. REVLIMID may cause serious side effects, including:

- **If you become pregnant while taking REVLIMID, stop taking it right away and call your healthcare provider.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including:

Do not have sex while taking REVLIMID. REVLIMID may cause serious side effects, including:

- **Do not have sex while taking REVLIMID.** REVLIMID may cause serious side effects, including: