

Notes in representation theory

Paul Martin

Dec 11, 2008 (printed: August 20, 2013)

Contents

1	Introduction	13
1.1	Representation theory preamble	13
1.1.1	Matrices	13
1.1.2	Notations for monoids and groups	14
1.1.3	Group representations	14
1.1.4	Unitary and normal representations	16
1.1.5	Group algebras, rings and algebras	17
1.2	Modules and representations	19
1.2.1	Examples	19
1.2.2	Simple modules and Jordan–Holder Theorem	20
1.2.3	Ideals, radicals, semisimplicities, and Artinian rings	21
1.2.4	Artin–Wedderburn Theorem	22
1.2.5	Projective modules over arbitrary rings	24
1.2.6	Structure of Artinian rings	24
1.3	Aims of representation theory	25
1.3.1	Radical series and socle of a module	25
1.3.2	The ordinary quiver of an algebra	26
1.4	Partition algebras — a quick example	29
1.4.1	Defining an algebra: by structure constants	29
1.4.2	Useful notation for set partitions	30
1.4.3	Defining an algebra: as a subalgebra	30
1.4.4	Defining an algebra: by a presentation	31
1.4.5	Exercises	31
1.5	Small categories and categories	31
1.5.1	Functors	32
1.5.2	Natural transformations and Morita equivalence	33
1.5.3	Special objects and arrows	33
1.5.4	Idempotents, Morita,	34
1.5.5	Aside: tensor products	36
1.5.6	Functor examples for module categories: globalisation	36
1.6	Modular representation theory	38
1.7	Modules and ideals for the partition algebra P_n	39
1.7.1	Ideals	39
1.7.2	Idempotents and idempotent ideals	40

1.8	Modules and ideals for T_n	40
1.8.1	Some module morphisms	41
1.8.2	Aside on Res-functors (exactness etc)	44
1.8.3	Functor examples for module categories: induction	45
1.8.4	Back to P_n	45
1.8.5	Back to T_n	46
1.8.6	The decomposition matrices	50
1.8.7	Odds and ends	50
1.9	Lie algebras	50
1.10	Eigenvalue problems	52
1.11	Notes and references	53
1.12	Exercises	53
1.12.1	Radicals	54
1.12.2	What is categorical?	54
2	Basic definitions, notations and examples	57
2.1	Preliminaries	57
2.1.1	Definition summary	57
2.1.2	Glossary	59
2.2	Elementary set theory notations and constructions	59
2.2.1	Functions	60
2.2.2	Composition of functions	60
2.2.3	Set partitions	62
2.2.4	Exercises	63
2.3	Basic tools: topology	63
2.4	Partial orders, lattices and graphs	64
2.4.1	Posets and lattices	64
2.4.2	Digraphs and graphs	66
2.5	Aside on quiver algebra characterisations of algebras	67
2.5.1	Ordinary quivers of algebras	68
2.5.2	Aside on centraliser algebras	69
2.6	Notes and references	70
3	Initial examples in representation theory	71
3.1	Initial examples in representation theory	71
3.1.1	The monoid $\text{hom}(\underline{2}, \underline{2})$	71
3.1.2	The monoid $\text{hom}(\underline{3}, \underline{3})$ and beyond	74
3.1.3	Quaternions	75
4	Reflection groups and geometry	77
4.1	Some basic geometry	77
4.1.1	Affine spaces and simplicial complexes	77
4.1.2	Hyperplane geometry, polytopes etc	78
4.2	Reflections, hyperplanes and reflection groups	80
4.2.1	Reflection group root systems	82
4.2.2	Coxeter systems and reflection groups by presentation	83

4.2.3	Some finite and hyperfinite examples and exercises	84
4.3	Reflection group chamber geometry	85
4.3.1	S_n as a reflection group, permutahedra, etc	88
4.3.2	Cayley and dual graphs, Bruhat order	88
4.4	Coxeter/Parabolic systems (W', W) and alcove geometry	89
4.5	Exercises and examples	90
4.5.1	Constructing $G_a(\mathcal{D}^-, \mathcal{D}_+)$ and $G_a(\mathcal{D}, \mathcal{D}_+)$, and beyond	90
4.5.2	Right cosets of \mathcal{D}_+ in \mathcal{D}^-	92
4.5.3	On connections of reflection groups with representation theory	93
4.6	Combinatorics of Kazhdan–Lusztig polynomials	93
4.6.1	The recursion for polynomial array $P(W'/W)$	93
4.6.2	Example	95
4.6.3	Alternative constructions: wall-alcove	99
4.7	Young graph combinatorics	99
4.7.1	Young diagrams and the Young lattice	99
4.8	Young graph via alcove geometry on $\mathbb{Z}^{\mathbb{N}}$	101
4.8.1	Nearest-neighbour graphs on \mathbb{Z}^n	102
4.8.2	Graphs on $\mathbb{Z}^{\mathbb{N}}$	103
5	Basic Category Theory	107
5.1	Categories I	107
5.1.1	Functors	109
5.1.2	Notes and Exercises (optional)	111
5.1.3	Natural transformations	111
5.2	R -linear and ab-categories	114
5.2.1	Abelian categories	115
5.3	Categories II	115
5.3.1	Adjunctions	116
5.4	Categories III	117
5.4.1	Tensor/monoidal categories	117
6	Rings in representation theory	119
6.1	Rings I	119
6.1.1	Examples	119
6.1.2	Properties of elements of a ring	120
6.2	Ideals and homomorphisms	122
6.2.1	Ring homomorphisms	122
6.2.2	Posets revisited	123
6.2.3	Properties of ideals: Artinian and Noetherian rings	123
6.2.4	Properties of ideals: Integral and Dedekind domains	123
6.3	Rings II	124
6.3.1	Order and valuation	124
6.3.2	Complete discrete valuation ring	126
6.3.3	p -adic numbers	128
6.3.4	Idempotents over the p -adics	129

7	Ring-modules	131
7.1	Ring-modules	131
7.1.1	The lattice of submodules of a module	131
7.2	R -homomorphisms and the category $R\text{-mod}$	132
7.2.1	quotients	132
7.2.2	Direct sums and simple modules	133
7.2.3	Free modules	134
7.2.4	Matrices over R and free module basis change	135
7.3	Finiteness issues	135
7.3.1	Radicals and semisimple rings	136
7.3.2	Composition series	138
7.3.3	More on chains of modules and composition series	139
7.4	Tensor product of ring-modules	139
7.4.1	Examples	140
7.4.2	R -lattices etc	141
7.5	Functors on categories of modules	141
7.5.1	Hom functors	141
7.5.2	Tensor functors and tensor-hom adjointness	143
7.5.3	Exact functors	144
7.6	Simple modules, idempotents and projective modules	145
7.6.1	Idempotents	145
7.6.2	Projective modules	148
7.6.3	Idempotent refinement	149
7.7	Structure of an Artinian ring	150
7.8	Homology, complexes and derived functors	150
7.9	More on tensor products	152
7.9.1	Induction and restriction functors	153
7.9.2	Globalisation and localisation functors	153
7.10	Morita equivalence	154
8	Algebras	155
8.1	Algebras and A -modules	155
8.2	Finite dimensional algebras over fields	156
8.2.1	Dependence on the field	156
8.2.2	Representation theory preliminaries	157
8.2.3	Structure of a finite dimensional algebra over a field	157
8.3	Cartan invariants (Draft)	158
8.3.1	Examples	159
8.3.2	Idempotent lifting revisited	160
8.3.3	Brauer reciprocity	160
8.4	Globalisation functors	162
8.4.1	Globalisation functors and projective modules	162
8.4.2	Brauer-modules in a Brauer-modular-system for A	164
8.5	On Quasi-heredity — an axiomatic framework	164
8.5.1	Definitions	164
8.5.2	Consequences for $A\text{-mod}$	165

8.5.3	Examples	166
8.6	Notes and References	166
8.7	More axiomatic frameworks	166
8.7.1	Summary of Donkin on finite dimensional algebras	166
8.7.2	Quasi-hereditary algebras	167
8.7.3	Cellular algebras	167
9	Forms, module morphisms and Gram matrices	169
9.1	Forms, module morphisms and Gram matrices (Draft)	169
9.1.1	Some basic preliminaries recalled: ordinary duality	169
9.1.2	Contravariant duality	171
9.1.3	A Schur Lemma for ‘standard’ modules	172
9.1.4	Bilinear forms	172
9.1.5	Contravariant forms on A -modules	174
9.1.6	Examples: contravariant forms	176
10	Basic representation theory of the symmetric group	179
10.1	Introduction	179
10.1.1	Integer partitions, Young diagrams and the Young lattice	180
10.1.2	Realisation of S_n as a reflection group	180
10.2	Representations of S_n from the category Set	182
10.2.1	Connection with Schur’s work and Schur functors	183
10.2.2	Idempotents and other elements in $\mathbb{Z}S_n$	185
10.2.3	Young modules	186
10.2.4	Specht modules	188
10.3	Characteristic p , Nakayama and the James abacus	189
10.4	James–Murphy theory	190
10.4.1	Murphy elements	192
10.5	Young forms for S_n irreducible representations	193
10.5.1	Hooks, diamond pairs and the Young Forms for S_n	193
10.5.2	Asides on geometry	195
10.6	Outer product and related representations of S_n	195
10.6.1	Multipartitions and their tableaux	195
10.6.2	Actions of S_n on tableaux	196
10.6.3	Generalised hook lengths and geometry	198
10.6.4	Connections to Lie theory and Yang–Baxter	198
10.7	Outer products continued — classical cases	198
10.7.1	Outer products over Young subgroups	198
10.7.2	Outer products over wreath subgroups	199
10.7.3	The Leduc–Ram–Wenzl representations	199
10.8	Finite group generalities	199
10.8.1	Characters	199

11 The Temperley–Lieb algebra	201
11.1 Ordinary Hecke algebras in brief	201
11.1.1 Geometric Braid groups	201
11.1.2 Artin braid groups	202
11.1.3 Braid group algebra quotients	203
11.1.4 Bourbaki generic algebras	204
11.2 Duality of Hecke algebras with quantum groups	204
11.3 Representations of Hecke algebras	205
11.4 Temperley–Lieb algebras from Hecke algebras	206
11.4.1 Presentation of Temperley–Lieb algebras as Hecke quotients	208
11.4.2 Tensor space representations	208
11.5 Diagram categories	208
11.5.1 Relation to quantum groups	208
11.6 Temperley–Lieb diagram algebras	209
11.6.1 TL diagram notations and definitions	210
11.6.2 Isomorphism with Temperley–Lieb algebras	211
11.6.3 TL diagram counting	211
11.6.4 Back to the TL isomorphism theorem	211
11.7 Representations of Temperley–Lieb diagram algebras	211
11.7.1 Tower approach: Preparation of small examples	213
11.8 Idempotent subalgebras, F and G functors	215
11.8.1 Aside on non-exactness of G	216
11.8.2 More general properties of F and G	217
11.8.3 Decomposition numbers	218
11.9 Decomposition numbers for the Temperley–Lieb algebra	219
11.10 Ringel dualities with U_qsl_2	219
11.10.1 Fusion of Temperley–Lieb algebras	220
12 On representations of the partition algebra	225
12.1 The partition category	225
12.1.1 Partition diagrams	226
12.1.2 Partition categories	227
12.2 Properties of partition categories	228
12.2.1 Δ -modules	229
12.3 Set partitions and diagrams	229
12.4 Representation theory	230
12.5 Representation theory via Schur algebras	232
12.5.1 Local notations	232
12.5.2 The Schur algebras	232
12.5.3 The global partition algebra as a localisation	234
12.5.4 Representation theory	234
12.5.5 Alcove geometric characterisation	239
12.5.6 More	239
12.6 Notes and references	240
12.6.1 Notes on the Yale papers on the partition algebra	240

13 On representations of the Brauer algebra	241
13.1 Context of the Brauer algebra	241
13.2 Introduction to Brauer algebra representations	241
13.2.1 Reductive and Brauer-modular representation theory	242
13.2.2 Globalisation and towers of recollement	243
13.2.3 Overview of the Chapter	244
13.3 Brauer diagrams and diagram categories	246
13.3.1 Remarks on the ground ring and Cartan matrices	248
13.4 Properties of the diagram basis	249
13.4.1 Manipulation of Brauer diagrams: lateral composition	249
13.4.2 Ket-bra diagram decomposition	249
13.5 Idempotent diagrams and subalgebras in $B_n(\delta)$	250
13.6 Appendix: Bibliographic notes	254
13.6.1 Summary	254
13.6.2 Preliminary generalities	255
13.6.3 Auslander: rep. thy. of small additive categories (as if rings)	255
13.6.4 Towers of recollement	256
13.7 Appendix: Overview of following Chapters	257
13.7.1 Blocks and the block graph $G_\delta(\lambda)$	258
13.7.2 Embedding the vertex set of $G_\delta(\lambda)$ in \mathbb{R}^N	258
13.7.3 Reflection group action on \mathbb{R}^N	259
13.7.4 Decomposition data: Hypercubical decomposition graphs	260
14 General representation theory of the Brauer algebra	263
14.1 Initial filtration of the left regular module	263
14.2 Brauer Δ -modules	265
14.2.1 Symmetric group Specht modules (a quick reminder)	265
14.2.2 Brauer Δ -module constructions	265
14.2.3 Brauer Δ -module examples	267
14.2.4 Simple head conditions for Δ -modules	267
14.2.5 Brauer algebra representations: The base cases	269
14.2.6 The case $k \supseteq \mathbb{Q}$	269
14.3 Δ -Filtration of projective modules	269
14.3.1 Some character formulae	269
14.3.2 General preliminaries	269
14.3.3 A Δ -filtration theorem	270
14.3.4 On simple modules, labelling and Brauer reciprocity	271
14.4 Globalisation functors	272
14.4.1 Preliminaries: \otimes versus category composition	273
14.4.2 G -functors	274
14.4.3 Idempotent globalisation	276
14.4.4 Simple head(Δ) conditions revisited using G -functors	278
14.4.5 Simple modules revisited using G -functors	279
14.5 Induction and restriction	280
14.6 Characters and Δ -filtration factors over \mathbb{C}	281
14.6.1 Aside on case $\delta = 0$	281

14.6.2	The main case	282
14.6.3	The n -independence of $(P(\lambda) : \Delta(\mu))$	285
15	Complex representation theory of the Brauer algebra	287
15.1	Blocks of $B_n(\delta)$	287
15.1.1	Blocks I: actions of central elements on modules	288
15.1.2	Easy Lemmas and the DWH Lemma	290
15.2	Blocks II: δ -balanced pairs of Young diagrams	292
15.2.1	Towards a constructive treatment: δ -charge and δ -skew	292
15.2.2	Sections and rims of Young diagrams	294
15.2.3	A constructive treatment: π -rotations and δ -pairs	294
15.2.4	Conditions for a width-1 δ -skew to have a section	295
15.2.5	Connections and properties of δ -skews and δ -pairs	297
15.2.6	The graph $G_\delta(\lambda)$	299
15.3	Brauer algebra Δ -module maps and block relations	300
15.4	On the block graph $G_\delta(\lambda)$	300
15.4.1	Embedding the vertex set of $G_\delta(\lambda)$ in $\mathbb{R}^{\mathbb{N}}$	301
15.4.2	Reflection group \mathcal{D} acting on $\mathbb{R}^{\mathbb{N}}$	302
15.4.3	Constructing graph morphisms for $G_\delta(\lambda)$: combinatorial approach	303
15.4.4	The graph G_{even}	306
15.5	Graph isomorphisms, via geometrical considerations	307
15.5.1	Dual graphs and alcove geometry	307
15.5.2	Group \mathcal{D} action on $\mathbf{e}_\delta(\Lambda)$	308
15.5.3	The graph isomorphism	310
15.5.4	Aside on alternative reflection group actions on Λ	313
15.6	The decomposition matrix theorem	314
15.6.1	Decomposition data: Hypercubical decomposition graphs	314
15.6.2	The main Theorem	318
15.6.3	Hypercubical decomposition graphs: examples	318
15.6.4	Hypercubical decomposition graphs: tools	320
15.7	Embedding properties of δ -blocks in Λ	322
15.7.1	The Relatively-regular-step Lemma	324
15.7.2	The Reflection Lemmas	324
15.7.3	The Embedding Theorem and the $\text{Proj}_\Lambda \text{Ind-}$ functor	326
15.7.4	The generic projective lemma	327
15.7.5	Properties of δ -pairs and rim-end removable boxes	329
15.7.6	The singularity lemma	331
15.8	Proof of The Decomposition Matrix Theorem	334
15.8.1	The generic inductive-step lemma	334
15.8.2	The rank-2 inductive-step lemma	334
15.8.3	Example for the rank-2 inductive step	338
15.9	Some remarks on the block graph	341
15.9.1	Yet more	341

16 Properties of Brauer block graphs	343
16.1 Kazhdan–Lusztig polynomials revisited	343
16.1.1 Overview	343
16.1.2 The recursion for $P(W'/W)$	343
16.2 The reflection group action \mathcal{D} on \mathbb{R}^N	344
16.3 Solving the polynomial recursion for $P(\mathcal{D}/\mathcal{D}_+)$	344
16.3.1 Hypercubes h^a revisited	345
16.3.2 Kazhdan–Lusztig polynomials for $\mathcal{D}/\mathcal{D}_+$	346
16.4 Related notes and open problems	349
16.5 Block labelling weights	351
16.6 Changing δ	351
17 More Brauer algebra modules	353
17.1 King’s polynomials and other results	353
17.2 Connection between King polynomials and D/A alcove geometry	356
17.3 Leduc–Ram representations of Brauer algebras and other results	357
17.3.1 Brauer diamonds	357
17.3.2 Leduc–Ram representations	359
17.3.3 Geometrical realisation	362
17.4 On ‘untruncating’ Leduc–Ram representations	369
17.5 Truncating Leduc–Ram representations (old version!)	370
17.6 JOBS	371
18 Example: the Temperley–Lieb algebra again	373
18.1 More on categories of modules	373
18.1.1 More fun with F and G functors	373
18.1.2 Saturated towers	375
18.1.3 Quasi-heredity of planar diagram algebras	375
19 Lie groups	377
19.1 Introduction (to algebraic groups etc)	377
19.2 Preliminaries	379
19.3 Lie group	380
19.3.1 Example: $SU(2)$ ‘polynomial’ representations	381
19.3.2 Lie algebra	382

Chapter 1

Introduction

ch:basic

Chapters 1 and 2 give a brief introduction to representation theory, and a review of some of the basic algebra required in later Chapters. A more thorough grounding may be achieved by reading the works listed in §1.11: *Notes and References*.

Section 1.1 (upon which later chapters do not depend) attempts to provide a sketch overview of topics in the representation theory of finite dimensional algebras. In order to bootstrap this process, we use some terms without prior definition. We assume you know what a vector space is, and what a ring is (else see Section 2.1.1). For the rest, either you know them already, or you must intuit their meaning and wait for precise definitions until after the overview.

1.1 Representation theory preamble

s:ov

ss:matrices1

1.1.1 Matrices

Let $M_{m,n}(R)$ denote the additive group of $m \times n$ matrices over a ring R , with additive identity $0_{m,n}$. Let $M_n(R)$ denote the ring of $n \times n$ matrices over R . Define a block diagonal composition (matrix direct sum)

$$\begin{aligned} \oplus : M_m(R) \times M_n(R) &\rightarrow M_{m+n}(R) \\ (A, A') &\mapsto A \oplus A' = \begin{pmatrix} A & 0_{m,n} \\ 0_{n,m} & A' \end{pmatrix} \end{aligned}$$

(sometimes we write \oplus for matrix/exterior \oplus for disambiguation). Define Kronecker product

$$\otimes : M_{a,b}(R) \times M_{m,n}(R) \rightarrow M_{am,bn}(R) \tag{1.1} \quad \text{eq:kronecker12}$$

$$(A, B) \mapsto \begin{pmatrix} a_{11}B & a_{12}B & \dots \\ a_{21}B & a_{22}B & \dots \\ \vdots & & \end{pmatrix} \tag{1.2}$$

In general $A \otimes B \neq B \otimes A$, but (if R is commutative then) for each pair A, B there exists a pair of permutation matrices S, T such that $S(A \otimes B) = (B \otimes A)T$ (if A, B square then $T = S$ — the *intertwiner* of $A \otimes B$ and $B \otimes A$).

1.1.2 Notations for monoids and groups

(See §2.2 for a more extended discussion of set theory notations.)

de:freemonoid

(1.1.1) Given a set S , then the *free monoid* S^* is the set of words in the alphabet S , together with the operation of juxtaposition: $a * b = ab$. (Note associativity.)

pr:f1

(1.1.2) If M is a monoid with generating subset in bijection with S then there is a map $f : S^* \rightarrow M$.

(1.1.3) Let ρ be a relation on set S , a monoid. Then ρ is *compatible* with monoid S if $(s, t), (u, v) \in \rho$ implies $(su, tv) \in \rho$.

We write $\rho\#$ for the intersection of all compatible equivalence relations ('congruences') on S containing ρ .

(1.1.4) If ρ is an equivalence relation on set S then S/ρ denotes the set of classes of S under ρ .

(1.1.5) If ρ is a congruence on semigroup S then S/ρ has a semigroup structure by:

$$\rho(a) * \rho(b) = \rho(a * b)$$

(Exercise: check well-definedness and associativity.)

(1.1.6) For set S finite we can define a monoid by presentation.

...

de:solvableg

(1.1.7) A group G is *solvable* if there is a chain of subgroups $\dots G_i \subset G_{i+1} \dots$ such that $G_i \leq G_{i+1}$ (normal subgroup) and G_{i+1}/G_i is abelian.

(1.1.8) EXAMPLE. $(\mathbb{Z}, +)$ and S_3 are solvable; S_5 is not.

1.1.3 Group representations

de:rep

(1.1.9) A matrix representation of a group G over a commutative ring R is a map

$$\rho : G \rightarrow M_n(R) \tag{1.3} \quad \text{try345}$$

such that $\rho(g_1 g_2) = \rho(g_1) \rho(g_2)$. In other words it is a map from the group to a different system, which nonetheless respects the extra structure (of multiplication) in some way. The study of representations — models of the group and its structure — is a way to study the group itself.

(1.1.10) The map ρ above is an example of the notion of representation that generalises greatly. A mild generalisation is the representation theory of R -algebras that we shall discuss, but one could go further. Physics consists in various attempts to model or represent the observable world. In a model, Physical entities are abstracted, and their behaviour has an image in the behaviour of the model. We say we understand something when we have a model or representation of it mapping to something we understand (better), which does not wash out too much of the detailed behaviour.

de:repIII

(1.1.11) Representation theory itself seeks to classify and construct representations (of groups, or other systems). Let us try to be more explicit about this.

(I) Suppose ρ is as above, and let S be an arbitrary invertible element of $M_n(R)$. Then one immediately verifies that

$$\rho_S : G \rightarrow M_n(R) \quad (1.4) \quad \boxed{\text{aaas}}$$

$$g \mapsto S\rho(g)S^{-1} \quad (1.5)$$

is again a representation.

(II) If ρ' is another representation (by $m \times m$ matrices, say) then

$$\rho \oplus \rho' : G \rightarrow M_{m+n}(R) \quad (1.6) \quad \boxed{\text{dsum}}$$

$$g \mapsto \rho(g) \oplus \rho'(g) \quad (1.7)$$

is yet another representation.

(III) For a finite group G let $\{g_i : i = 1, \dots, |G|\}$ be an ordering of the group elements. Each element g acts on G , written out as this list $\{g_i\}$, by multiplication from the left (say), to permute the list. That is, there is a permutation $\sigma(g)$ such that $gg_i = g_{\sigma(g)(i)}$. This permutation can be recorded as a matrix,

$$\rho_{\text{Reg}}(g) = \sum_{i=1}^{|G|} \epsilon_{i \sigma(g)(i)}$$

(where $\epsilon_{ij} \in M_{|G|}(R)$ is the i, j -elementary matrix) and one can check that these matrices form a representation, called the *regular representation*.

Clearly, then, there are unboundedly many representations of any group. However, these constructions also carry the seeds for an organisational scheme...

(1.1.12) Firstly, in light of the ρ_S construction, we only seek to classify representations *up to isomorphism* (i.e. up to equivalences of the form $\rho \leftrightarrow \rho_S$).

Secondly, we can go further (in the same general direction), and give a cruder classification, by *character*. (While cruder, this classification is still organisationally very useful.) We can briefly explain this as follows.

Let c_G denote the set of classes of group G . A *class function* on G is a function that factors through the natural set map from G to the set c_G . Thus an R -valued class function is completely specified by a c_G -tuple of elements of R (that is, an element of the set of maps from c_G to R , denoted R^{c_G}). For each representation ρ define a *character* map from G to R

$$\chi_\rho : G \rightarrow R \quad (1.8) \quad \boxed{\text{eq:ch1}}$$

$$g \mapsto \text{Tr}(\rho(g)) \quad (1.9)$$

(matrix trace). Note that this map is fixed up to isomorphism. Note also that this map is a class function. Fixing G and varying ρ , therefore, we may regard the character map instead as a map χ_- from the collection of representations to the set of c_G -tuples of elements of R .

Note that pointwise addition equips R^{c_G} with the structure of abelian group. Thus, for example, the character of a sum of representations isomorphic to ρ lies in the subgroup generated by the character of ρ ; and $\chi_{\rho \oplus \rho'} = \chi_\rho + \chi_{\rho'}$ and so on.

We can ask if there is a small set of representations whose characters ‘ \mathbb{N}_0 -span’ the image of the collection of representations in R^{c_G} . (We could even ask if such a set provides an R -basis for

R^{c_G} (in case R a field, or in a suitably corresponding sense — see later). Note that $|c_G|$ provides an upper bound on the size of such a set.)

(1.1.13) Next, conversely to the direct sum result, suppose $R_1 : G \rightarrow M_m(R)$, $R_2 : G \rightarrow M_n(R)$, and $V : G \rightarrow M_{m,n}(R)$ are set maps, and that a set map $\rho_{12} : G \rightarrow M_{m+n}(R)$ takes the form

$$\rho_{12}(g) = \begin{pmatrix} R_1(g) & V(g) \\ 0 & R_2(g) \end{pmatrix} \quad (1.10) \quad \boxed{\text{eq:plus}}$$

(a matrix of matrices). Then ρ_{12} a representation of G implies that both R_1 and R_2 are representations. Further, $\chi_{\rho_{12}} = \chi_{R_1} + \chi_{R_2}$ (i.e. the character of ρ_{12} lies in the span of the characters of the smaller representations). Accordingly, if the isomorphism class of a representation contains an element that can be written in this way, we call the representation *reducible*.

(1.1.14) For a finite group over $R = \mathbb{C}$ (say) we shall see later that there are only a finite set of ‘irreducible’ representations needed (up to equivalences of the form $\rho \leftrightarrow \rho_S$) such that every representation can be built (again up to equivalence) as a direct sum of these; and that all of these irreducible representations appear as direct summands in the regular representation.

We have done a couple of things to simplify here. Passing to a field means that we can think of our matrices as recording linear transformations on a space with respect to some basis. To say that ρ is equivalent to a representation of the form ρ_{12} above is to say that this space has a G -subspace (R_1 is the representation associated to the subspace). A representation is irreducible if there is no such proper decomposition (up to equivalence). A representation is *completely reducible* if for every decomposition $\rho_{12}(g)$ there is an equivalent identical to it except that $V(g) = 0$ — the direct sum.

Theorem [Mashke] Let ρ be a representation of a finite group G over a field K . If the characteristic of K does not divide the order of G , then ρ is completely reducible.

Corollary Every complex irreducible representation of G is a direct summand of the regular representation.

Representation theory is more complicated in general than it is in the cases to which Mashke’s Theorem applies, but the notion of irreducible representations as fundamental building blocks survives in a fair degree of generality. Thus the question arises:

Over a given R , what are the irreducible representations of G (up to $\rho \leftrightarrow \rho_S$ equivalence)?

There are other questions, but as far as physical applications (for example) are concerned, this is arguably the main interesting question.

(1.1.15) Examples: In this sense, of constructing irreducible representations, the representation theory of the symmetric groups S_n over \mathbb{C} is completely understood! (We shall review it.) On the other hand, over other fields we do not have even so much as a conjecture as to how to organise the statement of a conjecture! So there is work to be done.

1.1.4 Unitary and normal representations

A complex representation ρ of a group G in which every $\rho(g)$ is unitary is a *unitary representation* (see e.g. Boerner [11, III§6]). A representation equivalent to a unitary representation is *normal*.

(1.1.16) **THEOREM.** *Let G be a finite group. Every complex representation of G is normal. Every real representation of G is equivalent to a real orthogonal representation.*

1.1.5 Group algebras, rings and algebras

de: lset (1.1.17) For a set S , a map $\psi : G \times S \rightarrow S$ (written $\psi(g, s) = gs$ where no ambiguity arises) such that

$$(gg')s = g(g's),$$

equips S with the property of *left G -set*.

(1.1.18) For example, for a group $(G, *)$, then G itself is a left G -set by left multiplication: $\psi(g, s) = g * s$. (Cf. (1.1.11)(III).)

On the other hand, consider the map $\psi_r : G \times G \rightarrow G$ given by $\psi_r(g, s) = s * g$. This obeys $\psi_r(g * g', s) = s * (g * g') = (s * g) * g' = \psi_r(g', \psi_r(g, s))$. This ψ_r makes G a *right G -set*: in the notation of (1.1.17) we have

$$(gg')s = g'(gs). \quad (1.11) \quad \text{eq: rset}$$

The map $\psi_- : G \times G \rightarrow G$ given by $\psi_-(g, s) = g^{-1} * s$ obeys $\psi_-(g * g', s) = (g * g')^{-1} * s = (g'^{-1} * g^{-1}) * s = g'^{-1} * (g^{-1} * s) = \psi_-(g', \psi_-(g, s))$. This ψ_- makes G a *right G -set*.

(1.1.19) Remark: When working with R a *field* it is natural to view the matrix ring $M_n(R)$ as the ring of linear transformations of vector space R^n expressed with respect to a given ordered basis. The equivalence $\rho \leftrightarrow \rho_S$ corresponds to a change of basis, and so working up to equivalence corresponds to demoting the matrices themselves in favour of the underlying linear transformations (on R^n). In this setting it is common to refer to the linear transformations by which G acts on R^n as the representation (and to spell out that the matrices are a *matrix* representation, regarded as arising from a choice of ordered basis).

Such an action of a group G on a set makes the set a G -set. However, given that R^n is a set with extra structure (in this case, a vector space), it is a small step to want to try to take advantage of the extra structure.

(1.1.20) For example, continuing for the moment with R a field, we can define RG to be the R -vector space with basis G (see Exercise 1.12.1), and define a multiplication on RG by

$$\left(\sum_i r_i g_i \right) \left(\sum_j r'_j g_j \right) = \sum_{ij} (r_i r'_j) (g_i g_j) \quad (1.12) \quad \text{groupalgmult}$$

which makes RG a ring (see Exercise 1.12.2). One can quickly check that

$$\rho : RG \rightarrow M_n(R) \quad (1.13)$$

$$\sum_i r_i g_i \mapsto \sum_i r_i \rho(g_i) \quad (1.14)$$

extends a representation ρ of G to a representation of RG in the obvious sense. Superficially this construction is extending the use we already made of the multiplicative structure on $M_n(R)$, to make use not only of the additive structure, but also of the particular structure of ‘scalar’ multiplication (multiplication by an element of the centre), which plays no role in representing the group multiplication *per se*. The construction *also* makes sense at the G -set/vector space level, since linear transformations support the same extra structure.

de:RG-module

(1.1.21) The same formal construction of RG works when R is an arbitrary commutative ring (called the *ground ring*), except that RG is not then a vector space. Instead it is called (in respect of the vector-space-like aspect of its structure) a *free R -module with basis G* . The idea of matrix representation goes through unchanged. If one wants a generalisation of the notion of G -set for RG to act on, the additive structure is forced from the outset. This is called a (*left*) *RG -module*. This is, then, an abelian group $(M, +)$ with a suitable action of RG defined on it: $r(x + y) = rx + ry$, $(r + s)x = rx + sx$,

$$(rs)x = r(sx), \quad (1.15)$$

eq:lmodule

$1x = x$ ($r, s \in RG$, $x, y \in M$), just as the original vector space R^n was.

What is new at this level is that such a structure may not have a basis (a *free* module has a basis), and so may not correspond to any class of matrix representations.

(1.1.22) EXERCISE. Construct an RG -module without basis.

(Possible hints: 1. Consider $R = \mathbb{Z}$, G trivial, and look at §7.3. 2. Consider the ideal $\langle 2, x \rangle$ in $\mathbb{Z}[x]$.)

From this point the study of representation theory may be considered to include the study of both matrix representations and modules.

(1.1.23) What other kinds of systems can we consider representation theory for?

A natural place to start studying representation theory is in Physical modeling. Unfortunately we don't have scope for this in the present work, but we will generalise from groups at least as far as rings and algebras.

The generalisation from groups to *group algebras* RG over a commutative ring R is quite natural as we have seen. The most general setting within the ring-theory context would be the study of arbitrary ring homomorphisms from a given ring. However, if one wants to study this ring by studying its modules (the obvious generalisation of the RG -modules introduced above) then the parallel of the matrix representation theory above is the study of modules that are also free modules over the centre, or some subring of the centre. (For many rings this accesses only a very small part of their structure, but for many others it captures the main features. The property that *every* module over a commutative ring is free holds if and only if the ring is a field, so this is our most accessible case. We shall motivate the restriction shortly.) This leads us to the study of *algebras*.

To introduce the general notion of an algebra, we first write $\text{cen}(A)$ for the centre of a ring A

$$\text{cen } A = \{a \in A \mid ab = ba \ \forall b \in A\}$$

de: alg1

(1.1.24) An algebra A (over a commutative ring R), or an R -algebra, is a ring A together with a homomorphism $\psi : R \rightarrow \text{cen}(A)$, such that $\psi(1_R) = 1_A$.

Examples: Any ring is a \mathbb{Z} -algebra. Any ring is an algebra over its centre. The group ring RG is an R -algebra by $r \mapsto r1_G$. The ring $M_n(R)$ is an R -algebra.

Let $\psi : R \rightarrow \text{cen}(A)$ be a homomorphism as above. We have a composition $R \times A \rightarrow A$:

$$(r, a) = ra = \psi(r)a$$

so that A is a left R -module with

$$r(ab) = (ra)b = a(rb) \quad (1.16)$$

eq: alg12

Conversely any ring which is a left R -module with this property is an R -algebra.

(1.1.25) An R -representation of A is a homomorphism of R -algebras

$$\rho : A \rightarrow M_n(R)$$

(1.1.26) The study of RG depends heavily on R as well as G . The study of such R -algebras takes a relatively simple form when R is an algebraically closed field; and particularly so when that field is \mathbb{C} . We shall aim to focus on these cases. However there are significant technical advantages, even for such cases, in starting by considering the more general situation. Accordingly we shall need to know a little ring theory, even though general ring theory is not the object of our study.

Further, as we have said, neither applications nor aesthetics restrict attention to the study of representations of groups and their algebras. One is also interested in the representation theory of more general algebras.

1.2 Modules and representations

The study of algebra-modules and representations for an algebra over a field has some special features, but we start with some general properties of modules over an arbitrary ring R . (NB, this topic is covered in more detail in Chapter 7, and in our reference list §1.11.)

A module over an arbitrary ring R is defined exactly as for a module over a group ring — (1.1.21) (NB our ring R here has taken over from RG not the ground ring, so there is no requirement of commutativity).

We assume familiarity with exact sequences of modules. See Chapter 7, or say [75], for details.

1.2.1 Examples

ex:ring001

(1.2.1) EXAMPLE. Consider the ring $R = M_n(\mathbb{C})$. This acts on the space $M = M_{n,1}(\mathbb{C})$ of n -component column matrices by matrix multiplication from the left. Thus M is a left R -module.

ex:ring01

(1.2.2) EXAMPLE. Consider the ring $R = M_2(\mathbb{C}) \oplus M_3(\mathbb{C})$. A general element in R takes the form

$$r = r_1 \oplus r_2 = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \oplus \begin{pmatrix} e & f & g \\ h & i & j \\ k & l & m \end{pmatrix} \in R$$

Here, $M = \mathbb{C}\{(1,0)^T, (0,1)^T\} = \left\{ \begin{pmatrix} x \\ y \end{pmatrix} \mid x, y \in \mathbb{C} \right\}$ is a left R -module with r acting by left-

multiplication by $r_1 = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$; $M'' = M_2(\mathbb{C})$ is a left module with r acting in the same way;

$M' = \left\{ \begin{pmatrix} s \\ t \\ u \end{pmatrix} \mid s, t, u \in \mathbb{C} \right\}$ is a left module with r acting by r_2 ; and M'' is also a right module by right-multiplication by r_1 .

Note that the subset of M'' of form $\begin{pmatrix} x & 0 \\ y & 0 \end{pmatrix}$ is a left submodule.

(1.2.3) Our next example concerns a commutative ring, where the distinction between left and right modules is void. Consider the ring \mathbb{Q} . This acts on $(\mathbb{R}, +)$ in the obvious way, making $(\mathbb{R}, +)$ a

left (or right) \mathbb{Q} -module. Here $(\mathbb{Q}, +) \subset (\mathbb{R}, +)$ is a submodule — indeed it is a minimal submodule, in the sense that any submodule containing 1 must contain this one. Note that this submodule (generated by 1) and the submodule generated by $\sqrt{2} \in \mathbb{R}$ do not intersect non-trivially. Note that here there is no ‘maximal submodule’.

exe:funny1

(1.2.4) EXERCISE. Consider the ring R_χ of matrices of form $\begin{pmatrix} q & 0 \\ x & y \end{pmatrix} \in \begin{pmatrix} \mathbb{Q} & 0 \\ \mathbb{R} & \mathbb{R} \end{pmatrix}$. (Note that this is not an algebra over \mathbb{R} and is not a finite-dimensional algebra over \mathbb{Q} .) Determine some submodules of the left-regular module.

Answer: (See also (1.2.18).) Consider the submodules of the left-regular module R_χ generated by a single element. Firstly:

$$\begin{pmatrix} q & 0 \\ x & y \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ y & 0 \end{pmatrix}$$

— that is, there is a submodule of matrices of the form on the right, with $y \in \mathbb{R}$. Note that this submodule itself has no non-trivial submodules (indeed it is a 1-d \mathbb{R} -vector space). Then:

$$\begin{pmatrix} q & 0 \\ x & y \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & y \end{pmatrix}$$

is again a 1-d \mathbb{R} -vector space. Finally consider

$$\begin{pmatrix} q & 0 \\ x & y \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} q & 0 \\ x & 0 \end{pmatrix}$$

Note that the submodule generated here, while not an \mathbb{R} -vector space, itself has the first case above as a submodule. The quotient has no non-trivial submodule (and indeed is a 1-d \mathbb{Q} -vector space).

1.2.2 Simple modules and Jordan–Holder Theorem

(1.2.5) A left R -module (for R an arbitrary ring) is *simple* if it has no non-trivial submodules. (See §7.2 for more details.)

In Example 1.2.2 both M and M' are simple; while R is a left-module for itself which is not simple, and M'' is also not simple.

(1.2.6) Let M be a left R -module. A *composition series* for M is a sequence of submodules $M = M_0 \supset M_1 \supset M_2 \supset \dots \supset M_l = 0$ such that the section M_i/M_{i+1} is simple.

In particular if a composition series of M exists for some l then M_{l-1} is a simple submodule.

The sections of a composition series for M (if such exists) are *composition factors*. Their multiplicities up to isomorphism are called *composition multiplicities*. Write $(M : L)$ for the multiplicity of simple L .

th:JH

(1.2.7) **Theorem.** (Jordan–Holder) Let M be a left R -module. (A) All composition series for M (if such exist) have the same factors up to permutation; and (B) the following are equivalent:

- (I) M has a composition series;
- (II) every ascending and descending chain of submodules of M stops (these two stopping conditions separately are known as *ACC* and *DCC*);
- (III) every sequence of submodules of M can be refined to a composition series.

Proof. See §7.3.2.

1.2.3 Ideals, radicals, semisimplicities, and Artinian rings

de:semisim (1.2.8) A module M is *semisimple* if equal to the sum of its simple submodules.

de:ideal0 (1.2.9) A *left ideal* of R is a submodule of R regarded as a left-module for itself. A subset $I \subset R$ that is both a left and right ideal is a (*two-sided*) *ideal* of R . A *nil ideal* of R is a (left/right/two-sided) ideal in which every element r is nilpotent (there is an $n \in \mathbb{N}$ such that $r^n = 0$). A *nilpotent ideal* of R is an ideal I for which there is an $n \in \mathbb{N}$ such that $I^n = 0$. (So I nilpotent implies I nil.)

de:JacRad0 (1.2.10) The *Jacobson radical* of ring R is the intersection of its maximal left ideals.

(1.2.11) Ring R itself is a *semisimple ring* if its Jacobson radical vanishes.

Remark: This term is sometimes used for a ring that is semisimple as a left-module for itself. The two definitions coincide under certain conditions (but not always). See later.

de:lss (1.2.12) For the moment we shall say that a ring R is *left-semisimple* if it is semisimple as a left-module ${}_R R$ (cf. e.g. Adamson [2, §22]). There is then a corresponding notion of *right-semisimple*, however: THEOREM. A ring is right-semisimple if and only if left-semisimple.

The next Theorem is not trivial to show:

THEOREM. The following are equivalent:

- (I) ring R is left-semisimple.
- (II) every module is semisimple (as in (1.2.8)).
- (III) every module is projective (every short exact sequence splits — see also 1.2.32).

(1.2.13) THEOREM. The Jacobson radical of ring R contains every nil ideal of R . ■¹

Remark: In general the Jacobson radical is not necessarily a nil ideal. (But see Theorem 1.2.19.)

(1.2.14) An element $r \in R$ is *quasiregular* if $1_R + r$ is a unit. The element $r' = (1_R + r)^{-1} - 1$ is then the *quasiinverse* of r . (See e.g. Faith [?].)

(1.2.15) THEOREM. If J is the Jacobson radical of ring R and $r \in J$ then r is quasiregular. ■

(1.2.16) Ring R is *Artinian* (resp. *Noetherian*) if it has the DCC (resp. ACC, as in (1.2.7)) as a left and as a right module for itself.

th:fdalgebraa (1.2.17) Example: THEOREM. A finite dimensional algebra over a field is Artinian.

Proof. A left- (or right-)ideal here is a finite dimensional vector space. A proper subideal necessarily has lower dimension, so any sequence of strict inclusions terminates. □

de:funny ring (1.2.18) Aside: We say more about chain conditions in §7.3. Here we briefly show by an example that the left/right distinction is not vacuous (although, as the contrived nature of the example perhaps suggests, it will be largely irrelevant for us in practice). Consider the ring R_χ of matrices of form $\begin{pmatrix} q & 0 \\ x & y \end{pmatrix} \in \begin{pmatrix} \mathbb{Q} & 0 \\ \mathbb{R} & \mathbb{R} \end{pmatrix}$ as in (1.2.4). (Note that this is not an algebra over \mathbb{R} and is not a finite-dimensional algebra over \mathbb{Q} .) We claim that R_χ is Artinian and Noetherian as a left module for itself. However we claim that there are an infinite chain of right-submodules of R_χ as a

¹We shall use ■ to mean that the proof is left as an exercise.

right-module for itself between $\begin{pmatrix} 0 & 0 \\ \mathbb{Q} & 0 \end{pmatrix}$ and $\begin{pmatrix} 0 & 0 \\ \mathbb{R} & 0 \end{pmatrix}$. Thus R_χ is left Artinian but not right Artinian.

To prove the left-module claims one can show that all possible candidates are \mathbb{R} -vector spaces, and finite dimensional. To prove the infinite chain claim, recall that one can form a set of infinitely many \mathbb{Q} -linearly-independent elements in \mathbb{R} (else \mathbb{R} is countable!). Order the beginning of this set as $B_n = \{1, b_1, b_2, \dots, b_n\}$ (we have taken the first element as 1 WLOG), for $n = 0, 1, 2, \dots$. We have $\mathbb{Q}B_0 = \mathbb{Q}$ and $\mathbb{Q}B_n \subset \mathbb{Q}B_{n+1}$ for all n , thus an infinite ascending chain. On the other hand there is an inverse limit B of the sequence B_n contained in \mathbb{R} (perhaps this requires Zorn's Lemma/the axiom of choice!), so we can define a sequence B^n by eliminating 1 then b_1 and so on from $B = B^0$, giving an infinite descending chain $\mathbb{Q}B^n \supset \mathbb{Q}B^{n+1}$.

(1.2.19) THEOREM. If ring R Artinian then the Jacobson radical is the maximal two-sided nilpotent ideal of R (i.e. it is nilpotent and contains all other nilpotent ideals). ■

th:nilrad0

(1.2.20) THEOREM. If ring R Artinian then ideal I nil implies I nilpotent. ■

(1.2.21) THEOREM. If a ring is left-semisimple (as in 1.2.12) then it is (left and right) Artinian and left Noetherian, and is semisimple (i.e. has radical zero). ■ (See e.g. [2, Th.22.2].)

1.2.4 Artin–Wedderburn Theorem

(1.2.22) **Theorem.** (Schur's Lemma) Suppose M, M' are nonisomorphic simple R modules. Then the ring $\text{hom}_R(M, M)$ of R -module homomorphisms from M to itself is a division ring; and $\text{hom}_R(M, M') = 0$.

lem:Schur

Proof. (See also 7.2.11.) Let $f \in \text{hom}_R(M, M)$. M simple implies $\ker f = 0$ and $\text{im } f = M$ or 0 , so f nonzero is a bijection and hence has an inverse. Now let $g \in \text{hom}_R(M, M')$. M simple implies $\ker g = 0$ and M' simple implies $\text{im } g = M = M'$ or zero, so $g = 0$. □

(1.2.23) EXAMPLE. Let us return to ring R and module M from Example 1.2.2. In this case $\text{hom}_R(M, M) \subset \text{hom}_{\mathbb{C}}(M, M)$, and $\text{hom}_{\mathbb{C}}(M, M)$ is all \mathbb{C} -linear transformations, so realised by $M_2(\mathbb{C})$ in the given basis. We see that $\text{hom}_R(M, M)$ is the subset that commute with the action of R . This is the centre of $M_2(\mathbb{C})$, which is $\mathbb{C}1_2$, which is isomorphic to \mathbb{C} .

ex:ring01a

On the other hand, $\text{hom}(M, M')$ is realised by matrices $\tau \in M_{3,2}(\mathbb{C})$:

$$\begin{pmatrix} \cdot & \cdot \\ \cdot & \cdot \\ \cdot & \cdot \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \cdot \\ \cdot \\ \cdot \end{pmatrix}$$

Here in $\text{hom}_R(M, M')$ we look for matrices τ such that

$$\begin{pmatrix} \cdot & \cdot \\ \cdot & \cdot \\ \cdot & \cdot \end{pmatrix} r \begin{pmatrix} x \\ y \end{pmatrix} = r \begin{pmatrix} \cdot & \cdot \\ \cdot & \cdot \\ \cdot & \cdot \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$

for all r , that is

$$\begin{pmatrix} \cdot & \cdot \\ \cdot & \cdot \\ \cdot & \cdot \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} e & f & g \\ \cdots & & m \end{pmatrix} \begin{pmatrix} \cdot & \cdot \\ \cdot & \cdot \\ \cdot & \cdot \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$

but since a, b, c, d, e, \dots, m may be varied independently we must have $\tau = 0$.

(1.2.24) REMARK. Cf. the occurrence of the division ring in the general proof with the details in our example. We can consider the occurrence of the division ring in Schur's Lemma as one of the main reasons for studying division rings alongside fields.

de:ringdirectsum

(1.2.25) Suppose that ring R has a decomposition of 1 into orthogonal central idempotents: $1 = \sum_i e_i$. Then each $R_i = Re_i$ is an ideal of R and a ring with identity e_i . In this case we say that R is a *ring direct sum* of the rings R_i , and write $R = \oplus_i R_i$. (Note that this is consistent with Example (1.2.2).)

th:AWI

(1.2.26) **Theorem.** (Artin–Wedderburn) Suppose R is semisimple and Artinian. Then R is a direct sum of rings of form $M_{n_i}(K_i)$ ($i = 1, 2, \dots, l$, some l) where each K_i is a division ring.

Proof. Exercise. (See also §7.3 or e.g. Benson [7, Th.1.3.5].) ■

(1.2.27) Suppose M', M'' submodules of R -module M . They *span* M if $M' + M'' = M$; and are *independent* if $M' \cap M'' = 0$. If they are both independent and spanning we write

$$M = M' \oplus M''$$

((*module*) *direct sum*). A module is *indecomposable* if it has no proper direct sum decomposition.

(1.2.28) EXAMPLE. Suppose $e^2 = e \in R$, then

$$Re \oplus R(1 - e) = R \tag{1.17}$$

eq:projid1

as left-module.

Proof. For $r \in R$, $r = re + r(1 - e)$ so $Re + R(1 - e) = R$; and $re \in R(1 - e)$ implies $re = re(1 - e) = 0$. □

(1.2.29) Note that a central idempotent decomposition of 1_R leads to an ideal decomposition of R ; while an arbitrary orthogonal idempotent decomposition of 1_R leads to a left-module decomposition of R .

Evidently a central idempotent decomposition is an orthogonal idempotent decomposition, but such a decomposition may be refinable once the central condition is relaxed. The matrix algebra $M_n(K)$ has the n elementary matrix idempotents $\{e_i^n\}_i$, which are orthogonal and such that

$$1_{M_n(K)} = \sum_{i=1}^n e_i^n$$

so this gives us one way to refine the central idempotent decomposition of 1_R in a semisimple Artinian ring (as in 1.2.26) to an (ordinary) orthogonal idempotent decomposition:

$$1_R = \sum_{i=1}^l \sum_{j=1}^{n_i} e_j^{n_i}$$

(here the first sum needs interpretation — it comes formally from the direct sum). We say more about this in §1.5.4.

Krull

(1.2.30) **Theorem.** (Krull–Schmidt) If R is Artinian then as a left-module for itself it is a finite direct sum of indecomposable modules; and any two such decompositions may be ordered so that the i -th summands are isomorphic.

Proof. Exercise. (See also §7.3.2.)

1.2.5 Projective modules over arbitrary rings

ss:proj0001

(1.2.31) If $x : M \rightarrow M'$, $x' : M' \rightarrow M$ are R -module homomorphisms such that $x \circ x' = 1_{M'}$ then x is a *split surjection* (and x' a split injection).

de:iproj

(1.2.32) An R -module is *projective* if it is a direct summand of a free module (an R -module with a linearly independent generating set).

(1.2.33) EXAMPLE. $e^2 = e \in R$ implies left-module Re projective, since it is a direct summand of free module R , by (1.17).

th:proj intro

(1.2.34) **Theorem.** TFAE

(I) R -module P is projective;

(II) whenever there is an R -module surjection $x : M \rightarrow M'$ and a map $y : P \rightarrow M'$ then there is a map $z : P \rightarrow M$ such that $x \circ z = y$;

(III) every R -module surjection $t : M \rightarrow P$ splits.

Proof. Exercise. (See also §7.6.)

1.2.6 Structure of Artinian rings

:structArtinian1

th:ASTI

(1.2.35) If R is Artinian and J_R its radical then R/J_R is semisimple so by (1.2.26):

$$R/J_R = \oplus_{i \in l(R)} M_{n_i}(R_i)$$

for some set $l(R)$, numbers n_i and division rings R_i . There is a simple R/J_R -module (L_i say) for each factor, so that *as a left module*

$$R/J_R \cong \oplus_i n_i L_i$$

(i.e. n_i copies of L_i). There is a corresponding decomposition of 1 in R/J_R :

$$1 = \sum_i e_i$$

into orthogonal idempotents. One may find corresponding idempotents in R itself (see later) so that $1 = \sum_i e'_i$ there. This gives left module decomposition

$$R = \oplus_i n_i P_i$$

where (by (1.2.30)) the P_i s are a complete set of indecomposable projective modules up to isomorphism.

(See also §7.7.)

th:ASTIcaveat

(1.2.36) Caveat: Note that the above does not say, for an k -algebra over a field, that $\dim L_i = n_i$. For example, the \mathbb{Q} -algebra $A = \mathbb{Q}\{1, x\}/(x^2 - 2)$ is a simple module for itself of dimension 2. That is, Artin–Wedderburn here is rather trivial: $A = M_1(A)$. A sufficient condition for $\dim L_i = n_i$ is that k is algebraically closed.

1.3 Aims of representation theory

So, what are the aims of representation theory? For Artinian algebras they are, broadly and roughly speaking, to describe the (finite dimensional) modules, and their homomorphisms. One might also be looking for representations (i.e. module bases) with special properties (perhaps motivated by physics). But in any case, it is worth being a bit more specific about this ‘description’.

Typically, to start with, one is looking for *invariants* — properties of modules that would be manifested by any isomorphic algebra; so that one can, say, determine from representation theory whether two algebras are isomorphic (or more easily, that two algebras are *not* isomorphic).

An example of an invariant would be the number of isomorphism classes of simple modules — this would be the same for any isomorphic algebra... See (1.4.10) for a specific example.

de:fund inv

(1.3.1) Given an Artinian algebra R (let us say specifically a finite dimensional algebra over an algebraically closed field k , so that each $R_i = k$ in (1.2.35)), we are called on

(A0) to determine a suitable indexing set $l(R)$ as in (1.2.35),

(AI) to compute the fundamental invariants $\{n_i : i \in l(R)\}$,

(AII) to give a construction of the simple modules L_i ,

(AIII) to compute composition multiplicities for the indecomposable projective modules P_i ,

(AIV) to compute Jordan-Holder series for the modules P_i .

(AV) to compute some further invariants (see e.g. (1.3.9) below).

(1.3.2) Note that (AI) contains (A0), and completely determines the maximal semisimple quotient algebra up to isomorphism (by the Artin–Wedderburn Theorem). Aim (AII) is not an invariant, so does not have a unique answer; but having at least one such construction is clearly desirable in studying an algebra (and any answer for (AII) contains (AI)).

Of course there are unboundedly many nonisomorphic algebras with the same maximal semisimple quotient in general, so we need more information to classify non-semisimple algebras.

The aim (AIII) is an invariant, and tells us more about a non-semisimple algebra. Aim (AIV) contains (AIII). But still, (AIV) is not enough to classify algebras in general. It is very useful partial data, however. And we will usually consider this to be ‘enough’ for most purposes (applications, for example). We will say a little next about further (and possibly complete) invariants; before returning to study the above aims in detail.

(1.3.3) At a further level, we might also try the following. To investigate the isomorphism classes of indecomposable modules (beyond projective modules).

(1.3.4) Some invariants are invariants of isomorphism classes of algebras. Some are invariants of ‘Morita’ equivalence classes of algebras (see §1.5.2). This latter is a weaker (but very useful) notion. The number $l(R)$ is an invariant of Morita equivalence. The multiset $\{n_i\}$ is an invariant of isomorphism.

1.3.1 Radical series and socle of a module

(1.3.5) Fix an algebra A . Given an A -module M , its *radical* $\text{Rad}(M)$ is the intersection of maximal submodules. The *radical series* of M is

$$M \supset \text{Rad } M \supset \text{Rad Rad } M \supset \dots$$

The sections $\text{Rad}^i M / \text{Rad}^{i+1} M$ are the *radical layers*. In particular

$$\text{Head}(M) = M / \text{Rad } M$$

$$\text{Shoulder}(M) = \text{Rad } M / \text{Rad}^2 M = \text{Head}(\text{Rad } M)$$

pr:mradm

(1.3.6) PROPOSITION. (I) Module M is semisimple (of finite length) iff Artinian and $\text{Rad } M = 0$.
 (II) If a module M is Artinian then $M / \text{Rad } M$ is semisimple. ■

(1.3.7) The *socle* $\text{Soc}(M)$ of a module is the maximal semisimple submodule. One can form socle layers: $\text{Soc}(M)$, $\text{Soc}(M/\text{Soc}(M))$, $\text{Soc}((M/\text{Soc}(M))/\text{Soc}(M/\text{Soc}(M)))$, ... in the obvious way. These layers do not agree, in general, with the reverse of the radical layers; but the lengths of sequences agree if defined.

(1.3.8) Let A be a finite dimensional algebra over an algebraically closed field. (Then the radical series of any finite dimensional module terminates; and the sections are semisimple modules, by Prop.1.3.6.) Here we put indexing set $l(A) = \Lambda(A)$. For the indecomposable projective A -modules $\{P_i\}_{i \in \Lambda(A)}$ then

$$\{P_i\}_{i \in \Lambda(A)} \leftrightarrow \{S_i = \text{Head}(P_i)\}_{i \in \Lambda(A)}$$

is a bijection between indecomposable projectives and simples. In general we have

$$\text{Head}(M) \cong \bigoplus_{i \in \Lambda(A)} \underbrace{m_i^0(M)}_{\text{multiplicity}} S_i$$

$$\text{Shoulder}(M) \cong \bigoplus_{i \in \Lambda(A)} m_i^1(M) S_i$$

(and so on) for some multiplicities $m_i^l(M) \in \mathbb{N}_0$.

A *radical Loewy diagram* of an Artinian module M gives the radical layers:

$$\begin{array}{ccccccc} M & = & S_{0,1} & S_{0,2} & S_{0,3} & \cdots & S_{0,l_0} \\ & & S_{1,1} & S_{1,2} & S_{1,3} & S_{1,4} & \cdots & S_{1,l_1} \\ & & S_{2,1} & S_{2,2} & \cdots & & \\ & & \cdots & & & & \end{array}$$

(the multiset of simple modules $\{S_{0,1}, S_{0,2}, \dots\}$ encodes $\text{Head}(M)$ and so on). We give some examples in §1.3.2.

1.3.2 The ordinary quiver of an algebra

ss:quiv00

de:quiv1

(1.3.9) The ordinary quiver of an algebra. (...See §2.5 for details.)

How *do* we classify finite dimensional algebras (over an algebraically closed field) up to isomorphism; or up to Morita equivalence?

(1.3.10) An algebra is *connected* if it has no proper central idempotent. Every algebra is isomorphic to a direct sum of connected algebras, so it is enough to classify connected algebras (and then, for an arbitrary algebra, give its connected components).

de:basicalg0

(1.3.11) An algebra is *basic* if every simple module is one-dimensional. (See also (1.5.30).) Every algebra is Morita equivalent to (i.e. has an equivalent module category to) a basic algebra. So it is enough to classify basic connected algebras.

(1.3.12) The *Ext-matrix* $\mathcal{M}(A)$ of algebra A is given by the ‘shoulder data’

$$\mathcal{M}(A)_{ij} = m_i^1(P_j)$$

A necessary condition for algebra isomorphism $A \cong B$ is that there is an ordering of the index sets such that $\mathcal{M}(A) = \mathcal{M}(B)$.

The *Ext-quiver* or *ordinary quiver* $Q(A)$ of algebra A is the matrix $\mathcal{M}(A)$ expressed as a graph. Note that $Q(A)$ is connected as a graph if A is connected as an algebra. Isomorphism $A \cong B$ implies isomorphic Ext-quivers, but not v.v.. However one can characterise any connected basic algebra A up to isomorphism using a quotient of the *path algebra* $kQ(A)$ of $Q(A)$ (given a quiver Q , then kQ is the k -algebra with basis of walks on Q and composition on walks by concatenation where defined, and zero otherwise²), as we describe in §??. Specifically we have the following.

(1.3.13) THEOREM. [48, §4.3] *For any connected basic algebra A there is an ideal I_A in $kQ(A)$ (contained in $I_{\geq 2}$ and containing $I_{\geq m}$ for some m) such that*

$$A \cong kQ(A)/I_A$$

Proof. First note that there is a surjective algebra homomorphism $\Psi : kQ(A) \rightarrow A$. The walks of length-0 pass to a set of idempotents such that $P_i = Ae_i$. The walks of length-1 from i to j pass to a basis for $e_i J_A e_j / e_i J_A^2 e_j$.

Next we need to show that the kernel of Ψ has the required form. See e.g. [7, Prop.1.2.8]. ■

(1.3.14) Thus we can determine (characterise up to isomorphism) such a connected basic A by computing $Q(A)$ and then giving elements of $kQ(A)$ that generate I_A . (Note however that generators for I_A are not unique in general.)

More generally then, one can determine an arbitrary algebra A by giving the corresponding data for its connected components; together with the dimensions of the simple modules.

(1.3.15) Given $A \cong kQ(A)/I_A$, we can recover structural data about the indecomposable projective modules as follows. Write e_a for the path of length 0 from vertex a (sometimes we just write $a = e_a$ for this). This is an idempotent in $kQ(A)$. Then

$$P_a = Ae_a$$

(identifying A with $kQ(A)/I_A$ here without loss of generality). Thus a basis for P_a is the set of all paths from a ‘up to the quotient’. This is the path of length 0 (corresponding to the head); and all the paths of length 1 (the shoulder); and some paths of length 2; and so on.

Note that (the image of) $I_{\geq 1}$ lies in the radical of kQ/I_A , since the m -th power lies in $I_{\geq m} \equiv 0$. Hence the image of $I_{\geq 1}$ is the radical.

(1.3.16) Let us give some low-dimensional examples of algebras of form Q/I_A , where $I_A \subset I_{\geq 2}$ and $I_A \supset I_{\geq m}$ for some m .

²Note that walks of length at least l span an ideal in kQ . Write $I_{\geq l}$ for this ideal.

For Q a single point then kQ is one-dimensional and $I_{\geq 2} = 0$. Indeed any kQ with $I_{\geq 1} = 0$ is semisimple — the quiver is just a collection of points. Let us give some non-semisimple examples. For

$$\begin{array}{c} \textcircled{a} \\ \uparrow u \\ \textcircled{a} \end{array} \quad \text{with relation } u^2 = 0$$

we have a 2d algebra with 1 simple S_a . The corresponding projective P_a is $P_a = Aa = k\{a, ua\}$ (it terminates here since $aua = ua = u$ and $u^2a = 0$ and so on), in which $k\{ua\}$ is a submodule (of, in a suitable sense, length-1 elements) isomorphic to S_a . That is, a radical Loewy diagram for P_a is

$$P_a = \begin{array}{c} S_a \\ S_a \end{array}$$

There is a 1-simple algebra in each dimension obtained by replacing $u^2 = 0$ by $u^d = 0$.

Alternatively in 3d, we can take the quiver with 1 vertex and two loops u, v , together with the relations $uu = uv = vu = vv = 0$. The quiver

$$a \xleftarrow{x} b \quad \text{with no relations}$$

(again $I_{\geq 2} = 0$ here) gives another 3d algebra, this time with 2 simples.

The quiver

$$\begin{array}{ccc} & x & \\ a & \xrightarrow{\quad} & b \\ & s & \end{array} \quad \text{with } sx = 0$$

has basis $\{a, b, xa, sb, xsb\}$. (Note that the given relation is sufficient to make kQ/I_A finite, but otherwise an arbitrary choice for an example here.) The indecomposable projective Aa is generated by walks out of a : $a, xa, sxa = 0$, that is, it terminates after one step. The projective $P_b = Ab$ has walks $b, sb, xsb, xsxb = 0$.

(1.3.17) What about this?:


$$\begin{array}{ccccc} & x_{ab} & & x_{bc} & \\ a & \xrightarrow{\quad} & b & \xrightarrow{\quad} & c \\ & x_{ba} & & x_{cb} & \end{array} \quad \text{with } x_{bc}x_{ab}, x_{ba}x_{cb}, x_{ba}x_{ab} \text{ and } x_{ab}x_{ba} - x_{cb}x_{bc} \text{ in } I_A.$$

(These relations are another arbitrary finite choice here. However these particular relations will appear ‘in the wild’ later.) We have $P_a = Aa = k\{a, x_{ab}a\}$. Next $P_b = Ab = k\{b, x_{ba}b, x_{bc}b, x_{ab}x_{ba}b\}$. Finally $P_c = Ac$. Note the submodule structure of P_b . As ever there is a unique maximal submodule $\text{Rad } P_b = k\{x_{ba}b, x_{bc}b, x_{ab}x_{ba}b\}$. The intersection of the maximal submodules of this, in turn, is spanned by $x_{ab}x_{ba}b$. Thus the radical layers of the projectives look like this:

$$\begin{array}{ccccc} P_a = & S_a & P_b = & S_b & P_c = & S_c \\ & S_b & & S_a & S_c & S_b \\ & & & S_b & & S_c \end{array}$$

REMARK. This case exemplifies a very interesting point: that the presence of a simple module as a composition factor for a module always allows for a corresponding homomorphism from the indecomposable projective cover of that simple module. Here in particular there is no homomorphism from S_a to P_b , say, but there is a homomorphism from P_a to P_b . See later.

(1.3.18) What about this?:



Determine some conforming relations to make a finite quotient of kQ

1.4 Partition algebras — a quick example

ss:pa0001

Just so that we can have a glimpse of what is coming up, we use the partition algebra to generate some examples. The objective can be considered to be determining the data (A0-III) from (1.3.1) for various Artinian algebras. (The aim is to illustrate various tools for doing this kind of thing.) We follow directly the argument in [87].

We start by very briefly recalling the partition algebra construction but, essentially, we assume for now that you know the definition and some notations for the partition algebras (else see §2.2.3 and §12, or [87]).

Implicit in this section are a number of exercises, requiring the proof of the various claims.

1.4.1 Defining an algebra: by structure constants

Given a commutative ring k , how do we define an algebra over k ? One way is to give a basis and the ‘structure constants’ — the multiplication rule on this basis.


(1.4.1) EXAMPLE. A group algebra for a given group is a very simple example of this.

de:Pn

(1.4.2) Fix a commutative ring k , and $\delta \in k$. For S a set, P_S is the set of partitions of S . Let $n, m \in \mathbb{N}$. Define $N(n, m) = \{1, 2, \dots, n, 1', 2', \dots, m'\}$. We recall the *partition algebra*. Firstly, the partition algebra $P_n = P_n(\delta)$ over k is an algebra with a basis $P_{N(n, n)}$. Thus the rank of P_n as a free k -module is the Bell number B_{2n} . In particular if k is a field then P_n is Artinian.

We may draw a partition of $N(n, m)$ as an (n, m) -graph. An (n, m) -graph is a drawing of a graph in a box with vertex set including $N(n, m)$ on the frame — unprimed $1, 2, \dots, n$ left-to-right on the northern edge; primed $1', 2', \dots, m'$ on the southern. That is, if d is such a graph, then $\pi_{n, m}(d) \in P_{N(n, m)}$ is the partition with $i, j \in N(n, m)$ in the same part if they are in the same connected component in d . Any d such that $\pi_{n, n}(d) = p$, and such that every vertex is in a connected component with an element of $N(n, n)$, serves as a picture of p .

A one-picture summary (!) of the P_n diagram calculus (composition of partitions defined via concatenating diagrams) is:



(1.18) eq:Ppic1

Note that a connected component in such a graph is *internal* if it has vertices on neither external edge; and that a graph d with l internal components denotes an element $\delta^l \pi_{n, n}(d)$ of P_n .

1.4.2 Useful notation for set partitions

de:ideibp

(1.4.3) Given a partition p of some subset of $N(n, n)$, take p^* to be the image under toggling the prime.

Setting $v = \{\{1\}\}$ we have $v^* = \{\{1'\}\}$. Set $1 = \{\{1, 1'\}\}$, $u = \{\{1\}, \{1'\}\}$, $\cup = \{\{1, 2\}\}$, $\Gamma = \{\{1, 2, 1'\}\}$, $\square = \{\{1, 2, 1', 2'\}\}$, and $\sigma = \{\{1, 2'\}, \{2, 1'\}\}$. Define partition $p_1 \otimes p_2$ by side-by-side concatenation of diagrams (and hence renumbering the p_2 factor as appropriate). We have

$$u = v \otimes v^*$$

For given n we define $u_i \in P_{N(n, n)}$ by

$$u_1 := u \otimes 1 \otimes 1 \otimes \dots \otimes 1, \quad u_2 := 1 \otimes u \otimes 1 \otimes \dots \otimes 1, \quad \text{and so on.}$$

de:pnnotations

(1.4.4) Let $P_{n, m} := P_{N(n, m)}$. We say a part in $p \in P_{n, m}$ is *propagating* if it contains both primed and unprimed elements. Write $P_{n, l, m}$ for the subset of $P_{n, m}$ with l propagating parts; and $P_{n, m}^l$ for the subset of $P_{n, m}$ with at most l propagating parts. Thus

$$P_{n, m}^l = \sqcup_{l'=0}^l P_{n, l', m} \quad \text{and} \quad P_{n, m} = \sqcup_{l'=0}^n P_{n, l', m}.$$

E.g. $P_{2, 2, 2} = \{1 \otimes 1, \sigma\}$, $P_{2, 1, 1} = \{v \otimes 1, 1 \otimes v, \Gamma\}$, $P_{2, 0, 0} = \{v \otimes v, \cup\}$ and

$$P_{2, 1, 2} = P_{2, 1, 1} P_{1, 1, 2} = \{u \otimes 1, 1 \otimes u, v \otimes 1 \otimes v^*, v^* \otimes 1 \otimes v, \Gamma\Gamma^*, \dots\}.$$

Note that $P_{n, n, n}$ spans a multiplicative subgroup:

$$P_{n, n, n} \cong S_n \tag{1.19} \quad \text{eq:PnSnsb}$$

(1.4.5) We have $P_0 \cong k$, $P_1 = k\{1, u\}$ and

$$P_2 = k(P_{2, 2, 2} \cup P_{2, 1, 2} \cup P_{2, 0, 2}) = k(P_{2, 2, 2} \cup P_{2, 1, 2} \cup \{\cup \otimes \cup^*, (v \otimes v) \otimes \cup^*, (v \otimes v)^* \otimes \cup, u \otimes u\}).$$

We have $u^2 = \delta u$ (but see Ch.12 for the definition of the algebra/category composition) and $v^*v = \delta \emptyset$ and $vv^* = u$.

1.4.3 Defining an algebra: as a subalgebra

de:TLn

(1.4.6) We will also use the subalgebra

$$T_n = T_n(\delta)$$

of the k -algebra P_n with basis $T_{n, n} \subset P_{n, n}$ of non-crossing pair partitions (following [84, §9.5]). For example, $e := \cup \otimes \cup^* \in T_{2, 2}$; and for given n , $e_1 := e \otimes 1 \otimes 1 \otimes \dots \otimes 1 \in T_{n, n}$.

(1.4.7) EXERCISE. Show that there is such a subalgebra. And also a subalgebra with a basis of arbitrary pair-partitions.

(1.4.8) REMARK. Historically the subalgebra of P_n with basis of pair-partitions comes first [13] — the *Brauer algebra* B_n . We look at this in §?? et seq.

1.4.4 Defining an algebra: by a presentation

(1.4.9) EXERCISE. Determine generators and relations for P_n .

de:TLieb

(1.4.10) For k a commutative ring, and $\delta \in k$, define the Temperley–Lieb algebra TL_n as the quotient of the free k -algebra generated by the symbols U_1, U_2, \dots, U_{n-1} by the relations

$$U_i^2 = \delta U_i$$

$$U_i U_{i\pm 1} U_i = U_i$$

$$U_i U_j = U_j U_i \quad |i - j| \neq 1$$

Thus for example TL_2 has basis $\{1, U_1\}$; while $TL_3 = k\{1, U_1, U_2, U_1 U_2, U_2 U_1\}$ as a k -space. Note in the case TL_2 that the obvious bijection from this basis/generating set to $\{1, e\}$ extends to an isomorphism $TL_2 \cong T_2$. We have the following.

(1.4.11) THEOREM. (See e.g. [84, Co.10.1]) *Fix a commutative ring k and $\delta \in k$. For each n , $TL_n \cong T_n$. ■*

1.4.5 Exercises

(1.4.12) PROPOSITION. Assuming δ a unit,

$$P_{n-1} \cong u_1 P_n u_1 \tag{1.20}$$

eq:PUPU

$$P_n / P_n u_1 P_n \cong k S_n. \tag{1.21}$$

eq:PPUPx

■

Our idea is to determine the representation theory of P_n (over a suitable algebraically closed field k) inductively from that of P_m for $m < n$, using (1.20). To this end we need to connect the two algebras.

(1.4.13) PROPOSITION. Assuming δ a unit,

$$T_{n-2} \cong e_1 T_n e_1 \tag{1.22}$$

eq:UTU2

$$T_n / T_n e_1 T_n \cong k \tag{1.23}$$

eq:TTeT1

■

1.5 Small categories and categories

ss:cat0001

See §5.1 for more details. Categories are useful from at least two different perspectives in representation theory. One is in the idea of de-emphasising modules in favour of the (existence of) morphisms between them. Another is in embedding our algebraic structures (our objects of study) in yet more general settings.

A *small category* is a triple $(A, A(-, -), \circ)$ consisting of a set A (of ‘objects’); and for each element $(a, b) \in A \times A$ a set $A(a, b)$ (of ‘arrows’); and for each element $(a, b, c) \in A^{\times 3}$ a composition:

$A(a, b) \times A(b, c) \rightarrow A(a, c)$, satisfying associativity and identity conditions (for each a there is a 1_a in $A(a, a)$ such that $1_a \circ f = f = f \circ 1_b$ whenever these make sense).

(A *category* is a similar structure allowing larger classes of objects and arrows.)

(1.5.1) Example: A monoid is a category with one object.

(1.5.2) Example: $A = \mathbb{N}$ and $A(m, n)$ is $m \times n$ matrices over a ring R .

(1.5.3) Example: A is a set of R -modules and $A(M, N)$ is the set of R -module homomorphisms from M to N . (The category $R\text{-mod}$ is the category of all left R -modules.)

(1.5.4) The product in (1.18) generalises to a category \mathbf{P} in an obvious way, with object set \mathbb{N}_0 . There is a corresponding \mathbf{T} subcategory.

(1.5.5) We may construct an ‘opposite’ category A^o from category A , with the same object class, by setting $A^o(a, b) = A(b, a)$ and reversing the compositions.

1.5.1 Functors

(1.5.6) A *functor* is a map between (small) categories that preserves composition and identities.

de:functoreg0001

(1.5.7) Example: (I) If R is a ring and $e^2 = e \in R$ then there is a map $F_e : R\text{-mod} \rightarrow eRe\text{-mod}$ given by $M \mapsto eM$ that extends to a functor.

de:homfunctintro

(1.5.8) (II) If R is a ring and N a left R -module then there is a map

$$\text{Hom}(N, -) : R\text{-mod} \rightarrow \mathbb{Z}\text{-mod}$$

given by $M \mapsto \text{Hom}(N, M)$. This extends to a functor by $L \xrightarrow{f} M \mapsto (N \xrightarrow{g} L \mapsto N \xrightarrow{f \circ g} M)$.

de:homfunctproj

(1.5.9) The functor $\text{Hom}(N, -)$ has some nice properties. Consider a not-necessarily short-exact sequence $0 \rightarrow M' \xrightarrow{\mu} M \xrightarrow{\nu} M'' \rightarrow 0$ and its not-necessarily exact image

$$0 \rightarrow \text{Hom}(N, M') \xrightarrow{\mu_N = \text{Hom}(N, \mu)} \text{Hom}(N, M) \xrightarrow{\nu_N = \text{Hom}(N, \nu)} \text{Hom}(N, M'') \rightarrow 0.$$

$$N \xrightarrow{f} M' \quad \mapsto \quad N \xrightarrow{\mu \circ f} M$$

We can ask (i) if exactness at M' implies $\ker \mu_N = 0$; (ii) if exactness at M implies $\text{im } \mu_N = \ker \nu_N$; (ii') if $\nu \circ \mu = 0$ implies $\nu_N \circ \mu_N = 0$; (iii) if exactness at M'' implies $\text{im } \nu_N = \text{Hom}(N, M'')$?

(i) Since μ injective, $\mu \circ f = \mu \circ g$ implies $f = g$. But then $\mu \circ f = 0$ implies $f = 0$, so $\ker \mu_N = 0$.

(ii) See (7.5.6). (The answer is yes if exact at M' and M .)

(ii') $\text{Hom}(N, \nu) \circ \text{Hom}(N, \mu) = \text{Hom}(N, \nu \circ \mu) = 0$.

(iii) This does not hold in general. However if N is projective then by Th.1.2.34(II), given exactness at M'' , every $\gamma \in \text{Hom}(N, M'')$ can be expressed $\nu \circ g$ for some $g \in \text{Hom}(P, M)$, so then (iii) holds.

We will give some more examples shortly — see e.g. (1.5.10).

ex:functy

(1.5.10) Let $\psi : A \rightarrow B$ be a map of algebras over k . We define functor

$$\text{Res}_\psi : B\text{-mod} \rightarrow A\text{-mod}$$

by $\text{Res}_\psi M = M$, with action of $a \in A$ given by $am = \psi(a)m$ for $m \in M$; and by $\text{Res}_\psi f = f$ for $f : M \rightarrow N$.

We need to check that Res_ψ extends to a well-defined functor, i.e. that every B -module map $f : M \rightarrow N$ is also an A -module map. We have $bf(m) = f(bm)$ for $b \in B$ and $m \in M$. Consider $af(m) = \psi(a)f(m) = f(\psi(a)m)$, where the second identity holds since $\psi(a) \in B$. Finally $f(\psi(a)m) = f(am)$ and we are done.

See §1.8.2 for properties of Res_ψ .

(1.5.11) In order to develop a useful notion of equivalence of categories we need the notion of a natural transformation — a map between functors.

1.5.2 Natural transformations and Morita equivalence

ss:ME0

For now see (5.1.26) for natural transformations. A natural isomorphism is a natural transformation whose underlying maps are isomorphisms.

Two categories A, B are equivalent if there are functors $F : A \rightarrow B$ and $G : B \rightarrow A$ such that the composites FG and GF are naturally isomorphic to the corresponding identity functors.

1.5.3 Special objects and arrows

(1.5.12) An arrow f is *epi* if $gf = g'f$ implies $g = g'$ (see e.g. Mitchell [?]).

Given a category \mathcal{A} we write $A \xrightarrow{f} B$ if f is epi.

(1.5.13) An arrow f is *mono* if $fg = fg'$ implies $g = g'$.

Given a category \mathcal{A} we write $A \xrightarrow{f} B$ if f is mono.

If $A \xrightarrow{f} B$ then we say A is a *subobject* of B .

(1.5.14) Next we should define the notions of isomorphism; isomorphic subobject; and balanced category.

de:projincat1

(1.5.15) An object P is *projective* if for every $P \xrightarrow{h} B$ and $A \xrightarrow{f} B$ then $h = ff'$ for some $P \xrightarrow{f'} A$. (Cf. (1.2.34)(II).)

(1.5.16) A category \mathcal{A} *has enough projectives* if there is an $P \xrightarrow{f} A$, with P projective, for each object A .

de:zeroobject

(1.5.17) An object O in category \mathcal{A} is a *zero object* if every $\mathcal{A}(M, O)$ and $\mathcal{A}(O, M)$ contains a single element.

If there is a unique zero object we denote it 0 . In this case we also write $M \xrightarrow{0} 0$ and $0 \xrightarrow{0} M$ for all the ‘zero-arrows’ (even though they are distinct); and $M \xrightarrow{0} N$ for the arrow that factors through 0 .

de:kernelI

(1.5.18) Here we suppose that \mathcal{A} has a unique zero-object.

A *prekernel* of $A \xrightarrow{f} B$ is any pair $(K, K \xrightarrow{k} A)$ such that $fk = 0$.

A *kernel* of $A \xrightarrow{f} B$ is a prekernel $(K, K \xrightarrow{k} A)$ such that if $(K', K' \xrightarrow{k'} A)$ is another prekernel then there is a unique $K' \xrightarrow{g} K$ such that $kg = k'$.

(1.5.19) Note that if $(K, K \xrightarrow{k} A)$ is a kernel of f then k is mono, and K is an isomorphic subobject of A to every other kernel object of f (see later).

Exercise: consider the existence and uniqueness of kernels.

(1.5.20) Next we should define normal categories and exact categories; define exact sequences. —FINISH THIS SECTION!!!—

(1.5.21) A category of modules has a lot of extra structure and special properties compared to a generic category (see Freyd [45] or §?? for details). For example: (EI) The arrow set $A(M, N) = \text{Hom}(M, N)$ is an abelian group; composition of arrows is bilinear. (An *additive* functor between such categories respects this extra structure.) (SII) There is a unique object 0 such that $\text{Hom}(M, 0) \cong \text{Hom}(0, M) \cong \{0\}$ for all M (by $0 : M \rightarrow 0$ we mean this zero-arrow — an abuse of notation!). (SIII) Given objects M, N there is a categorical notion of an object $M \oplus N$, and these objects exist. (SIV) There is a function *ker* associating to each arrow $f \in \text{Hom}(M, N)$ an object K_f and an arrow $k_f \in \text{Hom}(K_f, A)$ such that $f \circ k_f = 0$ (in the sense above), and (K_f, k_f) is in a suitable sense universal (see later).

This extra structure is useful, and warrants the treatment of module categories almost separately from generic categories. This raises the question of what aspects of representation theory are ‘categorical’ — i.e. detectable from looking at the category alone, without probing the objects and arrows as modules and module morphisms per se.

For example, the property of projectivity is categorical. (Exercise. Hint: consider $\text{Hom}(P, -)$ and short exact sequences.) The property of an object being a set is not categorical (although this concreteness is a safe working assumption for module categories, fine details of the nature of this set are certainly not categorical).

(1.5.22) Two categories are equivalent if there are functors between them whose composite is in a suitable sense isomorphic to the identity functor. We talk about making this precise later. For now we will rather aim to build some illustrative examples.

de:adjointI

(1.5.23) Consider functors $C \xrightleftharpoons{F}_G C'$. Then (F, G) is an *adjoint pair* if for each suitable object pair M, N there are natural bijections $\text{Hom}(FM, N) \mapsto \text{Hom}(M, GN)$.

ss:xxid

1.5.4 Idempotents, Morita, ...

We started by thinking about matrix representations of groups, and this has led us naturally to consider modules over algebras. Two components of this progression have been (i) the passage to natural new algebraic structures (from groups to rings to algebras) on which to study representation theory; and (ii) the organisation of representations into equivalence classes (de-emphasising the basis). Representation theory studies algebras by studying the structure preserving maps between algebras (a map from the algebra under study to a known algebra gives us the modules for the known algebra as modules for the new algebra). We could go further and de-emphasise the modules in favour of the maps between them. This is one route into using ‘category theory’.

(1.5.24) Let A be an algebra over k and $e^2 = e \in A$. The *Peirce decomposition* (or Pierce decomposition! [30, 32, §6]) of A is

$$A = eAe \oplus (1 - e)Ae \oplus eA(1 - e) \oplus (1 - e)A(1 - e)$$

or

$$A = \oplus_{i,j} e_i A e_j$$

where $e_1 = e$ and $e_2 = 1 - e$. (Question: What algebraic structures are being identified here? This is an identification of vector spaces; but the algebra multiplication is also respected. On the other hand not every summand on the right is unital.)

This decomposition is non-trivial if $1 = e + (1 - e)$ is a non-trivial decomposition. Set $A(i, j) = e_i A e_j$. These components are not-necessarily-unital ‘algebras’, and non-unit-preserving subalgebras of A . The cases $A(i, i)$ are unital, with identity e_i .

Can we study A by studying the algebras $A(i, i)$?

(1.5.25) EXAMPLE. Consider $M_3(\mathbb{C})$ and the idempotent $e_{11} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$. We have the corresponding vector space decomposition (not confusing \oplus with $\oplus \cdot$)

$$\begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} = \begin{pmatrix} a_{11} & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \oplus \begin{pmatrix} 0 & a_{12} & a_{13} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \oplus \begin{pmatrix} 0 & 0 & 0 \\ a_{21} & 0 & 0 \\ a_{31} & 0 & 0 \end{pmatrix} \oplus \begin{pmatrix} 0 & 0 & 0 \\ 0 & a_{22} & a_{23} \\ 0 & a_{32} & a_{33} \end{pmatrix}$$

(which is not necessarily a particularly interesting decomposition, but see later).

(1.5.26) If we can further decompose e into orthogonal idempotents then there is a corresponding further Peirce decomposition. This decomposition process terminates when some $e = e_\pi$ has no decomposition in A (it is ‘primitive’). What special properties does $e_\pi A e_\pi$ have then?

(1.5.27) An orthogonal decomposition of 1 into primitive idempotents is called a ‘complete’ orthogonal decomposition.

For examples see §8.3.1.

(1.5.28) Aside: Let $1 = \sum_{i \in H} e_i$ be an orthogonal idempotent decomposition, and extend the definition of $A(i, j)$ to this case. Note that we have a composition $A(i, j) \times A(k, l) \rightarrow A(i, l)$ given by $a \circ b = ab$ in A . But in particular $ab = 0$ unless $j = k$. Thinking along these lines we see that the orthogonal idempotent decomposition of $1 \in A$ gives rise to a category (see §1.5, §5.1) ‘hiding’ in A . The category is $A_H = (H, A(i, j), \circ)$.

th:eRe-Re1

(1.5.29) THEOREM. If a ring R is left or right Artinian then it has a complete orthogonal idempotent decomposition of 1, $1 = \sum_{i=1}^l e_i$ say, with $e_i R e_i$ a local ring.

If $e_i R e_i$ is local then e_i is primitive and $R e_i$ is indecomposable projective. ■

de:basicalgebra

(1.5.30) An Artinian ring R , with complete set $\{e_1, e_2, \dots, e_l\}$ of orthogonal idempotents, is *basic* if $R e_i \cong R e_j$ as left- R -modules implies $i = j$. (Cf. also (1.3.11).)

(1.5.31) EXAMPLE. The k -subalgebra of $M_2(k)$ given by $A_{1,1} = \left\{ \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} \mid a, b \in k \right\}$ has a complete set $\{e_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, e_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}\}$. One easily checks that $A_{1,1} e_1 \not\cong A_{1,1} e_2$ (consider the action of e_1 on each side, say), so $A_{1,1}$ is basic.

On the other hand $M_2(k)$ has the same complete set, but $M_2(k) e_1 \cong M_2(k) e_2$, so $M_2(k)$ is not basic.

(1.5.32) One can check that if a finite-dimensional k -algebra A is basic then every simple R -module is 1-dimensional.

(1.5.33) (We will see shortly that) For every finite-dimensional k -algebra there is a basic algebra having an equivalent module category.

pr:eMsimple

(1.5.34) PROPOSITION. If M is a simple A -module; and $e^2 = e \in A$. Then eM is a simple eAe -module or zero. ■ (See e.g. §11.8.2.)

pr:eMJH

(1.5.35) PROPOSITION. Let $M \supset M_1 \supset \dots$ be a Jordan–Holder series for A -module M , with simple factors $L_i = M_i/M_{i+1}$; and $e^2 = e \in A$. Then $eM \supseteq eM_1 \supseteq \dots$ becomes a JH series for eM on deleting the terms for which $eM_i/eM_{i+1} = eL_i = 0$.

Thus in particular, if ${}_{eAe}eM$ is simple then the composition factors of M are a simple head factor appearing once, and any other factors L obey $eL = 0$. ■ (See e.g. (11.15).)

(1.5.36) Later we will provide detailed answers to the questions above. For now, our next step will be to construct some interesting algebras to play with, and hence some examples. We return to this discussion in (7.6.13) and §8.4.1 and §11.8.2.

1.5.5 Aside: tensor products

de:tensorprod0001

(1.5.37) Let R be a ring and $M = M_R$ and $N = {}_R N$ right and left R -modules respectively. Then there is a *tensor product* — an abelian group denoted $M \otimes_R N$ constructed as follows. Consider the formal additive group $\mathbb{Z}(M \times N)$, and the subgroup S_{MN} generated by elements of form $(m + m', n) - (m, n) - (m', n)$, $(m, n + n') - (m, n) - (m, n')$ and $(mr, n) - (m, rn)$ (all $r \in R$). We set $M \otimes_R N = \mathbb{Z}(M \times N)/S_{MN}$. (In essence $M \otimes_R N$ is equivalence classes of $M \times N$ under the relation $(mr, n) = (m, rn)$. See §7.4 for details.)

This construction is useful because it gives us, for each M_R , a functor $M_R \otimes -$ from R -mod to the category \mathbb{Z} -mod (of abelian groups). This has many useful generalisations.

1.5.6 Functor examples for module categories: globalisation

de:GF1

(1.5.38) Let A be an algebra over k and $e^2 = e \in A$ as in §1.5.4 above. We define functor $G = G_e$

$$G_e : eAe\text{-mod} \rightarrow A\text{-mod}$$


by $G_e M = Ae \otimes_{eAe} M$ (as defined in §7.4) and $F_e : A\text{-mod} \rightarrow eAe\text{-mod}$ by $F_e N = eN$. (Exercise: check that there are suitable mappings of module maps.)

ex:GF1

(1.5.39) EXERCISE. Show the following.

- (I) Pair (G_e, F_e) is an adjunction (as in (5.3.7)).
- (II) Functor F_e is exact.
- (III) Functor G_e is right exact, takes projectives to projectives and indecomposables to indecomposables. (See Th.7.5.19 et seq.)
- (IV) The composite $F_e \circ G_e : eAe\text{-mod} \rightarrow eAe\text{-mod}$ is a category isomorphism.

Note from these facts that there is an embedded image of $eAe\text{-mod}$ in $A\text{-mod}$ (the functorial version of an inclusion). Cf. Fig.1.1. Functor G_e does not take simples to simples in general. (One can see this either from the construction or ‘categorically’.) However since simples and indecomposable projectives are in bijective correspondence, we can effectively ‘count’ simples in

Figure 1.1: Schematic for the G -functor. fig:Pnmodembed1

A -mod by counting those in eAe -mod and then adding those which this count does not include. It is easy to see the following.

PROPOSITION. Functor F_e takes a simple module to a simple module or zero. ■

th:simp0001

(1.5.40) THEOREM. Let us write $\Lambda(A)$ for some index set for simple A -modules; and $\Lambda_e(A)$ for the subset on which e acts as zero. It follows from (1.5.39) that

$$\Lambda(A) = \Lambda(eAe) \sqcup \Lambda_e(A).$$

Of course simples on which e acts as zero are also the simples of the quotient algebra A/AeA , so $\Lambda_e(A) = \Lambda(A/AeA)$.

pr:lams

(1.5.41) PROPOSITION. Recall the partition algebra P_n from (1.4.2); and T_n from (1.4.6).

For $\delta \in k$ a unit, we may take $\Lambda(P_n) = \Lambda(P_{n-1}) \sqcup \Lambda(kS_n)$. Thus

$$\Lambda(P_n) = \sqcup_{i=0,1,\dots,n} \Lambda(kS_i).$$

Similarly $\Lambda(T_n) = \Lambda(T_{n-2}) \sqcup \Lambda(k)$. Thus

$$\Lambda(T_n) = \sqcup_{i=n,n-2,\dots,1/0} \Lambda(k).$$

Proof. (1.20) and (1.5.40). □

(1.5.42) Note that every simple module of P_n is associated to a symmetric group S_i irreducible for some $i \leq n$. Symmetric group irreducibles can be found in the heads of symmetric group Specht modules $\Delta_\lambda^S := kS_i v_\lambda$ (suitable $v_\lambda \in S_i$; these are classical constructions for irreducible modules over \mathbb{C} that are well defined over any ground ring). Accordingly we define P_n -module $\Delta(\lambda)$ by applying G -functors to Δ_λ^S as many times as necessary:

$$\Delta(\lambda) = G^{n-i} \Delta_\lambda^S$$

If $k \supset \mathbb{Q}$ then v_λ can be chosen idempotent (indeed primitive). It follows that $\Delta(\lambda)$ is indecomposable projective in a suitable quotient algebra of P_n . Thus it has simple head. It follows that every module's structure can be investigated by investigating morphisms from these modules.

(1.5.43) REMARK. The preceding example will be very useful for analysing P_n – mod by induction on n . But first we think about some other examples, and how module categories and functors work with representation theory in general.

1.6 Modular representation theory

ss:mod0001

Sometimes an algebra is defined over an arbitrary commutative ring k . We may focus on the representation theory over the cases of k a field in particular. But the idea of considering all cases together provides us with some useful tools (following ideas of Brauer [14]).

Let R be a commutative ring with a field of fractions (R_0) and quotient field k (quotient by some given maximal ideal). (Ring R a complete rank one discrete valuation ring would be sufficient to have such endowments.) Let A be an R -algebra that is a free R -module of finite rank. Let $A_0 = R_0 \otimes_R A$ and $A_k = k \otimes_R A$ (we call these constructions ‘base changes’ from R to R_0 and to k respectively).

The working assumption here is that A_0 is relatively easy to analyse. (The standard example would be a group algebra over a sufficiently large field of characteristic zero; which is semisimple by Maschke’s Theorem.) And that A_k is the primary object of study.

In particular, suppose that we have a complete set of simple modules for A_0 . One can see (e.g. in (??)) that:

LEMMA. For every A_0 -module M there is a finitely generated A -module (that is a free R -module) that passes to M by base change. ■

Note that there can be multiple non-isomorphic A -modules all passing to M . (We will give examples shortly.)

(1.6.1) Write

$$\mathcal{D} = \{D^R(l) : (l = 1, 2, \dots, m)\}$$

for a set of A -modules that passes by base change to a complete set of m simple A_0 -modules.

Write $D^k(l) = k \otimes D^R(l)$. Write L_λ^k ($\lambda \in \Lambda^k$) for a complete set of simple A_k -modules. Fixing k , this gives us a \mathcal{D} -decomposition matrix

$$D_{i\lambda} = [D^k(i) : L_\lambda^k]$$

(note that the index sets $1, 2, \dots, m$ and Λ^k are not the same).

Write P_λ^k for the projective cover of L_λ^k (the indecomposable projective with head L_λ^k), and e_λ^k for a corresponding primitive idempotent. One can show that there is a primitive idempotent in A that passes to e_λ^k , and an indecomposable projective A -module, $P_\lambda^{k,R}$ say, that passes to P_λ^k by base change (caveat: A is not Artinian in general).

(1.6.2) Since P_λ^k is projective, $D_{i\lambda} = \dim \text{hom}(P_\lambda^k, D^k(i))$. (Proof: For any indecomposable projective P_λ^k we have $\dim \text{hom}(P_\lambda^k, M) = [M : L_\lambda^k]$ by the exactness property (as in (1.5.9)) of the functor $\text{Hom}(P_\lambda^k, -)$. For example one can use exactness and an induction on the length of composition series.)

On the other hand the free R -module $\text{hom}(P_\lambda^{k,R}, D^R(i))$ has a basis which passes to a basis of $\text{hom}(P_\lambda^k, D^k(i))$; and to a basis of $\text{hom}(A_0 \otimes P_\lambda^{k,R}, A_0 \otimes D^R(i))$. A basis of the latter is the collection of maps, one for each simple factor of the direct sum $A_0 \otimes P_\lambda^{k,R}$ isomorphic to the simple module

$A_0 \otimes D^R(i)$. That is, the dimension is the multiplicity of the A_0 -simple module in $A_0 \otimes P_\lambda^{k,R}$. We have the following.

pr:mod recip

(1.6.3) PROPOSITION. (*Modular reciprocity*)

$$[D^k(i) : L_\lambda^k] = [A_0 \otimes P_\lambda^{k,R} : A_0 \otimes D^R(i)].$$

■

(1.6.4) For given k this says in particular that the Cartan decomposition matrix (with rows and columns indexed by Λ^k) is

$$C = D^T D \tag{1.24}$$

eq:Cartan0001

See e.g. §1.8.6.

Examples

1.7 Modules and ideals for the partition algebra P_n

1.7.1 Ideals

We continue to use the notations as in (1.4.4) and so on.

(1.7.1) Note that the number of propagating components cannot increase in the composition of partitions in P_n (the ‘bottleneck principle’). Hence $kP_{n,n}^m$ is an ideal of P_n for each $m \leq n$, and we have the following ideal filtration of P_n

$$P_n = kP_{n,n}^n \supset kP_{n,n}^{n-1} \supset \dots \supset kP_{n,n}^0.$$

Note that the sections $\mathfrak{P}_{n,n}^m := kP_{n,n}^m / kP_{n,n}^{m-1}$ of this filtration are bimodules, with bases $P_{n,m,n}$.

(1.7.2) Write

$$P_n^m := P_n / kP_{n,n}^m$$

for the quotient algebra.

(1.7.3) Note the natural inclusion

$$P_{n,l,m} \otimes v^* \hookrightarrow P_{n,l,m+1}$$

lem:natdecomp

(1.7.4) LEMMA. *For any $l \leq n$ there is a natural bijection*

$$P_{n,l,n} \xrightarrow{\sim} P_{n,l,l} \times P_{l,l,l} \times P_{l,l,n}$$

(the inverse map is essentially category composition).

1.7.2 Idempotents and idempotent ideals

(1.7.5) LEMMA. If $\delta \in k^*$ then $u_1 \in P_n$ is an unnormalised idempotent and

- (I) The ideal $kP_{n,n}^m = P_n(u^{\otimes(n-m)} \otimes 1_m)P_n$
- (II) $kP_{n,m} = kP_{n,m}^m \cong P_n(u^{\otimes(n-m)} \otimes 1_m)$ as a left P_n -module.

(1.7.6) Note that $kP_{n,l}^m$ is a left P_n -module (indeed a $P_n - P_l$ -bimodule) for each l, m , and $kP_{n,l}^{m-1} \subset kP_{n,l}^m$ (assuming $n \geq l \geq m$). Hence there is a quotient bimodule

$$\mathfrak{P}_{n,l}^l = kP_{n,l}^l / kP_{n,l}^{l-1}$$

with basis $P_{n,l,l}$.

There is a natural right action of the symmetric group S_l on this module (NB $S_l \subset P_l$), which we can use. Let $v_\lambda \in kS_l$ be such that $kS_l v_\lambda$ is a Specht S_l -module (an irreducible S_l -module over \mathbb{C}). Then define left P_n -module

$$D_\lambda = kP_{n,l,l} v_\lambda.$$

(1.7.7) If $k \supset \mathbb{Q}$ then v_λ can be chosen idempotent, and this module D_λ is a quotient of an indecomposable projective module, and hence has simple head. It follows that if P_n is semisimple then the modules of this form are a complete set of simple modules.

(1.7.8) EXERCISE. What can we say about $\text{End}_{P_n}(D_\lambda)$?

(1.7.9) EXERCISE. Construct some examples. What about contravariant duals?

(1.7.10) The case $n = 1$, $k = \mathbb{C}$. Fix δ . Artinian algebra P_1 has dimension 2. By (1.2.35) and (1.2.36) this tells us that either it is semisimple with two simple modules, or else it has one simple module.

Unless $\delta = 0$ then u/δ is idempotent so there are two simples. If $\delta = 0$ then u lies in the radical $J(P_1)$, and $P_1/J(P_1)$ is one-dimensional (semi)simple.

(1.7.11) The case $n = 2$, $k = \mathbb{C}$. Fix δ . Artinian algebra P_2 has dimension 15.

As we shall see, for most values of δ we have $P_2 \cong M_1(\mathbb{C}) \oplus M_1(\mathbb{C}) \oplus M_3(\mathbb{C}) \oplus M_2(\mathbb{C})$.

(1.7.12) We have $P_n \subset P_{n+1}$ via the injection given, say, by $p \mapsto p \cup \{\{n+1, (n+1)'\}\}$, which it will be convenient to regard as an inclusion.

1.8 Modules and ideals for T_n

Recall the definition (1.4.6) of T_n over k .

de:flippy

(1.8.1) Note that the flip map $t \mapsto t^*$ from (1.4.3) obeys $(t_1 t_2)^* = t_2^* t_1^*$. It follows that the flip \star defines an involutive antiautomorphism of T_n . Thus T_n is isomorphic to its opposite.

(1.8.2) Set $e_1^{2l-1} = e_1 e_3 \dots e_{2l-1}$ and (if $\delta \in k^*$) $\bar{e}_1^{2l-1} = \delta^{-l} e_1 e_3 \dots e_{2l-1}$. Then the ideal $T_n e_1 e_3 \dots e_{2l-1} T_n$ has basis $T_{n,n}^{n-2l}$ ($n - 2l$ or fewer propagating parts, as before). Write

$$T_n^{/n-2l} := T_n / (T_n e_1 e_3 \dots e_{2l-1} T_n)$$

for the quotient algebra by this ideal (with a basis of diagrams with more than $n - 2l$ propagating lines). In particular, (1.23) becomes $T_n^{/n-2} \cong k$.

Note that $\mathbf{e}_1 T_n^{/n-4} \mathbf{e}_1 \cong T_{n-2}^{/n-4} \cong k$ and $\mathbf{e}_1 \mathbf{e}_3 T_n^{/n-6} \mathbf{e}_1 \mathbf{e}_3 \cong T_{n-4}^{/n-6} \cong k$ and so on. By 1.5.29 this says that $\frac{1}{\delta} \mathbf{e}_1$ is a primitive idempotent in $T_n^{/n-4}$ and $\bar{\mathbf{e}}_1^3$ is primitive in $T_n^{/n-6}$ and so on:

(1.8.3) PROPOSITION. *The image of $\bar{\mathbf{e}}_1^{2l-1}$ is a primitive idempotent in the quotient algebra $T_n^{/n-2l-2}$.* \square

Hence the $T_n^{/n-4}$ -module $T_n^{/n-4} \mathbf{e}_1$ is indecomposable projective (we assume $\delta \in k^*$ for now); and hence also indecomposable with simple head as a T_n -module.

Generalising, define

$$D_n^{\text{tl}}(l) := T_n^{/n-2l-2} \mathbf{e}_1^{2l-1}$$

We have:

pr:DTL1 (1.8.4) PROPOSITION. *If $\delta \in k^*$, or $l \neq 0$, then $D_n^{\text{tl}}(l)$ is indecomposable with simple head as a T_n -module. Furthermore, by Prop.1.5.35 all the factors below the head obey $\mathbf{e}_1^{2l-1} L = 0$.* \blacksquare

pr:basisDTL (1.8.5) PROPOSITION. $\mathsf{T}_{n,l,l}$ is a basis for $D_n^{\text{tl}}(l)$. \blacksquare

Example: The case $\mathsf{T}_{4,2,2}$ is illustrated in Fig.1.2 (in the leftmost column).

(1.8.6) Note that there is a directly corresponding construction of indecomposable *right*-modules with analogous properties.

(1.8.7) There is also the construction of right-modules from the $D_n^{\text{tl}}(l)$ themselves by taking the ordinary duals, i.e. by applying the contravariant functor

$$()^* : T_n\text{-mod} \rightarrow \text{mod-}T_n$$

$$()^* : M \mapsto \text{Hom}_k(M, k)$$

(the duals $(D_n^{\text{tl}}(l))^*$ are also indecomposable on general grounds; but they need not have the other ‘standard’ properties from Prop.1.8.4 in general). We give a concrete example in (1.8.9).

One can ask how these two kinds of right modules are related. In general they are *not* isomorphic (but do have the same composition factors), as we shall see.

1.8.1 Some module morphisms

(1.8.8) It follows from (1.8.1) that every right T_n -module M can be made into a left T_n -module $\Pi_\star(M)$ by allowing T_n to act via the \star -map (the flip map). Note that a submodule of M passes to a submodule of $\Pi_\star(M)$. Indeed this map extends to a covariant functor between the categories of modules (in either direction):

$$\Pi_\star : \text{mod-}T_n \leftrightarrow T_n\text{-mod}$$

In particular, every exact sequence of right modules passes to an exact sequence of left modules.

Furthermore each module M has a contravariant (c-v) dual³, here denoted $\Pi^o(M)$:

$$\Pi^o(M) = \Pi_*(\text{Hom}_k(M, k)) = \Pi_*(M^*)$$

exa:422

(1.8.9) Example: What does the c-v dual of $M = D_n^{\text{tr}}(l)$ look like? As a k -module it is $\text{Hom}_k(M, k)$. Given a basis $\{b_1, b_2, \dots, b_r\}$ of M , the usual choice of basis of this ordinary dual is the set of linear maps f_i such that $f_i : b_j \mapsto \delta_{i,j}$. (The right-action of $a \in T_n$ is given by $(f_i a)(b_j) = f_i(ab_j)$. Thus $((f_i a)a')(b_j) = (f_i a)(a'b_j) = f_i(a'a'b_j)$ and $(f_i(aa'))(b_j) = f_i((aa')b_j) = f_i(a(a'b_j))$, so $((f_i a)a')(b_j) = (f_i(aa'))(b_j)$ as required.)

In our case let us order the basis of $M = D_n^{\text{tr}}(l)$ as in Fig.1.2. Then our basis for the dual is $\{f_1, f_2, \dots, f_{n-1}\}$.

Exercise: What is the right action of T_n on this k -module? For example, what is $f_1 U_1$?

de:headshot

(1.8.10) It follows from (1.8.4) that the only copy of the simple head L_l (say) of $D_n^{\text{tr}}(l)$ occurring in the c-v dual lies in the simple socle (note that e_1^{2l-1} is fixed under \star). It then follows from Schur's Lemma 1.2.22 that there is a unique T_n -module map, up to scalars, from $D_n^{\text{tr}}(l)$ to its contravariant dual — taking the simple head to the simple socle. (In theory the socle, which is the simple dual of the simple head, might not be isomorphic to it; allowing no map. But we will show the existence of at least one map explicitly.)

As we will see, it follows from this abstract representation theoretic argument that $D_n^{\text{tr}}(l)$ has a contravariant form (a bilinear form such that $\langle x, ay \rangle = \langle a^*x, y \rangle$, as in (??)) defined on it that is unique up to scalars.

Actually *finding* the explicit c-v form could be difficult in general. But in fact we can construct such a form here for all δ simultaneously (over a ring with δ indeterminate, as it were). We can use this to determine the structure of the module.

(1.8.11) For a, b in the basis $T_{n,l,l}$ (from (1.8.5)) then define $\alpha(a, b) \in k$ as follows. Note that $a^*b \in T_{l,l}$ (up to a scalar), thus either $a^*b = \alpha(a, b)c$ with $c \in T_{l,l,l}$ (indeed $c = 1_l$) for some $\alpha(a, b) \in k$; or $a^*b \in kT_{l,l}^{l-2}$, in which case set $\alpha(a, b) = 0$. Define an inner product on $kT_{n,l,l}$ by $\langle a, b \rangle = \alpha(a, b)$ and extending linearly.

ex:gramTL1

Example: Fig.1.2. The corresponding matrix of scalars is called the *gram matrix* with respect to this basis. From our example we have (in the handy alternative parameterisation $\delta = q + q^{-1}$):

$$\text{Gram}_n(n-2) = \begin{pmatrix} [2] & 1 & 0 & & \\ 1 & [2] & 1 & 0 & \\ 0 & 1 & [2] & 1 & \\ & & & \ddots & \\ 0 & \dots & 0 & 1 & [2] \end{pmatrix} \quad \text{so } |\text{Gram}_n(n-2)| = [n] = \frac{q^n - q^{-n}}{q - q^{-1}} \quad (1.25) \quad \text{eq:TLgram0001}$$

pr:innprodcov1


(1.8.12) PROPOSITION. The inner product defined by $\langle -, - \rangle$ is a contravariant form on $D_n^{\text{tr}}(l)$. ■

(1.8.13) Consider the k -space map

$$\phi_{\langle \rangle} : D_n^{\text{tr}}(l) \rightarrow \Pi^o(D_n^{\text{tr}}(l)) \quad (1.26)$$

$$\phi_{\langle \rangle} : m \mapsto \phi_{\langle \rangle}(m) \quad (1.27)$$

³The c-v dual of a module M over such a k -algebra is the ordinary dual right-module $\text{Hom}_k(M, k)$ made into a left-module via \star .

Figure 1.2: The array of diagrams $a \star b$ over the basis $\mathbf{T}_{4,2,2}$. fig:epud

where $\phi_{\chi}(m) \in \text{hom}(D_n^{\pi}(l), k)$ is given by

$$\phi_{\chi}(m)(m') = \langle m | m' \rangle .$$

(1.8.14) PROPOSITION. *The map ϕ_{χ} is a T_n -module homomorphism (unique up to scalars) from $D_n^{\pi}(l)$ to its contravariant dual.*

Proof. This map is a module morphism by Prop.1.8.12. To show uniqueness note that by (1.8.4) the contravariant dual must have the simple head of $D_n^{\pi}(l)$ as its simple socle (and only in the socle). Thus a head-to-socle map is the only possibility. \square

(1.8.15) EXAMPLE. In our example we have (from the gram matrix, using (1.8.9))

$$\begin{aligned} \phi_{\chi} : \cup || &\mapsto [2]f_1 + f_2 \\ \phi_{\chi} : | \cup | &\mapsto f_1 + [2]f_2 + f_3 \\ \phi_{\chi} : || \cup &\mapsto f_2 + [2]f_3 \end{aligned}$$

and for instance

$$\phi_{\chi} : \cup || - [2] | \cup | - [3] || \cup \mapsto [4]f_3$$

The point of this case is to show that the module map ϕ_{χ} has a kernel when $[4] = 0$. Obviously, in general,

PROPOSITION. If a T_n -module map has a kernel then the kernel is a submodule of the domain.

Thus in our case, when $[4] = 0$, the domain is not simple.

It will also be clear from the example that if the rank of the gram matrix is maximal then the morphism ϕ_{χ} has no kernel, and so is an isomorphism. This does not, of itself, show that the domain is a simple module, but we already showed in (1.8.10) that in our case the image must be simple, so the domain *is* simple.

(1.8.16) If $D_n^{\pi}(l)$ is in fact simple then ϕ_{χ} is an isomorphism and the contravariant form is non-degenerate. Otherwise the form is degenerate.

de:gramdetzero

It will be clear from our example that if the determinant of the gram matrix is non-zero then $D_n^{\text{tr}}(l)$ is simple; and otherwise it is not. (Note that the case $\delta = 0$ is excluded here, for brevity. It is easy to include it if desired, via a minor modification.) In particular if the determinant is zero then $D_n^{\text{tr}}(n-2)$ has composition length 2; and the other composition factor is the simple module $D_n^{\text{tr}}(n)$.

(1.8.17) PROPOSITION. *Given a c - v form (with respect to involutive antiautomorphism \star) on A -module M and $\text{Rad}_{<>} M = \{x \in M : \langle y, x \rangle = 0 \ \forall y\}$ then*

*(I) $\text{Rad}_{<>} M$ is a submodule, since $x \in \text{Rad}_{<>} M$ implies $\langle y, ax \rangle = \langle a^*y, x \rangle = 0$.*

(II) Thus $\dim \text{Rad}_{<>} M = \text{corank Gram}_{<>} M$. ■

(1.8.18) In our example rows 2 to $(n-1)$ of the $(n-1) \times (n-1)$ matrix $\text{Gram}_n(n-2)$ are clearly independent, while replacing $\boxed{\cup || \dots}$ (the basis element in the first row) by

$$w = \boxed{\cup || \dots} - [2] \boxed{|| \cup \dots} + [3] \boxed{|| \cup \dots} - \dots$$

(a sequence of elementary row operations adding to the first row multiples of each of the subsequent rows) replaces the first row of $\text{Gram}_n(n-2)$ with $(0, 0, \dots, 0, [n])$. That is, $\text{Rad}_{<>} D_n^{\text{tr}}(n-2) = 0$ unless $[n] = 0$. If $[n] = 0$ then w spans the Rad .

Explicit check in case $n = 4$: $U_1 w = ([2] - [2] + 0) \boxed{\cup ||} = 0$; $U_2 w = (1 - [2]^2 + [3]) \boxed{|| \cup} = 0$; $U_3 w = (0 - [2] + [2][3]) \boxed{|| \cup}$.

(1.8.19) It is easy to write down the form explicitly, particularly for $l = n-2$, and compute the determinant. We can use this to determine the structure of the algebra. First we will need a couple of functors.

(1.8.20) REMARK. In case M is a matrix over a PID, the *Smith form* of M (see e.g. [5]) is a certain diagonal matrix equivalent to M under elementary operations.

One sees from the proposition and example that the rank, or indeed a Smith form, of $\text{Gram} D$ is potentially more useful than the determinant. However note that working over $\mathbb{Z}[\delta]$ as we partly are, a Smith form may not exist until we pass specifically to \mathbb{C} , say (or at least to a PID $k[\delta]$ with k a field); and they are harder to compute when they do exist.

See §11.7 for more on this.

1.8.2 Aside on Res-functors (exactness etc)

ss:aside res

(1.8.21) Note the limits of what Res_ψ (from (1.5.10)) says in practice. For each B -module there is an A -module identical to it as a k -space. And for each B -module homomorphism there is an A -module homomorphism. It *does not* say that if $\text{Hom}_B(M, N) = 0$ then so is $\text{Hom}_A(M, N) = 0$.

In the particular case when ψ is surjective then M simple implies $\text{Res}_\psi M$ simple — i.e. M simple as an A -module (any A -submodule M' of M would also be a B -submodule, since in this case the B action is contained in the A action).

(1.8.22) We can also think about what happens to exact sequences under this functor Res_ψ . Suppose $M' \hookrightarrow M \twoheadrightarrow M''$ is a short-exact sequence of B -module maps. As we have just seen, it is again a sequence of A -module maps. The sequence is of the form $M' \hookrightarrow M \twoheadrightarrow M''$

since injection and surjection are properties of the underlying k -modules; but such a sequence is short-exact if $\dim(M') + \dim(M'') = \dim(M)$ — again a property of the underlying k -modules. In other words Res_ψ is an *exact* functor on finite dimensional modules.

We can also ask about split-ness. If the B -module sequence is split (i.e. $M = M' \oplus M''$) then there is another SES reversing the arrows, which again passes to an A -module sequence. If the B -module sequence is *non-split* what happens? Suppose that the A sequence is split. This means that there is an A -submodule of M isomorphic to M'' , i.e. (up to isomorphism) $aM'' \in M''$ for all a . Note that *if ψ is surjective*⁴ then every B action can be expressed as an A action (via ψ), so M'' is also a B -submodule, contradicting non-splitness. That is,

LEMMA. If ψ surjective then Res_ψ takes a non-split extension to a non-split extension. \square

1.8.3 Functor examples for module categories: induction

(1.8.23) Functor Res_ψ makes B a left- A right- B -bimodule; and there is a similar functor making B a left- B right- A -bimodule. Hence define

$$\text{Ind}_\psi : A\text{-mod} \rightarrow B\text{-mod}$$

by $\text{Ind}_\psi N = B \otimes_A N$ (cf. 1.5.38).

REMARK. This construction is typically used in case $\psi : A \rightarrow B$ is an inclusion of a subalgebra (in which case Res is called restriction).

(1.8.24) EXERCISE. Investigate these functors for possible adjunctions. Hints: Consider the map

$$a : \text{Hom}_B(\text{Ind}_\psi M, N) \rightarrow \text{Hom}_A(M, \text{Res}_\psi N)$$

given as follows. For $f \in \text{Hom}_B(\text{Ind}_\psi M, N)$ we define $a(f) \in \text{Hom}_A(M, \text{Res}_\psi N)$ by $a(f)(m) = f(1 \otimes m)$. Given $g \in \text{Hom}_A(M, \text{Res}_\psi N)$ we define $b(g) \in \text{Hom}_B(\text{Ind}_\psi M, N)$ by $b(g)(c \otimes m) = cg(m)$. One checks that $b = a^{-1}$, since $b(a(f)) = b(f(1 \otimes -)) = 1f = f$.

(1.8.25) EXAMPLE. We have in (1.21) above a surjective algebra map $\psi : P_n \rightarrow S_n$. It follows that every S_n -module is also a P_n -module via ψ . Of course every S_n -module map is also a P_n -module map.

(1.8.26) PROPOSITION. *The functor Ind_ψ takes projectives to projectives.* ■

1.8.4 Back to P_n

(1.8.27) Fix n . It follows from the results assembled above that for each $\lambda \vdash l \in \{n, n-1, \dots, 0\}$ we have a P_n -module $\Delta_\lambda = G^{n-l} S_\lambda$, where S_λ is a symmetric group Specht module. (Note that this notation omits n , so care is needed. We can write Δ_λ^n to emphasise n .)

Fix $k = \mathbb{C}$, so that every S_λ is simple. It follows that if P_n is semisimple for some given choice of δ (and some given n) then the set of Δ_λ modules is a complete set of simple modules for this algebra.

(1.8.28) More generally, if P_n is non-semisimple then at least one Δ_λ is not simple. Further, if Δ_λ^n is not simple, then Δ_λ^{n+1} is not simple. Thus, for fixed δ , we may think of the ‘first’ non-semisimple

⁴needed?




Figure 1.3: Orbits of an affine reflection group on \mathbb{Z} giving blocks for T_n with $l = 4$. fig:TLalcoves1

case (noting that P_0 is always simple), and hence a ‘first’ (one or more) non-simple Δ_λ — at level n say. We note that this first non-simple case is manifested by a homomorphism from some Δ_ν with $\nu \vdash n$.

There are a number of ways we can ‘detect’ these homomorphisms. One approach starts by noting another adjunction: the (ind,res) adjunction corresponding to the inclusion $P_n \hookrightarrow P_{n+1}$. One can work out $\text{Res}\Delta_\lambda$ by constructing an explicit basis for each Δ_λ . One can work out $\text{Ind}\Delta_\lambda$ by using the formula $\text{Ind} = \text{Res}G$. It then follows from the (ind,res) adjoint isomorphism that any such homomorphism implies a homomorphism to Δ_λ with $|\lambda|$ ‘close’ to n . These modules take a relatively simple form, and it is possible to detect morphisms to them explicitly by direct calculation.

Let D be the *decomposition matrix* for the Δ -modules (ordered in any way consistent with $\lambda > \mu$ if $|\lambda| > |\mu|$). It follows that D is upper unitriangular. It also follows that the *Cartan decomposition matrix* C is $C = DD^T$.

1.8.5 Back to T_n

To explain the P_n results it will be simpler to begin with T_n .

(1.8.29) Set $\Delta_l^T(l) = k$ (the trivial T_l -module) and

$$\Delta_n^T(l) = G_{\mathbf{e}_1} \Delta_{n-2}^T(l)$$

(1.8.30) EXAMPLE. $G_{\mathbf{e}_1} \Delta_{n-2}^T(n-2) = T_n \mathbf{e}_1 \otimes_{T_{n-2}} \Delta_{n-2}^T(n-2)$ (using the isomorphism to confuse $T_{n-2} \cong \mathbf{e}_1 T_n \mathbf{e}_1$). Noting $T_n \mathbf{e}_1 = k \mathbb{T}_{n,n-2} \otimes \cap$ (as in (1.4.3)); this is spanned by $\mathbb{T}_{n,n-2} \otimes_{T_{n-2}} 1_{n-2}$, where $\{1_{n-2}\}$ is acting as a basis for $\Delta_{n-2}^T(n-2)$. Note that $\mathbb{T}_{n,n-4,n-2} \otimes_{T_{n-2}} 1_{n-2} = 0$, so a basis is $\mathbb{T}_{n,n-2,n-2} \otimes_{T_{n-2}} 1_{n-2}$.

(1.8.31) LEMMA.

$$\Delta_n^T(l) \cong D_n^{\text{tr}}(l)$$

Proof. As above, a basis of $\Delta_n^T(l)$ is $\mathbb{T}_{n,l,l} \otimes_{T_l} 1_l$. Now cf. (1.8.5) and consider the obvious bijection between bases. The actions of $a \in T_n$ are the same — if (in the T category) $a * b \in k \mathbb{T}_{n,l,l}$ then $ab = a * b$ in both cases; otherwise $ab = 0$ in $\Delta_n^T(l)$ by the balanced map, and in $D_n^{\text{tr}}(l)$ by the quotient. \square

...See §?? for more details and treatment of the $\delta = 0$ case.

(1.8.32) THEOREM. [84, §7.3 Th.2] (Structure Theorem for T_n over \mathbb{C} .) Set $k = \mathbb{C}$. Fix $r \in \mathbb{N}$

(here we take $r \geq 3$ for now) and hence $q \in \mathbb{C}$ a primitive r -th root of unity. Fix $\delta = q + q^{-1}$. The simple content of $\Delta_n^T(\lambda)$ is given as follows.

Consider Fig.1.3. We give the positive real line two labellings for integral points: the natural labelling (with the origin labelled 0); and the *shifted* labelling. Points of form mr in the natural labelling ($mr - 1$ in the shifted labelling) are called *walls*. For given $\lambda \in \mathbb{N}_0$ determine m and b in \mathbb{N}_0 by $\lambda + 1 = mr + b$ with $0 \leq b < m$ (so b is the position of $\lambda + 1$ in the alcove above mr). For $b > 0$ set $\sigma_{m+1}.\lambda = \lambda + 2m - 2b$ — the image of λ under reflection in the wall above.

1) If $b = 0$ then $\Delta_n^T(\lambda) = L_n(\lambda)$.

2) Otherwise

$$0 \longrightarrow L_n(\lambda + 2m - 2b) \longrightarrow \Delta_n^T(\lambda) \longrightarrow L_n(\lambda) \longrightarrow 0$$

Here $L_n(\lambda + 2m - 2b)$ is to be understood as 0 if n is too small.

In particular the orbits of the reflection action describe the regular blocks (blocks of points not fixed by any non-trivial reflection); while the singular blocks (of points fixed by a non-trivial reflection) are singletons.

Proof. :

(1.8.33) By induction. We assume level $n = mr - 1$ and below. (And will work through a ‘cycle’ $n = mr, mr + 1, \dots, mr + r - 1$.) Thus, if $n' \equiv mr - 1 \pmod{2}$, we have $\Delta_{n'}^T(mr - 1) = L_{n'}(mr - 1) = P_{n'}(mr - 1)$.

Why? Firstly, We have ...

(1.8.34) LEMMA. (1) *Projective modules have filtrations by Δ modules; and the corresponding composition multiplicities are well defined.*

(2) *Once $n \geq \lambda$, so $P_n(\lambda)$ is defined, then the multiplicity $(P_n(\lambda) : \Delta_n^T(\lambda)) = 1$; $(P_n(\lambda) : \Delta_n^T(\mu)) = 0$ if $|\mu| \geq |\lambda|$ ($\mu \neq \lambda$); and otherwise $(P_n(\lambda) : \Delta_n^T(\mu))$ does not depend on n .*

Proof. (1) We can see this in various different ways. For now we note from §1.6 that both kinds of modules have lifts to the integral case, and hence corresponding modules in the ordinary case. But in the ordinary case the Δ -modules are simple.

(2) By (1.6.3) and (1.5.35). □

pr:resDeTL

(1.8.35) We have

$$0 \longrightarrow \Delta_{n-1}(l - 1) \longrightarrow \text{Res}\Delta_n(l) \longrightarrow \Delta_{n-1}(l + 1) \longrightarrow 0$$

pr:indresG

(1.8.36) PROPOSITION. *The functors Ind_ψ and $\text{Res}_\psi G$ are naturally isomorphic.*

(1.8.37) By 1.8.35 and 1.8.36 we have


$$\text{Ind } P(mr - 1) = \Delta^T(mr) + \Delta^T(mr - 2).$$

On the other hand

lem:Phwt1

(1.8.38) LEMMA. *Any projective is a direct sum of indecomposable projectives including those with the highest shifted label among those appearing in its Δ^T factors.*

(1.8.39) Hence $\text{Ind } P(mr - 1)$ contains $P(mr)$ as a direct summand.

Figure 1.4: fig:FRTL1

Suppose that $P(mr) = \Delta^T(mr)$. Then the remaining factor is also projective, so (again by 1.8.38) $P(mr-2) = \Delta^T(mr-2)$ — but this would imply $\Delta^T(mr-2) = L(mr-2)$ by reciprocity; and this contradicts the fact $\|\Delta_{ml}^T(ml-2)\| = 0$ from 1.8.11 (and an easy calculation).

Thus $\text{Ind } P(mr-1) = P(mr)$.

(1.8.40) Next we have

$$\text{Ind } P(mr) = \Delta^T(mr+1) + \Delta^T(mr-1) + \Delta^T(mr-1) + \Delta^T(mr-3)$$

Again this contains $P(mr+1)$ and the game is to determine which of the factors are in $P(mr+1)$.

If $\Delta^T(mr-1)$ is in $P(mr+1)$ then $L(mr+1)$ is in $\Delta^T(mr-1)$ by modular reciprocity (necessarily in the socle) which would imply $\|\Delta_{ml+1}^T(ml-1)\| = 0$ — a contradiction by (1.25).

de:TL901

(1.8.41) Next we will show by a contradiction that $P(mr+1) = \Delta^T(mr+1) + \Delta^T(mr-3)$. Suppose this sum splits. Then this would imply $P(mr-3) = \Delta^T(mr-3)$ and hence $L(mr-3) = \Delta^T(mr-3)$. However, consider the following.

de:TL902

(1.8.42) By Frobenius reciprocity we have

$$\text{Hom}(\text{Ind } A, B) \cong \text{Hom}(A, \text{Res } B)$$


in particular in the case in Fig.1.4: ⁵

$$\text{Hom}(\text{Ind } \Delta_{ml}^T(ml), \Delta_{ml+1}^T(ml-3)) \cong \text{Hom}(\Delta_{ml}^T(ml), \text{Res } \Delta_{ml+1}^T(ml-3))$$

Note that $\text{Res } \Delta_{ml+1}^T(ml-3) = \Delta_{ml}^T(ml-2) \oplus \Delta_{ml}^T(ml-4)$ (a direct sum by the block assumption), so that the RHS is nonzero by assumption, noting $??$. Thus the LHS is nonzero. There is no map from $\Delta^T(ml+1)$ to $\Delta^T(ml-1)$, so there is a map from $\Delta^T(ml+1)$ to $\Delta^T(ml-3)$. This demonstrates the contradiction needed in 1.8.41. Thus

$$P(mr+1) = \Delta^T(mr+1) + \Delta^T(mr-3)$$

⁵caveat: $l = r!!!$

Figure 1.5: fig:FRTL3

(1.8.43) Next we have

$$\text{Ind } P(mr + 1) = \Delta^T(mr + 2) + \Delta^T(mr) + \Delta^T(mr - 2) + \Delta^T(mr - 4)$$

We have $P(mr + 2) = \Delta^T(mr + 2) + \dots$. The question is, which of the factors above should be included? If we include $\Delta^T(mr)$ then $L(mr + 2)$ is in $\Delta^T(mr)$ by modular reciprocity. We can eliminate this possibility in a couple of ways. For example, we can compute a central element of T_n and show using this that the two shifted labels are in different blocks. Alternatively we can compute $||\Delta_{mr+2}^T(mr)||$ and check that it is nonzero in this case.

So far, then, we have that $\text{Ind } P(mr + 1) = P(mr + 2) \oplus P(mr) \oplus \dots$. However since $P(mr) = \Delta^T(mr) + \Delta^T(mr - 2)$ we have $P(mr + 2) = \Delta^T(mr + 2) + X$ where $X = \Delta^T(mr - 4)$ or zero.

In the latter case we would have $P(mr - 4) = \Delta^T(mr - 4)$. This contradicts the inductive assumption for every m value except $m = 1$. For $m = 1$ (or in general) we note instead that

$$\text{Hom}(\text{Ind } \Delta_{mr+1}^T(mr + 1), \Delta_{mr+2}^T(mr - 4)) \cong \text{Hom}(\Delta_{mr+1}^T(mr + 1), \text{Res } \Delta_{mr+2}^T(mr - 4))$$

and that the RHS is nonzero (for $r > 3$) by the inductive assumption (indeed we just showed this in 1.8.42 above). Thus the LHS is nonzero. But there is no map $\Delta^T(mr) \rightarrow \Delta^T(mr - 4)$ by the inductive assumption, so there is a map $\Delta^T(mr + 2) \rightarrow \Delta^T(mr - 4)$. This provides the required contradiction. That is

$$P(mr + 2) = \Delta^T(mr + 2) + \Delta^T(mr - 4)$$

(1.8.44) We may continue in the same way until $P(mr + (r - 2))$. At this point $\text{Res } \Delta^T(mr - r - 1)$ is not a direct sum (indeed it is indecomposable projective) and the argument for a nonzero RHS in Frobenius reciprocity fails. This tells us that there is no map on the LHS, so $P(mr + (r - 2)) = \Delta^T(mr + (r - 2))$ and we have completed the main inductive step. \square

1.8.6 The decomposition matrices

ss:decompmatex1

Note that the decomposition matrices (from §1.6 and (1.24)) are determined by the structure Theorem. The matrix for a single block (starting from the low-numbered weight) is of form

$$D_{block} = \begin{pmatrix} 1 & 1 & & & \\ & 1 & 1 & & \\ & & 1 & 1 & \\ & & & \ddots & \\ & & & & 1 & 1 \\ & & & & & 1 \end{pmatrix}$$

(this should be thought of as the n -dependent truncation of a semiinfinite matrix continuing down to the right), that is $\Delta^T(0)$ (say, from the first row) contains $L(0)$ and the next simple in the block, and so on; giving

$$C_{block} = D^T D = \begin{pmatrix} 1 & 1 & & & \\ 1 & 2 & 1 & & \\ & 1 & 2 & 1 & \\ & & & \ddots & \\ & & & & 1 & 2 & 1 \\ & & & & & 1 & 2 \end{pmatrix}$$

...

1.8.7 Odds and ends

(1.8.45) By 1.5.35 and 1.6.3 the $\Delta_n(l)$ content of $P_n(m)$ does not depend on n (once n is big enough for these modules to make sense). Thus $P_n(0) = \Delta_n(0)$; $P_n(1) = \Delta_n(1)$.

For $P_n(2)$ we have $\text{Ind } P_n(1) = \Delta_n(0) + \Delta_n(2)$; and $\text{Ind } P_n(1)$ contains $P_n(2)$ as a direct summand. If this is a proper direct sum then this is true in particular at $n = 2$ and there is a primitive idempotent decomposition of 1 in T_2 . It is easy to see that this depends on δ , but it true unless $\delta = 0$. (*We shall assume for now that $k = \mathbb{C}$ for definiteness.*)

Another way to look at the decomposition of $\text{Ind } P_n(1)$ is as follows. If it does not decompose then by ?? there is a homomorphism $\Delta(2) \rightarrow \Delta(0)$, so that the gram matrix of $\Delta(0)$ must be singular.

Let us assume $\delta \neq 0$. Proceeding to $P_n(3)$ we have $\text{Ind } P_n(2) = \Delta_n(1) + \Delta_n(3)$. Again this splits if and only if the gram matrix for $\Delta(1)$ is singular.

(1.8.46) TO DO:

Grothendieck group

1.9 Lie algebras

ss:Liealg0

We include a brief discussion of Lie algebras here,

(a) to provide some contrast with and hence context for our ‘associative’ algebras; and

(b) as a certain partner notion to the special case of (associative) finite group algebras. See 19.3.1 for a more detailed exposition. Here k is a field.

(1.9.1) A Lie algebra A over field k is a k -vector space and a bilinear operation $A \times A \rightarrow A$ denoted $[a, b]$ such that $[a, a] = 0$ and

$$[a, [b, c]] + [b, [c, a]] + [c, [a, b]] = 0 \quad (\text{'Jacobi identity'})$$

(1.9.2) From an associative algebra T we obtain a Lie algebra $Lie(T)$ by $[a, b] = ab - ba$.

(1.9.3) In particular, for V a vector space, the space of endomorphisms, sometimes denoted $gl(V)$, is a Lie algebra with $[a, b] = ab - ba$ (where ab is the composition of endomorphisms).

A representation of a Lie algebra A over k is a Lie algebra morphism $\rho : A \rightarrow gl(V)$ for some V .

An A -module is a space V and a map $A \times V \rightarrow V$ with

$$[a, b]v = a(bv) - b(av).$$

(1.9.4) Let V, V' be A -modules. Then the tensor product $V \otimes_k V'$ has a 'diagonal' action of A :

$$a(v \otimes v') = av \otimes v' + v \otimes av'$$

that makes $V \otimes_k V'$ an A -module.

Check: $[a, b](v \otimes v') = [a, b]v \otimes v' + v \otimes [a, b]v' = (a(bv) - b(av)) \otimes v' + v \otimes (a(bv') - b(av')) = \dots$

(1.9.5) The tensor algebra of Lie algebra A is the vector space

$$\tau = \bigoplus_{n \geq 0} A^{\otimes n}$$

with multiplication given by $(a \otimes b)(c \otimes d) = a \otimes b \otimes c \otimes d$ and so on. Set H to be the ideal in τ generated by the elements of form $a \otimes b - b \otimes a - [a, b]$, with $a, b \in A$. Define

$$U_A = \tau/H$$

(1.9.6) A *universal enveloping algebra* (UEA) of Lie algebra A is an associative algebra U together with a Lie algebra homomorphism $I : A \rightarrow Lie(U)$ such that every Lie algebra homomorphism of form $h : A \rightarrow Lie(B)$ has a unique 'factorisation through $Lie(U)$ ', that is, a unique morphism of associative (unital) algebras $f : U \rightarrow B$ such that $h = f \circ I$.

(1.9.7) U_A is a UEA for A , with the homomorphism $I : A \rightarrow Lie(U_A)$ given by $a \mapsto a + H$. It is unique as such up to isomorphism.

(1.9.8) There is a vector space bijection

$$Hom_{Lie}(A, Lie(B)) \cong Hom(U, B).$$

(1.9.9) Let V be an A -module and $\rho : A \rightarrow gl(V)$ the corresponding representation. Then ρ extends to a representation of a UEA U . This lifts to an 'isomorphism' of the categories of A -modules and U -modules (as subcategories of the category of vector spaces).

(1.9.10) THEOREM. (*Poincare–Birkoff–Witt*) Let $J = \{j_1, j_2, \dots\}$ be an ordered basis of A . Then the monomials of form $I(j_{i_1})I(j_{i_2})\dots I(j_{i_n})$ with $i_1 \leq i_2 \leq \dots$ and $n \geq 0$ are a basis for U_A .

(1.9.11) Recalling that k is fixed here, write sl_n for the Lie algebra of traceless $n \times n$ matrices. For example, sl_2 has k -basis:

$$x^+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad x^- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

These obey $[x^+, x^-] = h$, $[h, x^+] = 2x^+$, $[h, x^-] = -2x^-$.

1.10 Eigenvalue problems

(1.10.1) Operators acting on a space; their eigenvectors and eigenvalues.

Here we remark very briefly and generally on the kind of Physical problem that can lead us into representation theory.

A typical Physical problem has a linear operator Ω acting on a space H , with that action given by the action of the operator on a (spanning) subset of the space. One wants to find the eigenvalues of Ω .

The eigenvalue problem may be thought of as the problem of finding the one-dimensional subspaces of H as an $\langle \Omega \rangle$ -module, where $\langle \Omega \rangle$ is the (complex) algebra generated by Ω . That is, we want to find elements h_i in H such that:

$$\Omega h_i = \lambda_i h_i$$

— noting only that, usually, the object of primary physical interest is λ_i rather than h_i . If H is finite dimensional then (the complex algebra generated by) Ω will obey a relation of the form

$$\prod_i (\Omega - \lambda_i)^{m_i} = 0$$

Of course the details of this form are *ab initio* unknown to us. But, proceeding formally for a moment, if any $m_i > 1$ (necessarily) here, so that $S = \prod_i (\Omega - \lambda_i) \neq 0$, then S generates a non-vanishing nilpotent ideal (we say, the algebra has a radical). Obviously any such nilpotent object has 0-spectrum, so two operators differing by such an object have the same spectrum. In other words, the image of Ω in the quotient algebra by the radical has the same spectrum $\{\lambda_i\}$. An algebra with vanishing radical (such as the quotient of a complex algebra by its radical) has a particularly simple structural form, so this is a potentially useful step.

However, gaining *access* to this form may require enormously greater arithmetic complexity than the original algebra. In practice, a balance of techniques is most effective, even when motivated by physical ends. This balance can often be made by analysing the regular module (in which every eigenvalue is manifested), and thus subquotients of projective modules, but not more exotic modules. (Of course Mathematically other modules may well also be interesting — but this is a matter of aesthetic judgement rather than application.)

It may also be necessary to find the subspaces of H as a module for an algebra generated by a set of operators $\langle \Omega_i \rangle$. A similar analysis pertains.

A particularly nice (and Physically manifested) situation is one in which the operators Ω_i (whose unknown spectrum we seek to determine) are known to take the form of the representation matrices of elements of an abstract algebra A in some representation:

$$\Omega_i = \rho(\omega_i)$$

Of course any reduction of Ω_i in the form of (1.10) reduces the problem to finding the spectrum of $R_1(\omega_i)$ and $R_2(\omega_i)$. Thus the reduction of ρ to a (not necessarily direct) sum of irreducibles:

$$\rho(\omega_i) \cong \bigoplus_{\alpha} \rho_{\alpha}(\omega_i)$$

reduces the spectrum problem in kind. In this way, Physics drives us to study the representation theory of the abstract algebra A .

1.11 Notes and references

ss:refs

The following texts are recommended reading: Jacobson[61, 62], Bass[6], MacLane and Birkhoff[79], Green[52], Curtis and Reiner[30, 32], Cohn[24], Anderson and Fuller[3], Benson[7], Adamson[2], Cassels[20], Magnus, Karrass and Solitar[80], Lang[75], and references therein. .

1.12 Exercises

exe:gr01

(1.12.1) Let R be a commutative ring and S a set. Then RS denotes the ‘free R -module with basis S ’, the R -module of formal finite sums $\sum_i r_i s_i$ with the obvious addition and R action. Show that this is indeed an R -module.

exe:gr1

(1.12.2) Let R be a commutative ring and G a finite group. Show that the multiplication in (1.12) makes RG a ring.

Hints: We need to show associativity. We have

$$\left(\left(\sum_i r_i g_i \right) \left(\sum_j r'_j g_j \right) \right) \left(\sum_k r''_k g_k \right) = \left(\sum_{ij} (r_i r'_j) (g_i g_j) \right) \left(\sum_k r''_k g_k \right) = \sum_{ijk} ((r_i r'_j) r''_k) ((g_i g_j) g_k)$$

(1.28) **groupalgmult2**

and

$$\left(\sum_i r_i g_i \right) \left(\left(\sum_j r'_j g_j \right) \left(\sum_k r''_k g_k \right) \right) = \left(\sum_i r_i g_i \right) \left(\sum_{jk} (r'_j r''_k) (g_j g_k) \right) = \sum_{ijk} (r_i (r'_j r''_k)) (g_i (g_j g_k))$$

(1.29) **groupalgmult3**

These are equal by associativity of multiplication in R and G separately.

(1.12.3) Show that RG is still a ring as above if G is a not-necessarily finite monoid and RG means the free module of finite support as above.

Hints: Multiplication in monoid G is also associative.

1.12.1 Radicals

ss:radical0001

Write J_R for the radical of ring R .

(1.12.4) A ring is *semiprime* if it has no nilpotent ideal.

(1.12.5) THEOREM. A ring is left-semisimple if and only if every left ideal is a direct summand of the left regular module. ■

Show:

(1.12.6) THEOREM. If S a subring of ring R such that, regarded as an S -bimodule, R contains S as a direct summand, then R left-semisimple implies S is left-semisimple.

Hint:

Let S' be an S -bimodule complement of S in R : that is, $R = S \oplus S'$ as an S -bimodule. (For example if $R = \mathbb{C}$ and $S = \mathbb{R}$ then we can take $S' = \mathbb{R}z$ for any $z \in \mathbb{C} \setminus \mathbb{R}$.) If I is any left ideal of S then it is in particular a subset of R and RI makes sense as a left R -module, and hence as a left S -module by restriction. We claim $RI = (S \oplus S')I = SI \oplus S'I = S \oplus S'I$ as a left S -module. Now RI is a direct summand of ${}_R R$ by left-semisimplicity, so I is a direct summand of ${}_S S$.

(1.12.7) Let G be a finite group of automorphisms of ring R . Write r^g for the image of $r \in R$ under $g \in G$. Show that

$$R^G := \{r \in R \mid r^g = r \ \forall g \in G\}$$

is a subring of R .

Show:

(1.12.8) THEOREM. Suppose that $|G|$ is invertible in R . If R is semisimple Artinian (e.g. a semisimple algebra over a field) then R^G is semisimple Artinian. ■

Hints:

Show that $J_R \cap R^G \subseteq J_{R^G}$.

1.12.2 What is categorical?

ss:whatcat

(1.12.9) Prove: THEOREM. Let A be an Artinian algebra and I an ideal. Then A/I non-semisimple implies A non-semisimple. ■

Solution: (There are many ways to prove this. Here is one close to the idea of indecomposable matrix representations.) If A/I non-semisimple then not every module is a direct sum of simple modules (by definition), so there are a pair of modules with a non-split extension between them. That is, there is a short exact sequence

$$0 \longrightarrow M' \xrightarrow{i} M \xrightarrow{p} M'' \longrightarrow 0$$

such that there is no sequence with the arrows reversed. This sequence, indeed any sequence involving these modules, is also ‘in’ A -mod via $\psi : A \rightarrow A/I$. Now suppose (for a contradiction) that there is a sequence in A -mod involving the images of these modules but with the arrows reversed. This means that some $N \subset M$ obeys $N \cong M''$ as an A -submodule of M , i.e. $AN = N$ (keep in mind that the action of A on M and hence N comes by $am = \psi(a)m$, and the A/I -module property of M). But ψ is surjective, so every $x \in A/I$ is $\psi(a)$ for some a , so $(A/I)N = AN = N$

so N is also an A/I -submodule. This is a contradiction. Thus the original sequence is non-split in $A\text{-mod}$. \square

(1.12.10) Write $\text{Res}_\psi : A/I\text{-mod} \rightarrow A\text{-mod}$ for the functor associated to $\psi : A \rightarrow A/I$.

Let B be any algebra. Note that given a sequence of B -module maps

$$L \xrightarrow{f} M \xrightarrow{g} N$$

there is, trivially, an underlying sequence of maps of these objects as abelian groups. The exactness property at M , $\text{im}(f) = \ker(g)$, is defined at the level of abelian groups. Thus the sequence is exact for any B if and only if it is exact at the level of abelian groups.

Use this to show that Res_ψ is exact.