


Logistic Regression




sigmoid function

logistic function

outputs between 0 and 1

$$g(z) = \frac{1}{1 + e^{-z}} \quad 0 < g(z) < 1$$



$$f_{\vec{w}, b}(\vec{x}) = g(\vec{w} \cdot \vec{x} + b) = \frac{1}{1 + e^{-(\vec{w} \cdot \vec{x} + b)}}$$

logistic Regression

Example

x is "tumor size"

y is 0 (not malignant)

or 1 (malignant)

$$f_{\vec{w}, b}(\vec{x}) = 0.7$$


70% chance that y is 1

$$f_{\vec{w}, b}(\vec{x}) = P(y = 1 | \vec{x}; \vec{w}, b)$$

Probability that y is 1,

given input \vec{x} , parameters \vec{w}, b

Decision boundary



$$f_{\vec{w}, b}(\vec{x}) = g(z) = g(\underbrace{w_1 x_1}_{1} + \underbrace{w_2 x_2}_{1} + \underbrace{b}_{-3})$$

Decision boundary

$$z = \vec{w} \cdot \vec{x} + b = 0$$

$$z = x_1 + x_2 - 3 = 0$$

$$x_1 + x_2 = 3$$

Non-linear decision boundaries

Complex -- need Higher-order polynomial terms allow for more intricate decision boundaries

like: $f_{\vec{w}, b}(\vec{x}) = g(z) = g(w_1 x_1 + w_2 x_2 + w_3 x_1^2 + w_4 x_1 x_2 + w_5 x_2^2 + w_6 x_1^3 + \dots + b)$

Cost function for logistic regression

$$J(\vec{w}, b) = \frac{1}{m} \sum_{i=1}^m \underbrace{L(f_{\vec{w}, b}(\vec{x}^{(i)}), y^{(i)})}_{\text{loss}}$$

New definition

$$L(f_{\vec{w}, b}(\vec{x}^{(i)}), y^{(i)}) = \begin{cases} -\log(f_{\vec{w}, b}(\vec{x}^{(i)})) & \text{if } y^{(i)} = 1 \\ -\log(1 - f_{\vec{w}, b}(\vec{x}^{(i)})) & \text{if } y^{(i)} = 0 \end{cases}$$

convex
↳ can reach a
global minimum

Simplified loss function

$$L(f_{\vec{w}, b}(\vec{x}^{(i)}), y^{(i)}) = \underbrace{-y^{(i)} \log(f_{\vec{w}, b}(\vec{x}^{(i)})) - (1 - y^{(i)}) \log(1 - f_{\vec{w}, b}(\vec{x}^{(i)}))}_{\text{convex (single global minimum)}}$$

$$J(\vec{w}, b) = \frac{1}{m} \sum_{i=1}^m [L(f_{\vec{w}, b}(\vec{x}^{(i)}), y^{(i)})]$$

$$= \frac{1}{m} \sum_{i=1}^m [y^{(i)} \log(f_{\vec{w}, b}(\vec{x}^{(i)})) - (1 - y^{(i)}) \log(1 - f_{\vec{w}, b}(\vec{x}^{(i)}))]$$

Gradient Descent Implementation in Logistic Regression

cost

$$J(\vec{w}, b) = -\frac{1}{m} \sum_{i=1}^m \left[y^{(i)} \log(f_{\vec{w}, b}(\vec{x}^{(i)})) - (1 - y^{(i)}) \log(1 - f_{\vec{w}, b}(\vec{x}^{(i)})) \right]$$

repeat {
j = 1 ... n

$$w_j = w_j - \alpha \frac{\partial}{\partial w_j} J(\vec{w}, b)$$

$$b = b - \alpha \frac{\partial}{\partial b} J(\vec{w}, b)$$

} simultaneous updates

$$\frac{\partial}{\partial w_j} J(\vec{w}, b) = \frac{1}{m} \sum_{i=1}^m (f_{\vec{w}, b}(\vec{x}^{(i)}) - y^{(i)}) x_j^{(i)}$$

$$\frac{\partial}{\partial b} J(\vec{w}, b) = \frac{1}{m} \sum_{i=1}^m (f_{\vec{w}, b}(\vec{x}^{(i)}) - y^{(i)})$$

repeat {
j = 1 ... n

$$w_j = w_j - \alpha \left[\frac{1}{m} \sum_{i=1}^m (f_{\vec{w}, b}(\vec{x}^{(i)}) - y^{(i)}) x_j^{(i)} \right]$$

$$b = b - \alpha \left[\frac{1}{m} \sum_{i=1}^m (f_{\vec{w}, b}(\vec{x}^{(i)}) - y^{(i)}) \right]$$

} simultaneous updates

Same concepts:

- Monitor gradient descent (learning curve)
- Vectorized implementation
- Feature scaling

Linear Regression $f_{\vec{w}, b}(\vec{x}) = \vec{w} \cdot \vec{x} + b$

Logistic Regression $f_{\vec{w}, b}(\vec{x}) = \frac{1}{1 + e^{-(\vec{w} \cdot \vec{x} + b)}}$
Sigmoid function