

# QUANTUM COMPUTATION AND QUANTUM INFORMATION: THE QUANTUM FOURIER TRANSFORM

## 1

We consider the linear map in  $\mathbb{C}^N$  which acts on the computational basis as

$$|j\rangle \mapsto \frac{1}{\sqrt{N}} \sum_{k=0}^{N-1} e^{\frac{2i\pi jk}{N}} |k\rangle$$

Let  $A$  be the matrix of the transformation in the computational basis.

$$\forall (k, l) \in \llbracket 0, N-1 \rrbracket^2, \quad a_{kl} = \frac{1}{\sqrt{N}} e^{\frac{2i\pi kl}{N}}$$

The adjoint matrix  $A^\dagger$  is then

$$\begin{aligned} \forall (k, l) \in \llbracket 0, N-1 \rrbracket^2, \quad b_{kl} &= a_{lk}^* \\ &= \frac{1}{\sqrt{N}} e^{-\frac{2i\pi kl}{N}} \end{aligned}$$

We compute the coefficient  $k, l$  of the product  $AA^\dagger$ :

$$\begin{aligned} \forall (k, l) \in \llbracket 0, N-1 \rrbracket^2, \quad c_{kl} &= \sum_{j=0}^{N-1} a_{kj} b_{jl} \\ &= \frac{1}{N} \sum_{j=0}^{N-1} e^{\frac{2i\pi j}{N} (k-l)} \\ &= \frac{1}{N} \sum_{j=0}^{N-1} (e^{\frac{2i\pi}{N} (k-l)})^j \\ &= \begin{cases} \frac{1}{N} \frac{1 - (e^{\frac{2i\pi}{N} (k-l)})^N}{1 - e^{\frac{2i\pi}{N} (k-l)}} = 0 & \text{if } e^{\frac{2i\pi}{N} (k-l)} \neq 1, \\ 1 & \text{if } e^{\frac{2i\pi}{N} (k-l)} = 1. \end{cases} \\ &= \begin{cases} 0 & \text{if } k \neq l, \\ 1 & \text{if } k = l. \end{cases} \\ &= \delta_{kl} \end{aligned}$$

which shows that  $AA^\dagger = A^\dagger A = I$  i.e.  $A$  is unitary.

## 2

Here the dimension of the state space is  $N = 2^n$ . The Fourier transform of the  $n$  qubit state  $|00 \dots 0\rangle$  is

$$A|0\rangle = \frac{1}{\sqrt{N}} \sum_{k=0}^{N-1} |k\rangle$$

we can write  $k$  in binary  $k_{n-1} \dots k_1 k_0$

$$A|0\rangle = \frac{1}{2^{n/2}} \sum_{k_0, k_1, \dots, k_{n-1}=0}^1 |k_{n-1} \dots k_1 k_0\rangle$$

or in product representation,

$$= \frac{1}{2^{n/2}} \underbrace{(|0\rangle + |1\rangle)(|0\rangle + |1\rangle) \dots (|0\rangle + |1\rangle)}_{n \text{ qubits}}$$

## 3

Let  $N = 2^n$  and  $Y = (y_k)_{k \in \llbracket 0, N-1 \rrbracket}$  be the classical fourier transform of  $X = (x_k)_{k \in \llbracket 0, N-1 \rrbracket}$ .

$$\forall k \in \llbracket 0, N-1 \rrbracket, \quad y_k = \sum_{j=0}^{N-1} e^{\frac{2i\pi k j}{2^n}} x_j$$

The factor  $\frac{1}{\sqrt{N}}$  is omitted for clarity. We can write  $j$  in binary  $j_{n-1} \dots j_1 j_0$

$$\begin{aligned} y_k &= \sum_{j_0, j_1, \dots, j_{n-1}=0}^1 e^{\frac{2i\pi k (2^{n-1}j_{n-1} + \dots + 2j_1 + j_0)}{2^n}} x_j \\ &= \sum_{j_1, \dots, j_{n-1}=0}^1 e^{\frac{2i\pi k (2^{n-1}j_{n-1} + \dots + 2j_1)}{2^n}} x_{j_{n-1} \dots j_1 0} + \sum_{j_1, \dots, j_{n-1}=0}^1 e^{\frac{2i\pi k (2^{n-1}j_{n-1} + \dots + 2j_1 + 1)}{2^n}} x_{j_{n-1} \dots j_1 1} \\ &= \sum_{j_1, \dots, j_{n-1}=0}^1 e^{\frac{2i\pi k (2^{n-1}j_{n-1} + \dots + 2j_1)}{2^n}} x_{j_{n-1} \dots j_1 0} + e^{\frac{2i\pi k}{2^n}} \sum_{j_1, \dots, j_{n-1}=0}^1 e^{\frac{2i\pi k (2^{n-1}j_{n-1} + \dots + 2j_1)}{2^n}} x_{j_{n-1} \dots j_1 1} \\ &= \sum_{j_1, \dots, j_{n-1}=0}^1 e^{\frac{2i\pi k (2^{n-2}j_{n-1} + \dots + j_1)}{2^{n-1}}} x_{j_{n-1} \dots j_1 0} + e^{\frac{2i\pi k}{2^n}} \sum_{j_1, \dots, j_{n-1}=0}^1 e^{\frac{2i\pi k (2^{n-2}j_{n-1} + \dots + j_1)}{2^{n-1}}} x_{j_{n-1} \dots j_1 1} \end{aligned}$$

We see the first sum is the  $k^{th}$  coefficient of the FT of the sequence  $(x_{2k})_{k \in \llbracket 0, N/2-1 \rrbracket}$  and the second is the  $k^{th}$  coefficient of the FT of  $(x_{2k+1})_{k \in \llbracket 0, N/2-1 \rrbracket}$ . This shows that to compute FT of sequence of length  $N$ , we have to compute 2 FT of sequence of length  $\frac{N}{2}$  and do  $2N$  complex additions/multiplications. The complexity of the operation  $T(N)$  follows the recurrence:

$$T(N) = 2T\left(\frac{N}{2}\right) + 2N$$

We can use the Master theorem<sup>1</sup>:

**Theorem.** Let  $a \geq 1$  and  $b > 1$  be constants, let  $f(n)$  be a function, and let  $T(n)$  be defined on the non negative integers by the recurrence

$$T(n) = aT\left(\frac{n}{b}\right) + f(n)$$

where we interpret  $\frac{n}{b}$  to mean either  $\lfloor \frac{n}{b} \rfloor$  or  $\lceil \frac{n}{b} \rceil$ . Then  $T(n)$  has the following asymptotic bounds:

- (1) If  $f(n) = O(n^{\log_b a - \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$ .
- (2) If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \log n)$ .
- (3) If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(\frac{n}{b}) \leq cf(n)$  for some constant  $c < 1$  and  $n$  sufficiently large, then  $T(n) = \Theta(f(n))$ .

Here we are in the second case of the theorem, so  $T(N) = \Theta(N \log(N)) = \Theta(n 2^n)$ .

---

<sup>1</sup>Thomas H. Cormen and Charles E. Leiserson : *Introduction to algorithms*, MIT Press (2009)