everything we discussed.
I'll give you the baby's timeline

1. Molecular Induction and Neural Tube Formation (Weeks 3–4)

Category: Embryonic Signaling and Neurulation

• Primary Embryonic Induction: Around day 16, the notochord (a primitive structural rod in the embryo) releases specific signaling proteins, primarily Noggin, Chordin, and Follistatin. These proteins block Bone Morphogenetic Proteins (BMPs), which would normally turn the tissue into skin. By blocking BMPs, these signals force the overlying ectoderm to transform into the neural plate.
• Cellular Elongation and Constriction: The cells of the neural plate change shape via microtubule elongation. Microfilaments at the top of these cells contract (apical constriction), pinching the cells like a drawstring. This forces the flat neural plate to bend inward, creating the neural groove and neural folds.
• The Zipper-like Fusion: The neural folds elevate and meet in the center of the embryo (around the future neck area) and fuse together like a zipper, moving both upward toward the head (rostrally) and downward toward the spine (caudally). This creates the hollow neural tube.
• Neuropore Closure and Clinical Failures: The anterior neuropore (top opening) closes precisely on day 25, and the posterior neuropore (bottom opening) closes on day 27. If the anterior opening fails to close, the brain cannot form, causing anencephaly. If the posterior opening fails to close, it causes spina bifida.
• Primary and Secondary Vesicle Splitting: Before week 4 ends, the top of the tube expands into three primary vesicles: the Prosencephalon (forebrain), Mesencephalon (midbrain), and Rhombencephalon (hindbrain). By week 5, these split into five secondary vesicles: the forebrain becomes the telencephalon (cerebral cortex, basal ganglia) and diencephalon (thalamus, hypothalamus); the midbrain remains the mesencephalon; the hindbrain splits into the metencephalon (pons, cerebellum) and myelencephalon (medulla).

2. Cellular Genesis, Kinetics, and Migration (Weeks 5–25)

Category: Neurogenesis and Structural Architecture

• The Ventricular Zone Birthplace: All neurons and glial cells are born in the innermost lining of the neural tube, a highly active layer of stem cells called the ventricular zone (VZ).
• Interkinetic Nuclear Migration: As neural progenitor cells prepare to divide, their nuclei physically dance up and down within the VZ. The nucleus moves to the outer edge of the zone to replicate its DNA, then sinks back down to the innermost edge next to the fluid-filled ventricle to undergo mitosis.
• Symmetrical vs. Asymmetrical Division: In the early weeks, progenitor cells undergo symmetrical division (one stem cell splits into two identical stem cells) to rapidly expand the foundational pool. Around week 7, they switch to asymmetrical division: one stem cell splits

into one permanent stem cell and one immature neuron (neuroblast) or glial cell. This marks the peak where up to 250,000 neurons are created every minute.
• Radial Glial Scaffold Climbing: To build the cortex, neuroblasts must migrate centimeters away from their birthplace. Specialized stem cells called radial glia extend long, thin structural fibers spanning from the ventricular zone all the way to the outer surface of the brain. Immature neurons wrap around these fibers and use amoeba-like crawling motions, driven by adhesion molecules like astrotactin, to climb to the surface.
• Inside-Out Cortical Layering: The cerebral cortex is built from the inside out in six distinct layers. The very first wave of migrating neurons forms the deepest layer (Layer VI). The next wave climbs past Layer VI to form Layer V, and the next wave climbs past both to form Layer IV. This continue until all six layers are built, meaning the youngest neurons always occupy the outermost layer (Layer I).

3. Axon Guidance, Synaptogenesis, and Circuit Wiring (Weeks 20–35)

Category: Biochemical Navigation and Network Formation

• Growth Cone Navigation: Once a neuron reaches its destination, it extends a specialized, highly sensitive tip called a growth cone to find its target cells. The growth cone features finger-like extensions (filopodia) that actively sniff out chemical trails in the brain's extracellular matrix.
• Chemoattraction and Chemorepulsion: The growth cone navigate using a precise biological map governed by four families of guidance molecules:
    • Netrins and Semaphorins: Soluble proteins that diffuse through tissue, either attracting the growth cone toward a target or repelling it away.
    • Ephrins and CAMs (Cell Adhesion Molecules): Membrane-bound proteins that require direct physical contact to guide the growing axon over or around specific cell surfaces.
• The Synaptic Handshake: When the growth cone hits its exact target cell, it undergoes a radical transformation. Filopodia retract, and the membrane specializes into a presynaptic terminal. It releases proteins like neurexin, which bind to neuroligin on the receiving cell. This chemical handshake locks the two cells together, forming a functioning synapse capable of transmitting neurotransmitters.
• Trophic Factor Competition: The brain purposefully creates roughly 50% more neurons than will survive into childhood. As axons arrive at their target fields, they must compete for a limited supply of survival chemicals called neurotrophins (such as Nerve Growth Factor, or NGF, and Brain-Derived Neurotrophic Factor, or BDNF). Neurons that successfully absorb enough neurotrophins survive; those that fail undergo apoptosis (programmed cell death), cleanly pruning away the excess wiring to optimize circuit efficiency.

4. Gyrification, Myelination, and Functional Maturation (Week 26–Term)

Category: Mechanical Folding and Electrical Optimization

• Mechanical Tension and Gyrification: Around week 26, the brain transforms from a smooth surface (lissencephalic) to a highly folded surface (gyrencephalic). As billions of neurons

expand and connect, the physical tension of axons pulling between different regions forces the cortex to buckle. This creates the exact ridges (gyri) and valleys (sulci) that organize distinct functional maps, packing a vast surface area into a compact skull.
• Oligodendrocyte Proliferation and Myelination: Specialized glial cells called oligodendrocytes multiply rapidly. They target major axon tracks and begin wrapping their cell membranes around them layer after layer, creating a thick coating of lipid-rich myelin.
• The Spatiotemporal Wave of Myelination: Myelination does not happen all at once; it follows a strict evolutionary and functional hierarchy:
    • Peripheral and Brainstem Tracks: First to myelinate, securing basic life support systems like heart rate, swallowing, and respiration by week 24.
    • Sensory Pathways: Visual and auditory tracks myelinate next, allowing the late-stage fetus to process sounds and flashing lights.
    • Motor and Cortical Tracks: The pathways running down from the motor cortex to the spinal cord begin next, but complex higher-order regions (like the prefrontal cortex) will continue this myelination process well into early adulthood.
• The Birth of Thalamocortical Loops: By week 26 to 28, deep wiring pathways connect the thalamus (the sensory routing station) directly to the cerebral cortex. This is a profound milestone: for the first time, sensory inputs from the body are not just processed as primitive reflexes in the brainstem, but are successfully beamed up to the higher cortex, laying the literal neurological framework for early awareness, pain perception, and organized sleep-wake brain states.
moving forward, I want to discuss the intraparietal group. The intraparietal group has few str uctures. It acts as the split between the superior parietal lobe and inferior parietal lobe.

The dorsal visual stream hub, the IPS, is a core component of the dorsal visual stream. It tra nslates visual information from the occipital lobe into axonal space maps, allowing the brain t o understand where objects are located. It basically maps out the position of the objects. Aga in, we can use this to map out the position, location, or changes.

Through coordination, when we develop our eyes, a specialised subregion within the IPS cal culates the saccades. The rapid voluntary movement of the eyes when jumping from one vis ual target to another prioritises what object in the environment deserves immediate visual foc us. Again, prioritisation is also important.

Visually guided reaching and grasping is important too. The IPS contains dedicated neural cl usters that calculate the exact trajectory. It's somewhat related to the dorsal visual stream hu b, and it's connected to the internal number line. The IPS is the primary neural substrate for numerical processing and mathematical calculation. I think we discussed that, and this will b e good for the symbolic distribution when we discuss the frontal lobes, so the symbolic ones. I hope you remember that too, because I just woke up.

We can generally combine it with the inferior frontal gyrus and the medial frontal gyrus. This also pairs well with the dorsolateral prefrontal cortex, considering the amount of workload it t akes and the manipulation it does.

Non-symbolic estimation: long before the child learns to read numbers, the IPS allows them t o instantly estimate quantities. This is good. This is actually important because this is one of the core parts which develops in common sense. Estimation is also common sense, and I thi

nk we discussed that common sense is in the frontal lobes themselves. For common sense, we pass through DML, if I'm not wrong. I don't remember what the full form was, but I remem ber it was DML, and it was the default mode network for the superior frontal gyrus, the anter omedial region. We can pair this up with that.

Symbolic math calculation: as I discussed for the previous circuit, symbolism can play a reall y crucial role for this.

Working memory and multimodal integration: I think working memory, we discussed that exte nsively in the frontal lobe too, so we can integrate that extensively here too. It acts as a temp orary storage buffer for visual information, dictating a cognitive capacity for how many items you can track. The perceptual saliency mapping: it acts as a cognitive filter that builds salien cy mapping, so it basically filters out, focuses, and prioritises.
Category 1: Phonological Language Processing and Working Memory

• The Phonological Loop Buffer: In the dominant hemisphere (typically the left), the SMG acts as a temporary neural holding station for spoken words, a system known as the phonological loop. It keeps the distinct sound segments of a word active in your mind for several seconds, allowing you to repeat a sequence of words back or hold a phone number in your short-term memory before writing it down.
• Grapheme-to-Phoneme Conversion: The SMG is a vital bridge for reading. It is responsible for decoding written letters (graphemes) and translating them into their corresponding spoken sounds (phonemes), allowing you to successfully sounding out unfamiliar or completely novel words.
• Articulatory Loop Feedback: It works directly with frontal motor language networks (like Broca's area). The SMG monitors the auditory and motor feedback of your own voice to ensure that the words you are physically speaking match the words you intended to say.

Category 2: Complex Praxis and Motor Planning

• Tool-Use Ideation (Ideomotor Praxis): The SMG stores the learned motor plans and complex movement schemas required to use tools and everyday objects. When you pick up a pen, a key, or a hammer, the SMG retrieves the specific internal program that instructs your muscles on the precise sequence of finger movements and wrist angles needed to operate that specific tool.
• Action Execution Tracking: It translates abstract action intentions into real-time physical instructions. It calculates the necessary spatial-motor coordinates to make sure gestures (like waving goodbye or saluting) are executed accurately and fluidly.

Category 3: Somatosensory Integration and Spatial Self-Awareness

• Proprioceptive Egocentric Mapping: The SMG integrates tactile inputs with vestibular (balance) and visual signals to build an internal model of your own physical body. It calculates where your limbs are relative to your torso, forming the bedrock of your personal spatial self-awareness.
• Peripersonal Space Processing: It actively monitors your peripersonal space—the immediate physical zone directly surrounding your body within arms' reach. It registers when

objects or obstacles enter this protective bubble, allowing you to duck, dodge, or reach out to block an impact.

Category 4: Right-Hemisphere Specializations (Attention and Empathy)

• Stimulus-Driven Attentional Shifting: In the non-dominant hemisphere (typically the right), the SMG is a key node in the ventral attentional network. It acts as an environmental tripwire, instantly interrupting your current train of thought to reorient your focus toward unexpected, biologically relevant sensory changes, such as a sudden loud noise or a movement in your peripheral vision.
• Egocentric Empathy Regulation: The right SMG plays a highly specialized role in emotional processing by overcoming the emotional egocentricity bias. It helps you suppress your own current emotional state (e.g., being in a great mood) so that you can accurately perceive, understand, and empathize with someone else's conflicting emotional state (e.g., someone who is grieving). The angular gyrus plays a crucial role in the mathematical one. We have the cortical land. I'll go through the function breakdown.

It acts as visual-to-linguistic transduction. The dominant hemisphere, typically the left, the AG acts as the primary translator for written language. It takes the abstract visual shapes of written words received from the occipital lobe and converts them into internally auditory codes. For example: "hi," "hello, subtitles," "subtitles under a movie, perhaps." Subtitles under a movie generally help with both the occipital lobe and the parietal one. They work together.

Conversion of symbolism, facial expression, and letters, which are also symbols, and registering them as something meaningful, a substance, right? A substance which makes sense, a substance which is coherent.

Then we have grapheme-to-semantic mapping beyond translating letter shapes into sounds. The AG links words directly to their conceptual meaning, right? Again, I think I discussed this just now. It maps the visual form of a written word to its corresponding entry in your mental dictionary.

Then graphic motor output. The AG is a vital engine for autographic writing and spelling. It holds the internal visual and spatial template for letters, directing the motor cortex on how to move.

Then mathematical calculation and verbal memory. For element AG, I think we already discussed how the AG acts as a core part in translating letters, converting letters into coherent words and sentences. Then we have mathematical calculation and verbal memory, right? The left AG is heavily recruited in the exact mathematical calculation that relies on memorised verbal facts, such as multiplication tables or basic addition formulas. These are formulas. Again, this is not logic. This is memorised.

Then abstract symbol processing. I think we discussed symbolism, how the symbols are processed, and how it works up. After we discuss that, it processes numbers not just as visual shapes but as abstract concepts of magnitude and value, working closely with the intraparietal sulcus to manipulate numeric sequencing during multi-step math problems. I

think we discussed this in the frontal lobe too, where the inferior frontal groove, combined with the dorsolateral SRG and the superior frontal gyrus, work together. We can connect these two to make it more efficient.

In spatial navigation and body processing for egocentric and allocentric mapping in the non-dominant hemisphere, typically the right, the AG helps construct your internal spatial maps. It keeps track of where your body is positioned relative to the surroundings. Again, coordinate-wise, we discussed this before it. We discussed this in the marginal one, right?

Then stable visual alignment. The right AG integrates balance signals from the inner ear with the visual output to maintain the sense of physical equilibrium and gravity orientation. It helps with the stable body and locating where your body is relative to your environment and where your environment is.

In self-location and body ownership, it plays a foundational role in the conscious sense of occupying your physical body.

Then default mode network and social cognition. The AG is a core hub for the default mode network. I think we discussed that for the common sense, right? A system that activates when you are daydreaming, introspecting, or even focused on external work. Then theory of mind, that TOM for social interaction. now onto cerebellum.
- Tightly pack billions of computing neurons to run high-speed calculations for micro-adjustments to movemen.. we can use this a storage, it plays a really crucial role in data storage. Folia:
Cellular Micro-Architecture of a Folium

Every single folium across the cerebellum shares an identical, highly organized three-layer cellular matrix that processes information with mathematical precision.

Category 1: The Three Cortical Layers

• The Molecular Layer (Outer): This topmost layer contains a dense network of unmyelinated axons (parallel fibers) and the sprawling dendritic trees of Purkinje cells, alongside inhibitory interneurons (stellate and basket cells) that fine-tune local electrical signals.
• The Purkinje Cell Layer (Middle): This is a single, monolayer sheet of massive Purkinje cells. These are the primary computational units of the cerebellum, and their axons provide the sole inhibitory output leaving the cerebellar cortex.
• The Granular Layer (Inner): This densely packed bottom layer contains billions of tiny granule cells. It acts as the primary receiver station, taking raw sensory and motor inputs from the rest of the body and firing them upward into the molecular layer.

Category 2: Core Afferent and Efferent Wiring

• Mossy Fiber Inputs: These fibers bring raw sensory data from the spinal cord and motor plans from the cerebral cortex into the folium's granular layer, where the data is shattered into millions of parallel streams.
• Climbing Fiber Inputs: Arising strictly from the inferior olivary nucleus in the brainstem, a single climbing fiber wraps around a single Purkinje cell within the folium like ivy. It fires

powerful electrical impulses that act as an "error signal," instructing the folium when a movement mistake has been made.

Functional Breakdown of the Folia

Category 3: Motor Optimization and Micro-Timing

• Feedforward Motor Coordination: The folia do not initiate muscle movements, but they act as a rapid-fire processing grid that calculates exactly how much force, speed, and trajectory are needed for an action. They ensure that complex muscle groups fire in perfect, fluid succession rather than jerking or staggering.
• Microsecond Timing Calibration: The strict geometric layout of the folia allows them to act as internal clocks. They measure the exact microsecond intervals between sensory inputs and motor outputs, which is vital for fast, automated actions like typing, playing an instrument, or catching a flying object.

Category 4: Motor Learning and Error Correction

• Long-Term Depression (LTD): The folia are the biological seat of motor learning (muscle memory). When a climbing fiber delivers an error signal to a Purkinje cell, it structurally alters the sensitivity of that cell to future inputs. This process dampens incorrect movements and strengthens successful ones.
• Predictive Schema Tuning: As you practice a physical skill (like swinging a golf club or driving a car), the folia build internal predictive models. They compare what your brain intended to do with what your body actually did, constantly adjusting muscle commands in real time to erase the gap between intent and execution.

Category 5: Spatial and Cognitive Processing (Posterior Folia)

• Vestibulocerebellar Equilibrium: Folia located in the specialized flocculonodular lobe process inputs from the inner ear to adjust your core posture, stabilize your trunk, and coordinate your eye movements (vestibulo-ocular reflex) so your vision stays stable while you walk.
• Cognitive and Affective Filtering: Modern neuroscience shows that the posterior (rear) folia map directly to the cognitive and emotional centers of the cerebral cortex. They help smooth out transitions in thought, language sequencing, and emotional responses, preventing cognitive or behavioral overshoot in the same way they prevent physical overshooting.
Discussing the parietal lobe is one of the most important things, as you mentioned. We'll go through the grooves first. The grooves, or the spatial boundaries, are divided into three categories:
- the postcentral sulcus
- the intraparietal sulcus
- the parieto-occipital sulcus

Parieto-occipital, if I'm not wrong, is what you see. Again, it maps out the boundaries. I focus on the postcentral sulcus and the intraparietal sulcus.

The postcentral sulcus is a prominent groove located on the lateral surface of the parietal lobe. It serves as the definitive structural boundary that separates the postcentral gyrus anteriorly from the superior and inferior parietal lobules posteriorly.

Functional breakdown of the postcentral sulcus:
1. The somatosensory boundary and spatial organisation. Somatotopic mapping constraints: the anterior wall of the bank of the postcentral sulcus is the posterior limit of the somatosensory homunculus. This region acts as a structural container for the cortical mapping of the human body, where specific zones are dedicated to processing tactile data from the hands, face, torso, and limbs. This is good. This is basically the part which collects data from the human body, not understanding it, but collecting it.
2. Proprioceptive and nociceptive filtering. The sulcus serves as a functional gatekeeper between raw sensations and high-order spatial awareness. It helps isolate incoming signals regarding proprioception. We can focus on proprioception because it can allow self-understanding of the human body, like the body of this, or the understanding of this. We can modify it a bit because, instead of making it all about the body, we can make it about cognition again.
3. Higher-order tactile and spatial processing. The postcentral sulcus is the posterior limit of the somatosensory cortex for higher-order tactile and spatial processing and tactile feature synthesis. While the postcentral gyrus in front of it handles raw touch, the postcentral sulcus transitions into secondary somatosensory processing. It integrates basic tactile features such as texture, roughness, edge direction, and surface texture into cohesive sensory perception, so it helps us understand the configuration of objects.

This can go along with the parts we discussed for the configuration of the objects. I don't remember the parts properly. Let me check. There were two parts that we discussed that deal with the configuration of the object. I forgot it, and I cannot find it for some reason, but I think, I hope you remember, man. Moving forward, the stereognosis, the identification by touch, and the posterior ring of the postcentral sulcus facilitate stereognosis: this cognitive ability to close your eyes, place an unseen object in your hand, and manipulate it with your fingers. Identification from just touch of the objects is important because this will be used right now, if I'm not wrong. We can use this to understand ideas through touch. We can mould it.
We can mould it into ideas that it can touch. One of my ideas is sensory motor integration an d motor feedback, because of its proximity to the motor cortex. The postcentral sulcus plays a major role in sensory motor feedback loops that continuously feed real-time tactile and pos itional data toward the -- moving forward, I want to discuss the intraparietal one. which is above here.
everything we discussed.
I'll give you the baby's timeline

1. Molecular Induction and Neural Tube Formation (Weeks 3–4)

Category: Embryonic Signaling and Neurulation

• Primary Embryonic Induction: Around day 16, the notochord (a primitive structural rod in the embryo) releases specific signaling proteins, primarily Noggin, Chordin, and Follistatin. These proteins block Bone Morphogenetic Proteins (BMPs), which would normally turn the tissue into skin. By blocking BMPs, these signals force the overlying ectoderm to transform into the neural plate.
• Cellular Elongation and Constriction: The cells of the neural plate change shape via microtubule elongation. Microfilaments at the top of these cells contract (apical constriction), pinching the cells like a drawstring. This forces the flat neural plate to bend inward, creating the neural groove and neural folds.
• The Zipper-like Fusion: The neural folds elevate and meet in the center of the embryo (around the future neck area) and fuse together like a zipper, moving both upward toward the head (rostrally) and downward toward the spine (caudally). This creates the hollow neural tube.
• Neuropore Closure and Clinical Failures: The anterior neuropore (top opening) closes precisely on day 25, and the posterior neuropore (bottom opening) closes on day 27. If the anterior opening fails to close, the brain cannot form, causing anencephaly. If the posterior opening fails to close, it causes spina bifida.
• Primary and Secondary Vesicle Splitting: Before week 4 ends, the top of the tube expands into three primary vesicles: the Prosencephalon (forebrain), Mesencephalon (midbrain), and Rhombencephalon (hindbrain). By week 5, these split into five secondary vesicles: the forebrain becomes the telencephalon (cerebral cortex, basal ganglia) and diencephalon (thalamus, hypothalamus); the midbrain remains the mesencephalon; the hindbrain splits into the metencephalon (pons, cerebellum) and myelencephalon (medulla).

2. Cellular Genesis, Kinetics, and Migration (Weeks 5–25)

Category: Neurogenesis and Structural Architecture

• The Ventricular Zone Birthplace: All neurons and glial cells are born in the innermost lining of the neural tube, a highly active layer of stem cells called the ventricular zone (VZ).
• Interkinetic Nuclear Migration: As neural progenitor cells prepare to divide, their nuclei physically dance up and down within the VZ. The nucleus moves to the outer edge of the zone to replicate its DNA, then sinks back down to the innermost edge next to the fluid-filled ventricle to undergo mitosis.
• Symmetrical vs. Asymmetrical Division: In the early weeks, progenitor cells undergo symmetrical division (one stem cell splits into two identical stem cells) to rapidly expand the foundational pool. Around week 7, they switch to asymmetrical division: one stem cell splits into one permanent stem cell and one immature neuron (neuroblast) or glial cell. This marks the peak where up to 250,000 neurons are created every minute.
• Radial Glial Scaffold Climbing: To build the cortex, neuroblasts must migrate centimeters away from their birthplace. Specialized stem cells called radial glia extend long, thin structural fibers spanning from the ventricular zone all the way to the outer surface of the brain. Immature neurons wrap around these fibers and use amoeba-like crawling motions, driven by adhesion molecules like astrotactin, to climb to the surface.
• Inside-Out Cortical Layering: The cerebral cortex is built from the inside out in six distinct layers. The very first wave of migrating neurons forms the deepest layer (Layer VI). The next

wave climbs past Layer VI to form Layer V, and the next wave climbs past both to form Layer IV. This continue until all six layers are built, meaning the youngest neurons always occupy the outermost layer (Layer I).

3. Axon Guidance, Synaptogenesis, and Circuit Wiring (Weeks 20–35)

Category: Biochemical Navigation and Network Formation

• Growth Cone Navigation: Once a neuron reaches its destination, it extends a specialized, highly sensitive tip called a growth cone to find its target cells. The growth cone features finger-like extensions (filopodia) that actively sniff out chemical trails in the brain's extracellular matrix.
• Chemoattraction and Chemorepulsion: The growth cone navigate using a precise biological map governed by four families of guidance molecules:
    • Netrins and Semaphorins: Soluble proteins that diffuse through tissue, either attracting the growth cone toward a target or repelling it away.
    • Ephrins and CAMs (Cell Adhesion Molecules): Membrane-bound proteins that require direct physical contact to guide the growing axon over or around specific cell surfaces.
• The Synaptic Handshake: When the growth cone hits its exact target cell, it undergoes a radical transformation. Filopodia retract, and the membrane specializes into a presynaptic terminal. It releases proteins like neurexin, which bind to neuroligin on the receiving cell. This chemical handshake locks the two cells together, forming a functioning synapse capable of transmitting neurotransmitters.
• Trophic Factor Competition: The brain purposefully creates roughly 50% more neurons than will survive into childhood. As axons arrive at their target fields, they must compete for a limited supply of survival chemicals called neurotrophins (such as Nerve Growth Factor, or NGF, and Brain-Derived Neurotrophic Factor, or BDNF). Neurons that successfully absorb enough neurotrophins survive; those that fail undergo apoptosis (programmed cell death), cleanly pruning away the excess wiring to optimize circuit efficiency.

4. Gyrification, Myelination, and Functional Maturation (Week 26–Term)

Category: Mechanical Folding and Electrical Optimization

• Mechanical Tension and Gyrification: Around week 26, the brain transforms from a smooth surface (lissencephalic) to a highly folded surface (gyrencephalic). As billions of neurons expand and connect, the physical tension of axons pulling between different regions forces the cortex to buckle. This creates the exact ridges (gyri) and valleys (sulci) that organize distinct functional maps, packing a vast surface area into a compact skull.
• Oligodendrocyte Proliferation and Myelination: Specialized glial cells called oligodendrocytes multiply rapidly. They target major axon tracks and begin wrapping their cell membranes around them layer after layer, creating a thick coating of lipid-rich myelin.
• The Spatiotemporal Wave of Myelination: Myelination does not happen all at once; it follows a strict evolutionary and functional hierarchy:

• Peripheral and Brainstem Tracks: First to myelinate, securing basic life support systems like heart rate, swallowing, and respiration by week 24.

• Sensory Pathways: Visual and auditory tracks myelinate next, allowing the late-stage fetus to process sounds and flashing lights.

• Motor and Cortical Tracks: The pathways running down from the motor cortex to the spinal cord begin next, but complex higher-order regions (like the prefrontal cortex) will continue this myelination process well into early adulthood.

• The Birth of Thalamocortical Loops: By week 26 to 28, deep wiring pathways connect the thalamus (the sensory routing station) directly to the cerebral cortex. This is a profound milestone: for the first time, sensory inputs from the body are not just processed as primitive reflexes in the brainstem, but are successfully beamed up to the higher cortex, laying the literal neurological framework for early awareness, pain perception, and organized sleep-wake brain states.

moving forward, I want to discuss the intraparietal group. The intraparietal group has few structures. It acts as the split between the superior parietal lobe and inferior parietal lobe.

The dorsal visual stream hub, the IPS, is a core component of the dorsal visual stream. It translates visual information from the occipital lobe into axonal space maps, allowing the brain to understand where objects are located. It basically maps out the position of the objects. Again, we can use this to map out the position, location, or changes.

Through coordination, when we develop our eyes, a specialised subregion within the IPS calculates the saccades. The rapid voluntary movement of the eyes when jumping from one visual target to another prioritises what object in the environment deserves immediate visual focus. Again, prioritisation is also important.

Visually guided reaching and grasping is important too. The IPS contains dedicated neural clusters that calculate the exact trajectory. It's somewhat related to the dorsal visual stream hub, and it's connected to the internal number line. The IPS is the primary neural substrate for numerical processing and mathematical calculation. I think we discussed that, and this will be good for the symbolic distribution when we discuss the frontal lobes, so the symbolic ones. I hope you remember that too, because I just woke up.

We can generally combine it with the inferior frontal gyrus and the medial frontal gyrus. This also pairs well with the dorsolateral prefrontal cortex, considering the amount of workload it takes and the manipulation it does.

Non-symbolic estimation: long before the child learns to read numbers, the IPS allows them to instantly estimate quantities. This is good. This is actually important because this is one of the core parts which develops in common sense. Estimation is also common sense, and I think we discussed that common sense is in the frontal lobes themselves. For common sense, we pass through DML, if I'm not wrong. I don't remember what the full form was, but I remember it was DML, and it was the default mode network for the superior frontal gyrus, the anteromedial region. We can pair this up with that.

Symbolic math calculation: as I discussed for the previous circuit, symbolism can play a really crucial role for this.

Working memory and multimodal integration: I think working memory, we discussed that exte nsively in the frontal lobe too, so we can integrate that extensively here too. It acts as a temp orary storage buffer for visual information, dictating a cognitive capacity for how many items you can track. The perceptual saliency mapping: it acts as a cognitive filter that builds salien cy mapping, so it basically filters out, focuses, and prioritises.
Category 1: Phonological Language Processing and Working Memory

• The Phonological Loop Buffer: In the dominant hemisphere (typically the left), the SMG acts as a temporary neural holding station for spoken words, a system known as the phonological loop. It keeps the distinct sound segments of a word active in your mind for several seconds, allowing you to repeat a sequence of words back or hold a phone number in your short-term memory before writing it down.
• Grapheme-to-Phoneme Conversion: The SMG is a vital bridge for reading. It is responsible for decoding written letters (graphemes) and translating them into their corresponding spoken sounds (phonemes), allowing you to successfully sounding out unfamiliar or completely novel words.
• Articulatory Loop Feedback: It works directly with frontal motor language networks (like Broca's area). The SMG monitors the auditory and motor feedback of your own voice to ensure that the words you are physically speaking match the words you intended to say.

Category 2: Complex Praxis and Motor Planning

• Tool-Use Ideation (Ideomotor Praxis): The SMG stores the learned motor plans and complex movement schemas required to use tools and everyday objects. When you pick up a pen, a key, or a hammer, the SMG retrieves the specific internal program that instructs your muscles on the precise sequence of finger movements and wrist angles needed to operate that specific tool.
• Action Execution Tracking: It translates abstract action intentions into real-time physical instructions. It calculates the necessary spatial-motor coordinates to make sure gestures (like waving goodbye or saluting) are executed accurately and fluidly.

Category 3: Somatosensory Integration and Spatial Self-Awareness

• Proprioceptive Egocentric Mapping: The SMG integrates tactile inputs with vestibular (balance) and visual signals to build an internal model of your own physical body. It calculates where your limbs are relative to your torso, forming the bedrock of your personal spatial self-awareness.
• Peripersonal Space Processing: It actively monitors your peripersonal space—the immediate physical zone directly surrounding your body within arms' reach. It registers when objects or obstacles enter this protective bubble, allowing you to duck, dodge, or reach out to block an impact.

Category 4: Right-Hemisphere Specializations (Attention and Empathy)

• Stimulus-Driven Attentional Shifting: In the non-dominant hemisphere (typically the right), the SMG is a key node in the ventral attentional network. It acts as an environmental tripwire, instantly interrupting your current train of thought to reorient your focus toward

unexpected, biologically relevant sensory changes, such as a sudden loud noise or a movement in your peripheral vision.
• Egocentric Empathy Regulation: The right SMG plays a highly specialized role in emotional processing by overcoming the emotional egocentricity bias. It helps you suppress your own current emotional state (e.g., being in a great mood) so that you can accurately perceive, understand, and empathize with someone else's conflicting emotional state (e.g., someone who is grieving). The angular gyrus plays a crucial role in the mathematical one. We have the cortical land. I'll go through the function breakdown.

It acts as visual-to-linguistic transduction. The dominant hemisphere, typically the left, the AG acts as the primary translator for written language. It takes the abstract visual shapes of written words received from the occipital lobe and converts them into internally auditory codes. For example: "hi," "hello, subtitles," "subtitles under a movie, perhaps." Subtitles under a movie generally help with both the occipital lobe and the parietal one. They work together.

Conversion of symbolism, facial expression, and letters, which are also symbols, and registering them as something meaningful, a substance, right? A substance which makes sense, a substance which is coherent.

Then we have grapheme-to-semantic mapping beyond translating letter shapes into sounds. The AG links words directly to their conceptual meaning, right? Again, I think I discussed this just now. It maps the visual form of a written word to its corresponding entry in your mental dictionary.

Then graphic motor output. The AG is a vital engine for autographic writing and spelling. It holds the internal visual and spatial template for letters, directing the motor cortex on how to move.

Then mathematical calculation and verbal memory. For element AG, I think we already discussed how the AG acts as a core part in translating letters, converting letters into coherent words and sentences. Then we have mathematical calculation and verbal memory, right? The left AG is heavily recruited in the exact mathematical calculation that relies on memorised verbal facts, such as multiplication tables or basic addition formulas. These are formulas. Again, this is not logic. This is memorised.

Then abstract symbol processing. I think we discussed symbolism, how the symbols are processed, and how it works up. After we discuss that, it processes numbers not just as visual shapes but as abstract concepts of magnitude and value, working closely with the intraparietal sulcus to manipulate numeric sequencing during multi-step math problems. I think we discussed this in the frontal lobe too, where the inferior frontal groove, combined with the dorsolateral SRG and the superior frontal gyrus, work together. We can connect these two to make it more efficient.

In spatial navigation and body processing for egocentric and allocentric mapping in the non-dominant hemisphere, typically the right, the AG helps construct your internal spatial maps. It keeps track of where your body is positioned relative to the surroundings. Again, coordinate-wise, we discussed this before it. We discussed this in the marginal one, right?

Then stable visual alignment. The right AG integrates balance signals from the inner ear with the visual output to maintain the sense of physical equilibrium and gravity orientation. It helps with the stable body and locating where your body is relative to your environment and where your environment is.

In self-location and body ownership, it plays a foundational role in the conscious sense of occupying your physical body.

Then default mode network and social cognition. The AG is a core hub for the default mode network. I think we discussed that for the common sense, right? A system that activates when you are daydreaming, introspecting, or even focused on external work. Then theory of mind, that TOM for social interaction. now onto cerebellum.
- Tightly pack billions of computing neurons to run high-speed calculations for micro-adjustments to movemen.. we can use this a storage, it plays a really crucial role in data storage. Folia:
Cellular Micro-Architecture of a Folium

Every single folium across the cerebellum shares an identical, highly organized three-layer cellular matrix that processes information with mathematical precision.

Category 1: The Three Cortical Layers

• The Molecular Layer (Outer): This topmost layer contains a dense network of unmyelinated axons (parallel fibers) and the sprawling dendritic trees of Purkinje cells, alongside inhibitory interneurons (stellate and basket cells) that fine-tune local electrical signals.
• The Purkinje Cell Layer (Middle): This is a single, monolayer sheet of massive Purkinje cells. These are the primary computational units of the cerebellum, and their axons provide the sole inhibitory output leaving the cerebellar cortex.
• The Granular Layer (Inner): This densely packed bottom layer contains billions of tiny granule cells. It acts as the primary receiver station, taking raw sensory and motor inputs from the rest of the body and firing them upward into the molecular layer.

Category 2: Core Afferent and Efferent Wiring

• Mossy Fiber Inputs: These fibers bring raw sensory data from the spinal cord and motor plans from the cerebral cortex into the folium's granular layer, where the data is shattered into millions of parallel streams.
• Climbing Fiber Inputs: Arising strictly from the inferior olivary nucleus in the brainstem, a single climbing fiber wraps around a single Purkinje cell within the folium like ivy. It fires powerful electrical impulses that act as an "error signal," instructing the folium when a movement mistake has been made.

Functional Breakdown of the Folia

Category 3: Motor Optimization and Micro-Timing

• Feedforward Motor Coordination: The folia do not initiate muscle movements, but they act as a rapid-fire processing grid that calculates exactly how much force, speed, and trajectory are needed for an action. They ensure that complex muscle groups fire in perfect, fluid succession rather than jerking or staggering.
• Microsecond Timing Calibration: The strict geometric layout of the folia allows them to act as internal clocks. They measure the exact microsecond intervals between sensory inputs and motor outputs, which is vital for fast, automated actions like typing, playing an instrument, or catching a flying object.

Category 4: Motor Learning and Error Correction

• Long-Term Depression (LTD): The folia are the biological seat of motor learning (muscle memory). When a climbing fiber delivers an error signal to a Purkinje cell, it structurally alters the sensitivity of that cell to future inputs. This process dampens incorrect movements and strengthens successful ones.
• Predictive Schema Tuning: As you practice a physical skill (like swinging a golf club or driving a car), the folia build internal predictive models. They compare what your brain intended to do with what your body actually did, constantly adjusting muscle commands in real time to erase the gap between intent and execution.

Category 5: Spatial and Cognitive Processing (Posterior Folia)

• Vestibulocerebellar Equilibrium: Folia located in the specialized flocculonodular lobe process inputs from the inner ear to adjust your core posture, stabilize your trunk, and coordinate your eye movements (vestibulo-ocular reflex) so your vision stays stable while you walk.
• Cognitive and Affective Filtering: Modern neuroscience shows that the posterior (rear) folia map directly to the cognitive and emotional centers of the cerebral cortex. They help smooth out transitions in thought, language sequencing, and emotional responses, preventing cognitive or behavioral overshoot in the same way they prevent physical overshooting.
Discussing the parietal lobe is one of the most important things, as you mentioned. We'll go through the grooves first. The grooves, or the spatial boundaries, are divided into three categories:
- the postcentral sulcus
- the intraparietal sulcus
- the parieto-occipital sulcus

Parieto-occipital, if I'm not wrong, is what you see. Again, it maps out the boundaries. I focus on the postcentral sulcus and the intraparietal sulcus.

The postcentral sulcus is a prominent groove located on the lateral surface of the parietal lobe. It serves as the definitive structural boundary that separates the postcentral gyrus anteriorly from the superior and inferior parietal lobules posteriorly.

Functional breakdown of the postcentral sulcus:
1. The somatosensory boundary and spatial organisation. Somatotopic mapping constraints: the anterior wall of the bank of the postcentral sulcus is the posterior limit of the

somatosensory homunculus. This region acts as a structural container for the cortical mapping of the human body, where specific zones are dedicated to processing tactile data from the hands, face, torso, and limbs. This is good. This is basically the part which collects data from the human body, not understanding it, but collecting it.
2. Proprioceptive and nociceptive filtering. The sulcus serves as a functional gatekeeper between raw sensations and high-order spatial awareness. It helps isolate incoming signals regarding proprioception. We can focus on proprioception because it can allow self-understanding of the human body, like the body of this, or the understanding of this. We can modify it a bit because, instead of making it all about the body, we can make it about cognition again.
3. Higher-order tactile and spatial processing. The postcentral sulcus is the posterior limit of the somatosensory cortex for higher-order tactile and spatial processing and tactile feature synthesis. While the postcentral gyrus in front of it handles raw touch, the postcentral sulcus transitions into secondary somatosensory processing. It integrates basic tactile features such as texture, roughness, edge direction, and surface texture into cohesive sensory perception, so it helps us understand the configuration of objects.

This can go along with the parts we discussed for the configuration of the objects. I don't remember the parts properly. Let me check. There were two parts that we discussed that deal with the configuration of the object. I forgot it, and I cannot find it for some reason, but I think, I hope you remember, man. Moving forward, the stereognosis, the identification by touch, and the posterior ring of the postcentral sulcus facilitate stereognosis: this cognitive ability to close your eyes, place an unseen object in your hand, and manipulate it with your fingers. Identification from just touch of the objects is important because this will be used right now, if I'm not wrong. We can use this to understand ideas through touch. We can mould it.
We can mould it into ideas that it can touch. One of my ideas is sensory motor integration an d motor feedback, because of its proximity to the motor cortex. The postcentral sulcus plays a major role in sensory motor feedback loops that continuously feed real-time tactile and pos itional data toward the -- moving forward, I want to discuss the intraparietal one. which is above here.
Brain
1. connect the three brains
for sharpening of ideas, check-in results, Turn each big idea into a test that could fail, Let each life stage switch ideas on when the biology does
we will discuss that.
———————
Frontal Lobe:
Central Sulcus (Groove)
Physical boundary between the thinking/moving (frontal) & feeling (parietal). -- deeper central sulcus can contribute to extraordinary dexterity, faster motor learning, and a hyper-acute sense of touch. Which can help with the overall sensory understanding, which we discussed.

Precentral Sulcus
It serves as the physical border that separates your primary movement center from your high-level planning and decision-making centers. For premotor cortex & supplementary

motor area, I was thinking of enhancing it to push for a more possible complex chain of actions. when we will provide the baby with a library, and understanding of physical bounds. The Broca's area.. we will try to make it stronger so the interaction and overall clarity of ideas of novelty remains, pair this up with frontal eye speed -- possible hypothesis : "If Broca's area and the Frontal Eye Fields (FEF) were structurally deeper and had hyper-direct, heavily myelinated connections leading straight into the central sulcus".

Okay, moving forward to the temporal lobe. This one was the one which was lacking the most. Let's go ahead with it.

First, we have the grooves, the sulci, meaning enclaves. Again, enclaves, grooves, lateral fissure: the massive structural chasm protecting the deep language and the auditory centres. The superior and inferior temporal sulci demarcate the auditory processing zones from the visual object recognition zones. The collateral and rhinal sulci isolate the deep, ancient emotional structures from the outer evolutionary neocortex.

Okay, moving forward to the temporal lobe. This one was the one which was lacking the most. Let's go ahead with it.

First, we have the grooves, the sulci, meaning enclaves. Again, enclaves, grooves, lateral fissure: the massive structural chasm protecting the deep language and the auditory centres. The superior and inferior temporal sulci demarcate the auditory processing zones from the visual object recognition zones. The collateral and rhinal sulci isolate the deep, ancient emotional structures from the outer evolutionary neocortex.

Here, we can focus on the collateral and rhinal sulci, or the collateral and rhinal groove, which, again, will be useful for the development of both the emotional or the ethical development and the evolutionary cortex, which can look for processing of information. Moving forward, I would go for... wait, one second.
Here, we can focus on the collateral and rhinal sulci, or the collateral and rhinal groove, which, again, will be useful for the development of both the emotional or the ethical development and the evolutionary cortex, which can look for processing of information. Moving forward, I would go for... wait, one second. A few parts got cut: the lateral fissure and the superior and inferior temporal sulci, or, again, grooves. It's better when I say grooves.

A bit about the lateral fissure is the fact that, again, the lateral fissure is a part of the temporal lobe. It constitutes the sulci that are present. The lateral fissure, or the Sylvian fissure, I think we discussed this before, was actually quite unusual in Einstein's brain. It is the most prominent deep horizontal trench on the side of the human brain. If the central sulcus is the vertical line, the lateral fissure is the horizontal canyon that splits the temporal lobe completely away from the frontal and parietal lobes. It divides the three lobes.

It is the centre of human language, high-speed auditory processing, and internal self-regulation. Human language is a part of baby code's understanding of the whole thing, so we have to be using the Sylvian fissure. I would rather like to see how Einstein's Sylvian fissure was actually formed. It will be helpful for our brain formation. Moving forward, let's move on to:

- the language highway
- the parietal network
- the front bank: the inferior frontal gyrus. I think we discussed this: the IFG, Broca's area, which helps with the processing of language, specifically the pars opercularis. We discuss this too: it generates the motor plans for speech.
- the back bank: the Vernix area, the processing hub that translates heard or read words into meaningful thoughts.

We can use these specific areas for the recognition of patterns and interpretations of symbols into coherent codes, right? Codes which could be understood by it.

Auditory processing, we will discuss this later. Right now, I will just focus on cognition and touch.
Okay, then. The hidden island, the insula, deep within the depths of the Sylvian fissure, completely buried beneath the overlapping folds of the frontal, parietal, and temporal lobes, has a hidden fifth lobe called the insula.

The insula is the master centre of interoception. It's like your conscious awareness of your body's internal state, so basically awareness within the body. This is actually quite helpful because we can pair this with optogenetics so that the body can understand how it is working. Remember, we were talking about self-regulation earlier in the chat. We can base it off of that. Moving forward, we discuss the lateral fissure, the deeper part, obviously, then the superior and inferior temporal sulci. Let's discuss that, then we can discuss collateral and rhinal sulci.

Superior and inferior temporal sulci. I'll start a new chat window. Is this in detail? Please wait a second.

The superior temporal sulcus runs parallel to the Sylvian fissure and separates the superior temporal sulcus. Add again: the superior temporal sulcus and the inferior temporal sulcus are sulci, not circuses. They serve as the functional boundaries and processing hubs for vastly different neural networks.

The functional mapping of the superior temporal sulcus first helps in social cognition and the social brain when walking, eye gaze, and face processing. I think we discussed the theory of mind before too. The superior temporal sulcus somewhat assists in that too. Let me continue once I get okay, moving forward, let us discuss the STS, social perception, and body language. The STS is the absolute epicentre for reading other living beings. It activates heavily. You track using biological motions, such as reading someone's facial expression and watching their eyes.

This is quite similar to what we discussed previously, which was the Sylvian fissure, or part of it, not really the Sylvian fissure directly, if I'm not wrong. We connected that to the idea of symbolic representation and processing of that symbolic representation. We can use that here too.

We also talk about the specific part of the brain which analyses mistakes, if I'm not wrong. That specific part of the brain was the ACC, right? The ACC is the anterior cingulate cortex, so we can use that for hi there. Again, it is useful as well.

Moving forward, we move on to the inferior temporal sulcus, the object cataloger, and it forms the crucial part of the ventral stream. While the upper network tracks movement, the tissue around the inferior temporal sulcus works to identify exactly what an object is. It processes shapes. I think we discussed this before too: there is a specific part in the brain which processes the configuration of an object, isolating the object's configuration and understanding it. We even comprehensively discussed that too, so we can connect the inferior temporal sulcus and that to hi there, right?

Moving forward, let us move on to the rhinal and the collateral one. Okay, the memory and recognition, the entorhinal hub, the spatial GPS.

The tissue around the rhinal sulcus is the primary interface between the cerebral cortex and the hippocampus. It contains the grid cells, which act like your brain's internal GPS, creating a cognitive map of its physical environment so you can navigate through space.

This is good. This can be a method for us, for the web code to navigate itself through the region of the internet. Once we open source it, we can give it the ability to understand where it is. We need to make it a mix so that it can be self-sufficient as long as it develops like a human. We cannot have a mother to cordle it, right?

Then we have the parietal cortex. The parietal region is responsible for identifying whether you have encountered an object or environment before. It gives you the distinct flash of recognition. It's memory, right? Again, wire it up to the memory one.

Then the collateral sulcus, the scene and the landscape decoder, the context and environment recognizer. The outer bank of the collateral sulcus houses the parahippocampal place area, the PPA. This specialises region activates intensely when you look at scenes, landscape, buildings, or a room. It does not look at an isolated object. It looks at the larger picture.

Looking at the larger picture can be used as a way of problem solving, which I think we discussed in the frontal lobe. In that frontal lobe, we discussed the fluidity of how something can be solved, right? That was the middle frontal gyrus, which constructs DAN and BAN, so we can work on that. Then memory encoding, again, we discuss this. Okay. Okay, the memory and recognition, the entorhinal hub, the spatial GPS.

The tissue around the rhinal sulcus is the primary interface between the cerebral cortex and the hippocampus. It contains the grid cells, which act like your brain's internal GPS, creating a cognitive map of its physical environment so you can navigate through space.

This is good. This can be a method for us, for the web code to navigate itself through the region of the internet. Once we open source it, we can give it the ability to understand where it is. We need to make it a mix so that it can be self-sufficient as long as it develops like a human. We cannot have a mother to cordle it, right?

Then we have the parietal cortex. The parietal region is responsible for identifying whether you have encountered an object or environment before. It gives you the distinct flash of recognition. It's memory, right? Again, wire it up to the memory one.

Then the collateral sulcus, the scene and the landscape decoder, the context and environment recognizer. The outer bank of the collateral sulcus houses the parahippocampal place area, the PPA. This specialises region activates intensely when you look at scenes, landscape, buildings, or a room. It does not look at an isolated object. It looks at the larger picture.

Looking at the larger picture can be used as a way of problem solving, which I think we discussed in the frontal lobe. In that frontal lobe, we discussed the fluidity of how something can be solved, right? That was the middle frontal gyrus, which constructs DAN and BAN, so we can work on that. Then memory encoding, again, we discuss this. Okay.
1. Heschl’s Gyri (Primary Auditory Cortex)

Category: Tonotopic Mapping and Spatial Acoustics

• Primary Auditory Processing: Slipped inside the lateral fissure on the upper surface of the temporal lobe, Heschl's gyri serve as the brain's gateway for sound. It is the first cortical structure to receive raw, unprocessed auditory signals traveling directly from the ears via the cochlea and the medial geniculate body of the thalamus.
• Tonotopic Organization: The tissue within Heschl's gyri is organized precisely by sound frequency, functioning like an internal piano keyboard. Neurons at one end are highly tuned to fire only to high-pitched sounds, while neurons at the opposite end fire strictly in response to low-pitched frequencies.
• Basic Feature Extraction: This gyrus breaks down sound waves into their foundational physical properties. It isolates the exact pitch (frequency), volume (amplitude), timbre, and baseline duration of a sound before passing these individual building blocks to adjacent areas for comprehension.
• Binaural Localization: By calculating the microsecond differences in when a sound reaches the left ear versus the right ear, Heschl's gyri calculate the directional origin of a sound, letting you map where a noise is coming from in your environment.

2. Superior Temporal Gyrus (STG)

Category: Secondary Auditory Processing and Language Comprehension

• Complex Acoustic Synthesis: Wrapping around Heschl's gyri, the STG takes the raw physical building blocks of sound and fuses them into recognizable, complex auditory patterns. It allows you to tell the difference between footsteps, a dog barking, a musical instrument, or a human voice.
• Phonological Processing: The STG acts as the primary deciphering engine for spoken language. It analyzes changing frequencies over time to identify phonemes—the specific, individual sounds that make up words (such as distinguishing the sound of a "B" from a "P").

• Wernicke's Area Integration: The posterior (rear) portion of the left STG houses Wernicke's area, the critical neural hub for language comprehension. It decodes the syntactic structure of spoken sentences, allowing you to understand the actual meaning behind words spoken by others.
• Prosody and Emotional Mapping: While the left STG handles the linguistic meaning of words, the right STG evaluates the melody, rhythm, and intonation of speech (prosody). It decodes whether a speaker is angry, sarcastic, happy, or sad based entirely on their vocal tone.

3. Middle Temporal Gyrus (MTG)

Category: Lexical Storage and Multimodal Integration

• The Mental Lexicon: The MTG serves as the brain's primary library for words. It stores your vocabulary, acting as a massive neural dictionary where word forms are linked directly to their definitions and grammatical profiles.
• Semantic Context Retrieval: During a conversation, the MTG rapidly retrieves words from memory so you can parse sentences in real time. It resolves ambiguity, ensuring that if someone says the word "bank," your brain uses the surrounding conversation to instantly know if they mean a riverbank or a financial institution.
• Action and Tool Concepts: The MTG is heavily involved in processing verbs, action words, and tool concepts. When you look at an object like a hammer or read the word "run," neurons in the MTG fire to retrieve the conceptual memory of how that object is physically manipulated or how that action is performed.
• Multimodal Sensory Integration: Positioned symmetrically between the auditory cortex above and the visual cortex below, the MTG acts as a processing bridge. It takes visual inputs (like reading a word or seeing a picture) and auditory inputs (like hearing a word) and merges them into a single, unified conceptual idea.

4. Inferior Temporal Gyrus (ITG)

Category: Ventral Stream Visual Identification

• High-Level Visual Form Synthesis: The ITG occupies the lower lateral ridge of the temporal lobe and represents the visual processing peak of the ventral stream (the "what" pathway). It takes highly complex visual inputs—such as varying lines, shadows, colors, and textures—and synthesizes them into highly structured, recognizable shapes.
• Viewpoint Invariant Object Recognition: A core marvel of the ITG is its ability to recognize objects regardless of the angle from which they are viewed. If you look at a coffee mug from the top, the side, or upside down, the ITG ensures your brain still instantly identifies it as the exact same object.
• Perceptual Constancy: It maintains visual stability by tracking the structural reality of an item rather than its lighting environment. The ITG allows you to recognize an object whether it is cloaked in deep shadow, brightly illuminated by sunlight, or partially hidden behind something else.

5. Fusiform Gyrus (Lateral Occipitotemporal Gyrus)

Category: Expert Visual Recognition and Category Specialization

• Holistic Face Processing (Fusiform Face Area): A specialized region on the lateral side of the fusiform gyrus is explicitly dedicated to facial recognition. Instead of analyzing eyes, noses, and mouths as isolated parts, it processes faces holistically, mapping the exact spatial geometry of a face to tell you if you are looking at a stranger, a celebrity, or a family member.
• Visual Word Recognition (Visual Word Form Area): Located on the left side of the fusiform gyrus, this area is highly specialized for reading. It memorizes the visual shape of letters, character strings, and whole words, instantly recognizing written language configurations and passing them to language centers for rapid reading.
• Within-Category Visual Expertise: Beyond faces and text, the fusiform gyrus adapts to become a hub for deep visual expertise. If a person becomes an expert car mechanic, an avid birdwatcher, or a chess master, this gyrus structurally adapts to rapidly identify specific car models, bird species, or complex chess board configurations.

6. Parahippocampal Gyrus

Category: Spatial Navigation and Contextual Memory Encoding

• Environmental Scene Processing (Parahippocampal Place Area): The PPA is a highly specialized zone within this gyrus that lights up when you look at environmental scenes. It analyzes the structural layout of landscapes, city streets, building interiors, and rooms, mapping the physical geometry of your surroundings rather than individual objects.
• Memory Gateway to the Hippocampus: Wrapping directly around the hippocampus, this gyrus acts as the primary physical conduit for memory. It gathers sensory information from across the entire cerebral cortex, organizes it, and feeds it into the hippocampus so it can be packaged and stored as a long-term episodic memory.
• Spatial Navigation and Cognitive Mapping: The parahippocampal gyrus creates and maintains internal cognitive maps of the physical world. It keeps track of where you are located within a city or house, helping you navigate through space, remember routes, and orient yourself when you walk into an unfamiliar environment.
• Contextual Association: It links memories to specific contexts. When you encounter a specific smell, sight, or sound, the parahippocampal gyrus retrieves the holistic memory of the specific place, time, and situational backdrop where you originally experienced it.

1. Primary Auditory Cortex (A1)

Category: Sensory Transduction and Acoustic Analysis

• Acoustic Wave Decomposition: Located within Heschl's gyri, A1 is the ultimate terminal for raw auditory data. It receives neural impulses sent from the cochlea via the auditory nerve and thalamus, translating physical sound vibrations into conscious auditory perceptions.

• Spectral Frequency Mapping: A1 relies on a highly organized tonotopic grid where neighboring columns of neurons process neighboring sound frequencies. The anterior section maps low-frequency bass tones, while the posterior section decodes high-frequency treble tones.
• Intensity and Temporal Gating: Beyond pitch, A1 isolates the precise amplitude (volume) and microsecond timing of sound waves. It notes exactly when a sound starts, fluctuates, and stops, feeding this temporal structure to higher-level language and music centers.

2. Wernicke’s Area

Category: Semantic Decoding and Syntactic Assembly

• Linguistic Meaning Extraction: Situated in the posterior portion of the superior temporal gyrus (typically in the left hemisphere), Wernicke's area transforms acoustic phonemes into meaningful words. It acts as the brain's internal linguistic dictionary, matching auditory signals to stored conceptual definitions.
• Syntactic Structure Parsing: This area decodes the grammatical architecture of spoken and written sentences. It allows you to process the relationships between nouns, verbs, and prepositions so you can understand complex statements like "The dog chased the cat" versus "The cat chased the dog."
• Speech Generation Blueprinting: Before you speak, Wernicke's area constructs the conceptual
Brain
1. connect the three brains
for sharpening of ideas, check-in results, Turn each big idea into a test that could fail, Let each life stage switch ideas on when the biology does
we will discuss that.
———————
Frontal Lobe:
Central Sulcus (Groove)
Physical boundary between the thinking/moving (frontal) & feeling (parietal). -- deeper central sulcus can contribute to extraordinary dexterity, faster motor learning, and a hyper-acute sense of touch. Which can help with the overall sensory understanding, which we discussed.

Precentral Sulcus
It serves as the physical border that separates your primary movement center from your high-level planning and decision-making centers. For premotor cortex & supplementary motor area, I was thinking of enhancing it to push for a more possible complex chain of actions. when we will provide the baby with a library, and understanding of physical bounds. The Broca's area.. we will try to make it stronger so the interaction and overall clarity of ideas of novelty remains, pair this up with frontal eye speed -- possible hypothesis : "If Broca’s area and the Frontal Eye Fields (FEF) were structurally deeper and had hyper-direct, heavily myelinated connections leading straight into the central sulcus".

Okay, moving forward to the temporal lobe. This one was the one which was lacking the most. Let's go ahead with it.

First, we have the grooves, the sulci, meaning enclaves. Again, enclaves, grooves, lateral fissure: the massive structural chasm protecting the deep language and the auditory centres. The superior and inferior temporal sulci demarcate the auditory processing zones from the visual object recognition zones. The collateral and rhinal sulci isolate the deep, ancient emotional structures from the outer evolutionary neocortex.

Okay, moving forward to the temporal lobe. This one was the one which was lacking the most. Let's go ahead with it.

First, we have the grooves, the sulci, meaning enclaves. Again, enclaves, grooves, lateral fissure: the massive structural chasm protecting the deep language and the auditory centres. The superior and inferior temporal sulci demarcate the auditory processing zones from the visual object recognition zones. The collateral and rhinal sulci isolate the deep, ancient emotional structures from the outer evolutionary neocortex.

Here, we can focus on the collateral and rhinal sulci, or the collateral and rhinal groove, which, again, will be useful for the development of both the emotional or the ethical development and the evolutionary cortex, which can look for processing of information. Moving forward, I would go for... wait, one second.
Here, we can focus on the collateral and rhinal sulci, or the collateral and rhinal groove, which, again, will be useful for the development of both the emotional or the ethical development and the evolutionary cortex, which can look for processing of information. Moving forward, I would go for... wait, one second. A few parts got cut: the lateral fissure and the superior and inferior temporal sulci, or, again, grooves. It's better when I say grooves.

A bit about the lateral fissure is the fact that, again, the lateral fissure is a part of the temporal lobe. It constitutes the sulci that are present. The lateral fissure, or the Sylvian fissure, I think we discussed this before, was actually quite unusual in Einstein's brain. It is the most prominent deep horizontal trench on the side of the human brain. If the central sulcus is the vertical line, the lateral fissure is the horizontal canyon that splits the temporal lobe completely away from the frontal and parietal lobes. It divides the three lobes.

It is the centre of human language, high-speed auditory processing, and internal self-regulation. Human language is a part of baby code's understanding of the whole thing, so we have to be using the Sylvian fissure. I would rather like to see how Einstein's Sylvian fissure was actually formed. It will be helpful for our brain formation. Moving forward, let's move on to:
- the language highway
- the parietal network
- the front bank: the inferior frontal gyrus. I think we discussed this: the IFG, Broca's area, which helps with the processing of language, specifically the pars opercularis. We discuss this too: it generates the motor plans for speech.
- the back bank: the Vernix area, the processing hub that translates heard or read words into meaningful thoughts.

We can use these specific areas for the recognition of patterns and interpretations of symbols into coherent codes, right? Codes which could be understood by it.

Auditory processing, we will discuss this later. Right now, I will just focus on cognition and touch.
Okay, then. The hidden island, the insula, deep within the depths of the Sylvian fissure, completely buried beneath the overlapping folds of the frontal, parietal, and temporal lobes, has a hidden fifth lobe called the insula.

The insula is the master centre of interoception. It's like your conscious awareness of your body's internal state, so basically awareness within the body. This is actually quite helpful because we can pair this with optogenetics so that the body can understand how it is working. Remember, we were talking about self-regulation earlier in the chat. We can base it off of that. Moving forward, we discuss the lateral fissure, the deeper part, obviously, then the superior and inferior temporal sulci. Let's discuss that, then we can discuss collateral and rhinal sulci.

Superior and inferior temporal sulci. I'll start a new chat window. Is this in detail? Please wait a second.

The superior temporal sulcus runs parallel to the Sylvian fissure and separates the superior temporal sulcus. Add again: the superior temporal sulcus and the inferior temporal sulcus are sulci, not circuses. They serve as the functional boundaries and processing hubs for vastly different neural networks.

The functional mapping of the superior temporal sulcus first helps in social cognition and the social brain when walking, eye gaze, and face processing. I think we discussed the theory of mind before too. The superior temporal sulcus somewhat assists in that too. Let me continue once I get okay, moving forward, let us discuss the STS, social perception, and body language. The STS is the absolute epicentre for reading other living beings. It activates heavily. You track using biological motions, such as reading someone's facial expression and watching their eyes.

This is quite similar to what we discussed previously, which was the Sylvian fissure, or part of it, not really the Sylvian fissure directly, if I'm not wrong. We connected that to the idea of symbolic representation and processing of that symbolic representation. We can use that here too.

We also talk about the specific part of the brain which analyses mistakes, if I'm not wrong. That specific part of the brain was the ACC, right? The ACC is the anterior cingulate cortex, so we can use that for hi there. Again, it is useful as well.

Moving forward, we move on to the inferior temporal sulcus, the object cataloger, and it forms the crucial part of the ventral stream. While the upper network tracks movement, the tissue around the inferior temporal sulcus works to identify exactly what an object is. It processes shapes. I think we discussed this before too: there is a specific part in the brain which processes the configuration of an object, isolating the object's configuration and

understanding it. We even comprehensively discussed that too, so we can connect the inferior temporal sulcus and that to hi there, right?

Moving forward, let us move on to the rhinal and the collateral one. Okay, the memory and recognition, the entorhinal hub, the spatial GPS.

The tissue around the rhinal sulcus is the primary interface between the cerebral cortex and the hippocampus. It contains the grid cells, which act like your brain's internal GPS, creating a cognitive map of its physical environment so you can navigate through space.

This is good. This can be a method for us, for the web code to navigate itself through the region of the internet. Once we open source it, we can give it the ability to understand where it is. We need to make it a mix so that it can be self-sufficient as long as it develops like a human. We cannot have a mother to cordle it, right?

Then we have the parietal cortex. The parietal region is responsible for identifying whether you have encountered an object or environment before. It gives you the distinct flash of recognition. It's memory, right? Again, wire it up to the memory one.

Then the collateral sulcus, the scene and the landscape decoder, the context and environment recognizer. The outer bank of the collateral sulcus houses the parahippocampal place area, the PPA. This specialises region activates intensely when you look at scenes, landscape, buildings, or a room. It does not look at an isolated object. It looks at the larger picture.

Looking at the larger picture can be used as a way of problem solving, which I think we discussed in the frontal lobe. In that frontal lobe, we discussed the fluidity of how something can be solved, right? That was the middle frontal gyrus, which constructs DAN and BAN, so we can work on that. Then memory encoding, again, we discuss this. Okay. Okay, the memory and recognition, the entorhinal hub, the spatial GPS.

The tissue around the rhinal sulcus is the primary interface between the cerebral cortex and the hippocampus. It contains the grid cells, which act like your brain's internal GPS, creating a cognitive map of its physical environment so you can navigate through space.

This is good. This can be a method for us, for the web code to navigate itself through the region of the internet. Once we open source it, we can give it the ability to understand where it is. We need to make it a mix so that it can be self-sufficient as long as it develops like a human. We cannot have a mother to cordle it, right?

Then we have the parietal cortex. The parietal region is responsible for identifying whether you have encountered an object or environment before. It gives you the distinct flash of recognition. It's memory, right? Again, wire it up to the memory one.

Then the collateral sulcus, the scene and the landscape decoder, the context and environment recognizer. The outer bank of the collateral sulcus houses the parahippocampal place area, the PPA. This specialises region activates intensely when you look at scenes,

landscape, buildings, or a room. It does not look at an isolated object. It looks at the larger picture.

Looking at the larger picture can be used as a way of problem solving, which I think we discussed in the frontal lobe. In that frontal lobe, we discussed the fluidity of how something can be solved, right? That was the middle frontal gyrus, which constructs DAN and BAN, so we can work on that. Then memory encoding, again, we discuss this. Okay.
1. Heschl’s Gyri (Primary Auditory Cortex)

Category: Tonotopic Mapping and Spatial Acoustics

• Primary Auditory Processing: Slipped inside the lateral fissure on the upper surface of the temporal lobe, Heschl's gyri serve as the brain's gateway for sound. It is the first cortical structure to receive raw, unprocessed auditory signals traveling directly from the ears via the cochlea and the medial geniculate body of the thalamus.
• Tonotopic Organization: The tissue within Heschl's gyri is organized precisely by sound frequency, functioning like an internal piano keyboard. Neurons at one end are highly tuned to fire only to high-pitched sounds, while neurons at the opposite end fire strictly in response to low-pitched frequencies.
• Basic Feature Extraction: This gyrus breaks down sound waves into their foundational physical properties. It isolates the exact pitch (frequency), volume (amplitude), timbre, and baseline duration of a sound before passing these individual building blocks to adjacent areas for comprehension.
• Binaural Localization: By calculating the microsecond differences in when a sound reaches the left ear versus the right ear, Heschl's gyri calculate the directional origin of a sound, letting you map where a noise is coming from in your environment.

2. Superior Temporal Gyrus (STG)

Category: Secondary Auditory Processing and Language Comprehension

• Complex Acoustic Synthesis: Wrapping around Heschl's gyri, the STG takes the raw physical building blocks of sound and fuses them into recognizable, complex auditory patterns. It allows you to tell the difference between footsteps, a dog barking, a musical instrument, or a human voice.
• Phonological Processing: The STG acts as the primary deciphering engine for spoken language. It analyzes changing frequencies over time to identify phonemes—the specific, individual sounds that make up words (such as distinguishing the sound of a "B" from a "P").
• Wernicke's Area Integration: The posterior (rear) portion of the left STG houses Wernicke’s area, the critical neural hub for language comprehension. It decodes the syntactic structure of spoken sentences, allowing you to understand the actual meaning behind words spoken by others.
• Prosody and Emotional Mapping: While the left STG handles the linguistic meaning of words, the right STG evaluates the melody, rhythm, and intonation of speech (prosody). It decodes whether a speaker is angry, sarcastic, happy, or sad based entirely on their vocal tone.

3. Middle Temporal Gyrus (MTG)

Category: Lexical Storage and Multimodal Integration

• The Mental Lexicon: The MTG serves as the brain's primary library for words. It stores your vocabulary, acting as a massive neural dictionary where word forms are linked directly to their definitions and grammatical profiles.
• Semantic Context Retrieval: During a conversation, the MTG rapidly retrieves words from memory so you can parse sentences in real time. It resolves ambiguity, ensuring that if someone says the word "bank," your brain uses the surrounding conversation to instantly know if they mean a riverbank or a financial institution.
• Action and Tool Concepts: The MTG is heavily involved in processing verbs, action words, and tool concepts. When you look at an object like a hammer or read the word "run," neurons in the MTG fire to retrieve the conceptual memory of how that object is physically manipulated or how that action is performed.
• Multimodal Sensory Integration: Positioned symmetrically between the auditory cortex above and the visual cortex below, the MTG acts as a processing bridge. It takes visual inputs (like reading a word or seeing a picture) and auditory inputs (like hearing a word) and merges them into a single, unified conceptual idea.

4. Inferior Temporal Gyrus (ITG)

Category: Ventral Stream Visual Identification

• High-Level Visual Form Synthesis: The ITG occupies the lower lateral ridge of the temporal lobe and represents the visual processing peak of the ventral stream (the "what" pathway). It takes highly complex visual inputs—such as varying lines, shadows, colors, and textures—and synthesizes them into highly structured, recognizable shapes.
• Viewpoint Invariant Object Recognition: A core marvel of the ITG is its ability to recognize objects regardless of the angle from which they are viewed. If you look at a coffee mug from the top, the side, or upside down, the ITG ensures your brain still instantly identifies it as the exact same object.
• Perceptual Constancy: It maintains visual stability by tracking the structural reality of an item rather than its lighting environment. The ITG allows you to recognize an object whether it is cloaked in deep shadow, brightly illuminated by sunlight, or partially hidden behind something else.

5. Fusiform Gyrus (Lateral Occipitotemporal Gyrus)

Category: Expert Visual Recognition and Category Specialization

• Holistic Face Processing (Fusiform Face Area): A specialized region on the lateral side of the fusiform gyrus is explicitly dedicated to facial recognition. Instead of analyzing eyes, noses, and mouths as isolated parts, it processes faces holistically, mapping the exact

spatial geometry of a face to tell you if you are looking at a stranger, a celebrity, or a family member.
• Visual Word Recognition (Visual Word Form Area): Located on the left side of the fusiform gyrus, this area is highly specialized for reading. It memorizes the visual shape of letters, character strings, and whole words, instantly recognizing written language configurations and passing them to language centers for rapid reading.
• Within-Category Visual Expertise: Beyond faces and text, the fusiform gyrus adapts to become a hub for deep visual expertise. If a person becomes an expert car mechanic, an avid birdwatcher, or a chess master, this gyrus structurally adapts to rapidly identify specific car models, bird species, or complex chess board configurations.

6. Parahippocampal Gyrus

Category: Spatial Navigation and Contextual Memory Encoding

• Environmental Scene Processing (Parahippocampal Place Area): The PPA is a highly specialized zone within this gyrus that lights up when you look at environmental scenes. It analyzes the structural layout of landscapes, city streets, building interiors, and rooms, mapping the physical geometry of your surroundings rather than individual objects.
• Memory Gateway to the Hippocampus: Wrapping directly around the hippocampus, this gyrus acts as the primary physical conduit for memory. It gathers sensory information from across the entire cerebral cortex, organizes it, and feeds it into the hippocampus so it can be packaged and stored as a long-term episodic memory.
• Spatial Navigation and Cognitive Mapping: The parahippocampal gyrus creates and maintains internal cognitive maps of the physical world. It keeps track of where you are located within a city or house, helping you navigate through space, remember routes, and orient yourself when you walk into an unfamiliar environment.
• Contextual Association: It links memories to specific contexts. When you encounter a specific smell, sight, or sound, the parahippocampal gyrus retrieves the holistic memory of the specific place, time, and situational backdrop where you originally experienced it.

1. Primary Auditory Cortex (A1)

Category: Sensory Transduction and Acoustic Analysis

• Acoustic Wave Decomposition: Located within Heschl's gyri, A1 is the ultimate terminal for raw auditory data. It receives neural impulses sent from the cochlea via the auditory nerve and thalamus, translating physical sound vibrations into conscious auditory perceptions.
• Spectral Frequency Mapping: A1 relies on a highly organized tonotopic grid where neighboring columns of neurons process neighboring sound frequencies. The anterior section maps low-frequency bass tones, while the posterior section decodes high-frequency treble tones.
• Intensity and Temporal Gating: Beyond pitch, A1 isolates the precise amplitude (volume) and microsecond timing of sound waves. It notes exactly when a sound starts, fluctuates, and stops, feeding this temporal structure to higher-level language and music centers.

2. Wernicke’s Area

Category: Semantic Decoding and Syntactic Assembly

• Linguistic Meaning Extraction: Situated in the posterior portion of the superior temporal gyrus (typically in the left hemisphere), Wernicke's area transforms acoustic phonemes into meaningful words. It acts as the brain's internal linguistic dictionary, matching auditory signals to stored conceptual definitions.
• Syntactic Structure Parsing: This area decodes the grammatical architecture of spoken and written sentences. It allows you to process the relationships between nouns, verbs, and prepositions so you can understand complex statements like "The dog chased the cat" versus "The cat chased the dog."
• Speech Generation Blueprinting: Before you speak, Wernicke's area constructs the conceptual
Looking at the larger picture can be used as a way of problem solving, which I think we discussed in the frontal lobe. In that frontal lobe, we discussed the fluidity of how something can be solved, right? That was the middle frontal gyrus, which constructs DAN and BAN, so we can work on that. Then memory encoding, again, we discuss this. Okay. Okay, the memory and recognition, the entorhinal hub, the spatial GPS.

The tissue around the rhinal sulcus is the primary interface between the cerebral cortex and the hippocampus. It contains the grid cells, which act like your brain's internal GPS, creating a cognitive map of its physical environment so you can navigate through space.

This is good. This can be a method for us, for the web code to navigate itself through the region of the internet. Once we open source it, we can give it the ability to understand where it is. We need to make it a mix so that it can be self-sufficient as long as it develops like a human. We cannot have a mother to cordle it, right?

Then we have the parietal cortex. The parietal region is responsible for identifying whether you have encountered an object or environment before. It gives you the distinct flash of recognition. It's memory, right? Again, wire it up to the memory one.

Then the collateral sulcus, the scene and the landscape decoder, the context and environment recognizer. The outer bank of the collateral sulcus houses the parahippocampal place area, the PPA. This specialises region activates intensely when you look at scenes, landscape, buildings, or a room. It does not look at an isolated object. It looks at the larger picture.

Looking at the larger picture can be used as a way of problem solving, which I think we discussed in the frontal lobe. In that frontal lobe, we discussed the fluidity of how something can be solved, right? That was the middle frontal gyrus, which constructs DAN and BAN, so we can work on that. Then memory encoding, again, we discuss this. Okay.
1. Heschl’s Gyri (Primary Auditory Cortex)

Category: Tonotopic Mapping and Spatial Acoustics

• Primary Auditory Processing: Slipped inside the lateral fissure on the upper surface of the temporal lobe, Heschl's gyri serve as the brain's gateway for sound. It is the first cortical structure to receive raw, unprocessed auditory signals traveling directly from the ears via the cochlea and the medial geniculate body of the thalamus.
• Tonotopic Organization: The tissue within Heschl's gyri is organized precisely by sound frequency, functioning like an internal piano keyboard. Neurons at one end are highly tuned to fire only to high-pitched sounds, while neurons at the opposite end fire strictly in response to low-pitched frequencies.
• Basic Feature Extraction: This gyrus breaks down sound waves into their foundational physical properties. It isolates the exact pitch (frequency), volume (amplitude), timbre, and baseline duration of a sound before passing these individual building blocks to adjacent areas for comprehension.
• Binaural Localization: By calculating the microsecond differences in when a sound reaches the left ear versus the right ear, Heschl's gyri calculate the directional origin of a sound, letting you map where a noise is coming from in your environment.

2. Superior Temporal Gyrus (STG)

Category: Secondary Auditory Processing and Language Comprehension

• Complex Acoustic Synthesis: Wrapping around Heschl's gyri, the STG takes the raw physical building blocks of sound and fuses them into recognizable, complex auditory patterns. It allows you to tell the difference between footsteps, a dog barking, a musical instrument, or a human voice.
• Phonological Processing: The STG acts as the primary deciphering engine for spoken language. It analyzes changing frequencies over time to identify phonemes—the specific, individual sounds that make up words (such as distinguishing the sound of a "B" from a "P").
• Wernicke's Area Integration: The posterior (rear) portion of the left STG houses Wernicke’s area, the critical neural hub for language comprehension. It decodes the syntactic structure of spoken sentences, allowing you to understand the actual meaning behind words spoken by others.
• Prosody and Emotional Mapping: While the left STG handles the linguistic meaning of words, the right STG evaluates the melody, rhythm, and intonation of speech (prosody). It decodes whether a speaker is angry, sarcastic, happy, or sad based entirely on their vocal tone.

3. Middle Temporal Gyrus (MTG)

Category: Lexical Storage and Multimodal Integration

• The Mental Lexicon: The MTG serves as the brain’s primary library for words. It stores your vocabulary, acting as a massive neural dictionary where word forms are linked directly to their definitions and grammatical profiles.
• Semantic Context Retrieval: During a conversation, the MTG rapidly retrieves words from memory so you can parse sentences in real time. It resolves ambiguity, ensuring that if

someone says the word "bank," your brain uses the surrounding conversation to instantly know if they mean a riverbank or a financial institution.
• Action and Tool Concepts: The MTG is heavily involved in processing verbs, action words, and tool concepts. When you look at an object like a hammer or read the word "run," neurons in the MTG fire to retrieve the conceptual memory of how that object is physically manipulated or how that action is performed.
• Multimodal Sensory Integration: Positioned symmetrically between the auditory cortex above and the visual cortex below, the MTG acts as a processing bridge. It takes visual inputs (like reading a word or seeing a picture) and auditory inputs (like hearing a word) and merges them into a single, unified conceptual idea.

4. Inferior Temporal Gyrus (ITG)

Category: Ventral Stream Visual Identification

• High-Level Visual Form Synthesis: The ITG occupies the lower lateral ridge of the temporal lobe and represents the visual processing peak of the ventral stream (the "what" pathway). It takes highly complex visual inputs—such as varying lines, shadows, colors, and textures—and synthesizes them into highly structured, recognizable shapes.
• Viewpoint Invariant Object Recognition: A core marvel of the ITG is its ability to recognize objects regardless of the angle from which they are viewed. If you look at a coffee mug from the top, the side, or upside down, the ITG ensures your brain still instantly identifies it as the exact same object.
• Perceptual Constancy: It maintains visual stability by tracking the structural reality of an item rather than its lighting environment. The ITG allows you to recognize an object whether it is cloaked in deep shadow, brightly illuminated by sunlight, or partially hidden behind something else.

5. Fusiform Gyrus (Lateral Occipitotemporal Gyrus)

Category: Expert Visual Recognition and Category Specialization

• Holistic Face Processing (Fusiform Face Area): A specialized region on the lateral side of the fusiform gyrus is explicitly dedicated to facial recognition. Instead of analyzing eyes, noses, and mouths as isolated parts, it processes faces holistically, mapping the exact spatial geometry of a face to tell you if you are looking at a stranger, a celebrity, or a family member.
• Visual Word Recognition (Visual Word Form Area): Located on the left side of the fusiform gyrus, this area is highly specialized for reading. It memorizes the visual shape of letters, character strings, and whole words, instantly recognizing written language configurations and passing them to language centers for rapid reading.
• Within-Category Visual Expertise: Beyond faces and text, the fusiform gyrus adapts to become a hub for deep visual expertise. If a person becomes an expert car mechanic, an avid birdwatcher, or a chess master, this gyrus structurally adapts to rapidly identify specific car models, bird species, or complex chess board configurations.

6. Parahippocampal Gyrus

Category: Spatial Navigation and Contextual Memory Encoding

• Environmental Scene Processing (Parahippocampal Place Area): The PPA is a highly specialized zone within this gyrus that lights up when you look at environmental scenes. It analyzes the structural layout of landscapes, city streets, building interiors, and rooms, mapping the physical geometry of your surroundings rather than individual objects.
• Memory Gateway to the Hippocampus: Wrapping directly around the hippocampus, this gyrus acts as the primary physical conduit for memory. It gathers sensory information from across the entire cerebral cortex, organizes it, and feeds it into the hippocampus so it can be packaged and stored as a long-term episodic memory.
• Spatial Navigation and Cognitive Mapping: The parahippocampal gyrus creates and maintains internal cognitive maps of the physical world. It keeps track of where you are located within a city or house, helping you navigate through space, remember routes, and orient yourself when you walk into an unfamiliar environment.
• Contextual Association: It links memories to specific contexts. When you encounter a specific smell, sight, or sound, the parahippocampal gyrus retrieves the holistic memory of the specific place, time, and situational backdrop where you originally experienced it.

1. Primary Auditory Cortex (A1)

Category: Sensory Transduction and Acoustic Analysis

• Acoustic Wave Decomposition: Located within Heschl's gyri, A1 is the ultimate terminal for raw auditory data. It receives neural impulses sent from the cochlea via the auditory nerve and thalamus, translating physical sound vibrations into conscious auditory perceptions.
• Spectral Frequency Mapping: A1 relies on a highly organized tonotopic grid where neighboring columns of neurons process neighboring sound frequencies. The anterior section maps low-frequency bass tones, while the posterior section decodes high-frequency treble tones.
• Intensity and Temporal Gating: Beyond pitch, A1 isolates the precise amplitude (volume) and microsecond timing of sound waves. It notes exactly when a sound starts, fluctuates, and stops, feeding this temporal structure to higher-level language and music centers.

2. Wernicke’s Area

Category: Semantic Decoding and Syntactic Assembly

• Linguistic Meaning Extraction: Situated in the posterior portion of the superior temporal gyrus (typically in the left hemisphere), Wernicke's area transforms acoustic phonemes into meaningful words. It acts as the brain's internal linguistic dictionary, matching auditory signals to stored conceptual definitions.
• Syntactic Structure Parsing: This area decodes the grammatical architecture of spoken and written sentences. It allows you to process the relationships between nouns, verbs, and

prepositions so you can understand complex statements like "The dog chased the cat" versus "The cat chased the dog."
• Speech Generation Blueprinting: Before you speak, Wernicke's area constructs the conceptual
I know many things are repeated here. Common sense in the baby's behavior will look a bit different from common sense from a normal human. Common sense for a baby's behavior would be the feeling of touch, right? Till now I kept discussing about cognition, but I have also written few notes about the frontal lobes work in like in motor coordination. Few of the things that I have jotted down are the pyramidal neurons, the bed cell, BETZ cells. These connect from precentral gyrus to the spinal cord. I cannot really give it a spinal cord because it really won't make sense, because we are making not a human, we are making an AI, so it does not make sense. But when we are giving it like the sense of touch, right? So for touch, it would really make sense if we give it some sort of a motor mechanism which allows it to understand. So for example, for an infant or for a newborn, for a fetus which is there in the mother's womb, the common sense is the fact that when the amniotic globule is felt on the lips of the fetus opens its mouth and takes it in, right? That is common sense. Another common sense is when the occasional kicking or sucking of thumb, that is also common sense, because sucking provides an understanding of how the baby will be sucking the milk out of the mother's breast, right? So that is also understood, if I'm not wrong. Another part of the common sense will be about the types of pars. So like the pars orbitalis, pars triangulis, all these, right? These are also some of the important things. I worked on like a research, I am working on it, about the timeline on how the neurological brain develops of a baby, along with the sensory details. I'll get to it, but till now think on how does this work out and how it's going to work out. The inferior frontal groove for cognitive flexibility, adaptation and rule switching with working memory for logic and words. It keeps track of goals, numbers and abstract concepts when solving problems. Superior frontal sulcus or the groove. Spatial working memory, top-down attention control, which facilitates for dorsal attention network, guides the FEF. Besides that, we have superior frontal gyrus. The superior frontal gyrus for anteromedial default mode network, the DMN, for self-awareness and introspection, navigating complex social dynamics. Then Broca's connection. It's back end of the inferior frontal sulcus runs into Broca's area. Correct word from memory during speech. This then the middle frontal gyrus, which allows for fluid reasoning, which allows for devising novel strategies and works with limbic system to understand the threat or regulate emotions, understand threat in the environment. Then dorsolateral SFG, superior frontal gyrus. It holds and manipulates complex concepts, numbers, rules, suppress distraction for focus. Then inferior frontal gyrus, the forms the core of Broca's area, transform abstract language and vocabulary into complex sequences. Then dorsolateral prefrontal cortex, manipulating working memory, strategic under pressure. Then premotor cortex, that calculating spatial orientation of objects. Then the work of gyrus rectus, which coordinates with the orbitofrontal cortex, which works with the amygdala for its calculation, and anterior cingulate cortex for error related negativity impulse and choose between conflicting options. A few approaches I found here through which we may be able to manipulate the brain. Again, there are a few other parts which I'll list them here only.
Brain
1. connect the three brains
for sharpening of ideas, check-in results, Turn each big idea into a test that could fail, Let each life stage switch ideas on when the biology does
we will discuss that.

_____

Frontal Lobe:
Central Sulcus (Groove)
Physical boundary between the thinking/moving (frontal) & feeling (parietal). -- deeper central sulcus can contribute to extraordinary dexterity, faster motor learning, and a hyper-acute sense of touch. Which can help with the overall sensory understanding, which we discussed.

Precentral Sulcus
It serves as the physical border that separates your primary movement center from your high-level planning and decision-making centers. For premotor cortex & supplementary motor area, I was thinking of enhancing it to push for a more possible complex chain of actions. when we will provide the baby with a library, and understanding of physical bounds. The Broca's area.. we will try to make it stronger so the interaction and overall clarity of ideas of novelty remains, pair this up with frontal eye speed -- possible hypothesis : "If Broca's area and the Frontal Eye Fields (FEF) were structurally deeper and had hyper-direct, heavily myelinated connections leading straight into the central sulcus". Extending to this, few other ideas I had was the fact that we see again Broca's connection is again connecting to Broca's area, which collects fluid from memory, and you know, overall the Broca's area, it in general works in correcting speech, working with the more articulate sentences, like I'm speaking right now, right? So I was thinking that we can pair, like, what if the brain had deeper grooves which connected superior frontal sulcus, the spatial working memory, and top-down attention control which guides the FEF, inferior frontal groove, the spatial working memory paired with the cognitive flexibility allows for a sharper understanding of the environment, which will also give the working memory for logic and words, keeping in track of goals, numbers, abstract concepts, and also guiding the FEF, which will be extremely extremely efficient for vision, optical understanding. Then close, not closer to this, but a bit different was my thought for default mode network, which we can use for a path towards common sense with how the newborn will understand, right? Then with the other idea I had was superior frontal sulcus, inferior frontal groove, the dorsolateral SFG, and connecting it to dorsolateral prefrontal cortex, which will work extremely well as we can connect these deeply. If I'm not wrong, these are not deeply connected, but rather a bit scattered. So connecting them deeply will. The inferior frontal groove for cognitive flexibility, adaptation and rule switching with working memory for logic and words. It keeps track of goals, numbers and abstract concepts when solving problems. Superior frontal sulcus or the groove. Spatial working memory, top-down attention control, which facilitates for dorsal attention network, guides the FEF. Besides that, we have superior frontal gyrus. The superior frontal gyrus for anteromedial default mode network, the DMN, for self-awareness and introspection, navigating complex social dynamics. Then Broca's connection. It's back end of the inferior frontal sulcus runs into Broca's area. Correct word from memory during speech. This then the middle frontal gyrus, which allows for fluid reasoning, which allows for devising novel strategies and works with limbic system to understand the threat or regulate emotions, understand threat in the environment. Then dorsolateral SFG, superior frontal gyrus. It holds and manipulates complex concepts, numbers, rules, suppress distraction for focus. Then inferior frontal gyrus, the forms the core of Broca's area, transform abstract language and vocabulary into complex sequences. Then dorsolateral prefrontal cortex, manipulating working memory, strategic under pressure. Then premotor cortex, that calculating spatial orientation of objects. Then the work of gyrus rectus, which coordinates with the orbitofrontal cortex, which works with

the amygdala for its calculation, and anterior cingulate cortex for error related negativity impulse and choose between conflicting options. A few approaches I found here through which we may be able to manipulate the brain. Again, there are a few other parts which I'll list them here only. "The Biological Blueprint: How the Mind Creates Reality

This is the comprehensive, interconnected summary of the journey we mapped out—from a single cell in the dark of the womb to a high-level adult leader navigating the world.

Phase 1: The Blueprint and The Separation (The Womb)

The human brain begins as a flat disk of cells that curls into a tube (neurulation). Deep inside this tube, stem cells enter a hyper-drive state, generating 250,000 neurons every single minute until reaching roughly 86 billion cells. These newborn neurons literally crawl along glial ladders (neural migration) to build the layers of the brain from the inside out.
To prevent catastrophic sensory overload, nature activates the senses in a strict, protective sequence:
1. Touch (The Somatic Baseline): Activated at just 8 weeks, the embryo's first action is a primitive flinch. This tiny reflex loop is the literal origin point of self-awareness. It establishes the binary feedback loop ("There is a Me, and there is a Not-Me") that maps the physical boundaries of the body onto the parietal lobe.
2. Chemical Senses: Taste and smell follow, introducing the chemistry of the outside environment.
3. Hearing: The baby hears the steady, double-thump rhythm of the mother’s heartbeat. This massive, repetitive data stream becomes so deeply wired that after birth, skin-to-skin contact instantly calms the reactive limbic system through structural familiarity.
4. Vision: Kept completely dark until birth, protecting the massive visual cortex from premature data flooding.

Phase 2: The Physical Construction Site (Infancy to Youth)

A baby is born with a massive pile of unlinked "cellular bricks." The architecture of the brain literally builds itself through real-world experiences.
• The Sucking Reflex: Months of practicing the touch-and-open sequence with amniotic fluid in the womb hardwires a sensory-motor highway between the parietal lobe (detecting the touch) and the prefrontal cortex (coordinating the action). At birth, this ingrained loop allows the infant to nurse automatically.
• Serve and Return: Back-and-forth interactions between a baby and a caregiver act as the primary construction workers, physically sprouting branches (axons and dendrites) to forge millions of new connections (synapses) every second.
• Pouring the Concrete: Through neuroplasticity, repetition cements these paths. Heavily used neural tracks are wrapped in a protective, fatty insulation called myelin, turning slow dirt roads into high-speed fiber-optic cables (myelination).
• The Sculpting (Pruning): By age 3, the brain has twice as many connections as an adult. Neuroplasticity triggers a massive clean-up (synaptic pruning), physically dismantling unused connections to leave a streamlined brain perfectly adapted to its specific environment.

Phase 3: The Birth of Logic and Neural Topography

A child’s brain does not start with abstract logic; it is a pattern-seeking machine.
• Action and Consequence: Through repetition (like dropping a spoon 100 times), the brain links sensory data to outcomes, building physical "If/Then" rules.
• Retinotopy (Geometric Mapping): The brain physically arranges its cells in geometric layout patterns that mimic external inputs. When the retina sees a circle, the neurons in the visual cortex fire in the exact physical shape of a circle. The brain literally arranges its own internal shape to map the external world.
• Executive Control: As a child grows, the Frontoparietal Network (FPN)—the physical highway linking sensory processing (parietal) to executive planning (prefrontal)—matures. This allows the prefrontal cortex to grow inhibitory pathways that can suppress the raw emotional alarms of the limbic system, transitioning the child from purely emotional reactions to abstract, rational logic.

Phase 4: The Lifelong Shifting of the Tides (Adulthood to Aging)

The brain never stops physically changing its structure, a lifelong ability driven by neuroplasticity.

[Ages 10-14]	[Ages 30-40]	[Ages 38-45]	[Early 70s]	
MAX PHYSICAL SIZE	MAX WIRING EFFICIENCY	MAX SYSTEM INTEGRATION	ACCELERATED DECAY	> Time
Total brain volume peaks early.	White matter insulation (myelin) reaches its peak.	Limbic & Prefrontal networks perfectly sync.	Brain volume shrinkage rate more than doubles.	

• The Muscle Effect: Regional areas can physically grow or shrink based on use (like the expanded spatial hippocampus found in London taxi drivers).
• The Horsepower Peak (Early 20s): Raw mental processing speed peaks at 18–20, and working memory (holding short-term information) peaks around 25.
• The Leadership Peak (40s to 60s): While raw processing speed slows down, the brain reaches maximum pathway integration, structural stability, and emotional regulation around age 32. The deeply mature prefrontal-to-limbic connections allow mid-life brains to manage intense stress, resist cognitive biases, and deploy a lifetime database of crystallized intelligence (wisdom). This is why society deeply prefers leaders and CEOs in the 40–60 age bracket.
• The Structural Shift (Early 70s): A scientifically documented turning point occurs where natural brain shrinkage dramatically accelerates (doubling to over 0.4%+ per year). Myelin insulation begins to degrade, causing long-distance communication networks like the FPN to slow down. However, because neuroplasticity never truly turns off, rigorous cognitive challenges and physical health can actively cushion this decline.

The Grand Conclusion: "Eating" the World

Ultimately, we realized that the line between abstract thought and physical biology doesn't exist. Human language perfectly mirrors our neurology: when we "chew on food," we break down external items to physically build our bodies; when we "chew on an idea," the FPN

breaks down external data and uses neuroplasticity to physically weave that concept into the wiring of our brain cells.

Whether it is a fetus touching its lips to map reality, a toddler categorizing a cat versus a dog, a CEO managing a global crisis, or a future human using spatial computing to manipulate 3D data—learning is simply the physical brain "eating" its environment to sculpt its own shape.

We have woven together an incredible biological narrative. As you look over this complete map, which specific chapter feels like the most fascinating spark for your own thinking—the sensory training of the womb, the mechanics of how logic forms, or the philosophy of how the brain literally "eats" information?"”