

# A multivariate skew-normal finite mixture model for analysis of infant development trajectories

**Carter Allen**

Department of Public Health Sciences, Medical University of South Carolina, Charleston, SC, U.S.A

*email:* allecart@musc.edu

**and**

**Brian Neelon, PhD**

Department of Public Health Sciences, Medical University of South Carolina, Charleston, SC, U.S.A

**and**

**Sara Benjamin-Neelon, PhD, MPH, RD**

Department of Health, Behavior and Society, Johns Hopkins University, Baltimore, MD, U.S.S

**SUMMARY:** In studies of infant motor development, a crucial research goal is to identify latent classes of infants that experience delayed development, as this is a known risk factor for adverse outcomes later in life. However, there are a number of statistical challenges in modeling infant development: the data are typically skewed, exhibit intermittent missingness, and are highly correlated across the repeated measurements collected during infancy. Using data from the Nurture study, a cohort of over 600 mother-infant pairs followed from pregnancy to 12 months postpartum, we develop a flexible Bayesian latent class model for the analysis infant motor development. Our model has a number of attractive features. First, we adopt the multivariate skew normal distribution with class-specific parameters that accommodate the inherent correlation and skewness in the data. Second, we model the class membership probabilities using a novel Plya-Gamma data-augmentation scheme, thereby improving predictions of the class membership allocations. Lastly, we impute missing responses under missing at random assumption by drawing from appropriate conditional skew normal distributions. Bayesian inference is achieved through straightforward Gibbs sampling, and can be carried out in available software such as R. Through simulation studies, we show that the proposed model yields improved inferences over models that ignore skewness. In addition, our imputation method yields improvements compared to conventional missing data methods, including multiple imputation and complete or available case analysis. When applied to Nurture data, we identified two distinct development classes: one characterized by delayed U-shaped development and a higher percentage of male infants and another characterized by more steady development and a

December 2008

lower percentage of males. The classes also differed in terms of key demographic variables, such as infant race and maternal pre-pregnancy body mass index. These findings can aid investigators in targeting interventions during this critical early-life developmental window.

KEY WORDS: A key word; But another key word; Still another key word; Yet another key word.

## CONTENTS

- 1 Introduction
- 2 Nurture Study
  - 2.1 Baseline Demographics and Description of Variables
  - 2.2 Statistical Challenges
    - 2.2.1 Skewness of Bayley score residuals
    - 2.2.2 Attrition and Intermittent Missingness
- 3 Model
  - 3.1 Multivariate Skew Normal Regression
  - 3.2 Multinomial Regression on Class Probabilities
  - 3.3 Conditional MVSN Imputation
  - 3.4 Bayesian Inference
    - 3.4.1 MCMC Algorithm
    - 3.4.2 Assessment of MCMC Convergence
    - 3.4.3 Label Switching
    - 3.4.4 Prior Choice
- 4 Simulation Studies
  - 4.1 Simulation to Compare to Multivariate Normal
  - 4.2 Simulation to Compare Imputation Methods
  - 4.3 Simulation to Assess Sensitivity to Misspecified H
- 5 Application
- 6 Discussion
- 7 Appendix
  - 7.1 Glossary of Notation
  - 7.2 Derivation of Full Conditional Distributions

7.2.1 Multivariate Skew-Normal Regression

7.2.2 Multinomial Logit Regression

7.2.3 Multivariate Normal Conditional Imputation

References

## 1. Introduction

## 2. Nurture Study

### 2.1 Baseline Demographics and Description of Variables

### 2.2 Statistical Challenges

#### 2.2.1 Skewness of Bayley score residuals.

#### 2.2.2 Attrition and Intermittent Missingness.

## 3. Model

### 3.1 Multivariate Skew Normal Regression

We model the effect of covariates on longitudinal development outcomes through the use of a MSN regression model. The MSN distribution can be represented as the superposition of a MN random variable with a latent truncated normal random effect. Let  $\mathbf{Y}_{n \times k}$  be the observation matrix such that  $Y_{ij}$  is the observation for subject  $i$  at timepoint  $j$ .

$$\mathbf{Y}_{n \times k} = \mathbf{X}_{n \times p} \boldsymbol{\beta}_{p \times k} + t_{n \times 1} \boldsymbol{\psi}_{1 \times k} + \boldsymbol{\epsilon}_{n \times k}$$

where  $X_i$  is the  $1 \times p$  vector of covariate values for subject  $i$ ,  $\beta_j$  is the  $i \times k$  vector of fixed effects coefficients for timepoint  $j$ ,  $t_i \stackrel{iid}{\sim} N_{[0, \infty)}(0, 1)$  is a truncated normal random effect,  $\psi$  is the vector containing skewness parameters for each timepoint, and  $\boldsymbol{\epsilon}_i \sim N_k(0, \boldsymbol{\Sigma}_{k \times k})$  is the correlated error term.

### 3.2 Multinomial Regression on Class Probabilities

A primary concern of our model is with identification of latent infant development clusters. We accomplish this via multinomial logit regression model on cluster membership, which utilizes Pólya-Gamma data-augmentation to allow for updating of all parameters using Gibbs sampling. The multinomial logit model is as follows for  $l = 1, \dots, h$ .

$$P(Z_i = l | w_i) = \pi_{il} = \frac{e^{w_i^T \delta_l}}{\sum_{r=1}^h e^{w_i^T \delta_r}}$$

where  $w_i$  is the vector of class probability covariates for subject  $i$ ,  $\delta_l$  contains the multinomial regression parameters for class  $l$ , and  $h$  is the number of putative clusters specified *a priori*.

During our MCMC estimation procedure, the class labels  $z_i$  are updated from their multinomial full conditional distribution and used in the remaining MCMC steps as class assignments.

### 3.3 Conditional MVSN Imputation

We allow for missingness of outcomes in the MSN mixture model by imputing missing values from their conditional multivariate normal distributions. We note that

$$Y_i | X_i, t_i, \beta, \psi \sim N_k(X_i \beta + t_i \psi, \Sigma)$$

This allows us to appeal to standard conditional forms of the multivariate normal distribution. Let  $Y_i = [Y_{i_q \times 1}^{miss} | Y_{i_{k-q} \times 1}^{obs}]^T$ . We have

$$Y_i^{miss} | Y_i^{obs}, X_i, t_i, \beta, \psi \sim N(\mu^{miss}, \Sigma^{miss})$$

where  $\mu^{miss}$  and  $\Sigma^{miss}$  take standard forms. Each missing outcome is imputed "online", i.e. once per MCMC iteration. This provides more opportunities to explore the parameter space than multiple imputation and avoids multiplicative run-time scaling in  $m$ , the number of imputations.

### 3.4 Bayesian Inference

#### 3.4.1 MCMC Algorithm.

### 3.4.2 *Assessment of MCMC Convergence.*

### 3.4.3 *Label Switching.*

### 3.4.4 *Prior Choice.*

## 4. Simulation Studies

### 4.1 *Simulation to Compare to Multivariate Normal*

### 4.2 *Simulation to Compare Imputation Methods*

### 4.3 *Simulation to Assess Sensitivity to Misspecified $H$*

## 5. Application

## 6. Discussion

## 7. Appendix

Put your final comments here.

### ACKNOWLEDGEMENTS

### SUPPLEMENTARY MATERIALS

#### 7.1 *Glossary of Notation*

- **Y**: A  $n \times p$  matrix containing all multivariate skew-normal regression outcomes such that  $y_{ij}$  is the  $j^{th}$  outcome observed for subject  $i$ , where  $i = 1, \dots, n$  and  $j = 1, \dots, p$ .
- **X**: A  $n \times m$  matrix containing all multivariate skew-normal regression covariates such that  $x_{ij}$  is the  $j^{th}$  covariate value for subject  $i$ .
- **B**: A  $m \times p$  matrix containing all multivariate skew-normal regression coefficients such that  $\mathbf{B} = [\boldsymbol{\beta}_1, \dots, \boldsymbol{\beta}_p]$ , where  $\beta_{ij}$  is interpreted as the effect of covariate  $i$  on outcome  $j$  for  $i = 1, \dots, m$  and  $j = 1, \dots, p$ .

- **E**: A  $n \times p$  matrix of error terms in the multivariate skew-normal regression model component. **E** is made up of row vectors  $\boldsymbol{\epsilon}_i = (\epsilon_{i1}, \dots, \epsilon_{ip})$ , where  $\boldsymbol{\epsilon}_i \stackrel{iid}{\sim} N_p(0, \boldsymbol{\Sigma})$  for  $i = 1, \dots, n$ .
- **$\boldsymbol{\Sigma}$** : A  $p \times p$  covariance matrix that defines the correlation between the  $p$  multivariate normal outcomes.
- **$\boldsymbol{\Omega}$** : A  $p \times p$  covariance scale matrix that defines the correlation between the  $p$  multivariate skew-normal outcomes.
- **$\boldsymbol{\psi}$** : A  $p \times 1$  vector containing the skewness parameter for each outcome.
- **$\boldsymbol{\alpha}$** : A  $p \times 1$  vector containing the skewness parameter for each outcome.
- **t**: A  $n \times 1$  vector of truncated normal random effects used in the stochastic representation of the multivariate skew-normal distribution. For  $i = 1, \dots, n$ ,  $t_i \stackrel{iid}{\sim} T[0, \infty)(0, 1)$
- **X\***: A  $n \times (m + 1)$  matrix constructed by column binding **t** to **X**
- **B\***: A  $(m + 1) \times p$  matrix constructed by row binding  $\boldsymbol{\psi}^T$  to **B**.

## 7.2 Derivation of Full Conditional Distributions

**7.2.1 Multivariate Skew-Normal Regression.** Without loss of generality, we derive the full conditional distributions for the multivariate skew-normal regression model component under the assumption that all observations belong to a single cluster. To make the extension to the case where more than one cluster is specified, simply apply these distributional forms to cluster specific parameters and data.

### 7.2.2 Multinomial Logit Regression.

### 7.2.3 Multivariate Normal Conditional Imputation.

## REFERENCES

- Cox, D. R. (1972). Regression models and life tables (with discussion). *Journal of the Royal Statistical Society, Series B* **34**, 187–200.



Hastie, T., Tibshirani, R., and Friedman, J. (2001). *The Elements of Statistical Learning: Data Mining, Inference, and Prediction*. New York: Springer.

*Received October 2007. Revised February 2008. Accepted March 2008.*