

3 Spectral Theory V: Diagonalization

Prof.Dr.P.M.Bajracharya

School of Mathematical Sciences
T.U., Kirtipur

April 25, 2024

Summary

① Diagonalization

Diagonalization

Similar matrices

A square matrix B is **similar** to a matrix A if there exists an invertible matrix P such that $P^{-1}AP = B$.

We have proved in “3_Spectral theory II”:

Theorem

Let A and B be similar matrices. Then the eigenvalues of these matrices are identical.

Diagonalizable matrix

An $n \times n$ matrix A is said to **diagonalizable** if it is similar to a diagonal matrix D .

Theorem

An $n \times n$ matrix A is **diagonalizable** iff it has n linearly independent eigenvectors.

Example

Let $A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. Then A cannot be diagonalized.

Example

Let $A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. Then A cannot be diagonalized.

Solution. A is diagonalizable implies A is similar to a diagonal matrix D with diagonal entries $\{d_1, d_2\} = \{0, 0\}$.

Example

Let $A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. Then A cannot be diagonalized.

Solution. A is diagonalizable implies A is similar to a diagonal matrix D with diagonal entries $\{d_1, d_2\} = \{0, 0\}$. Hence

$$D = 0 \Rightarrow A = PDP^{-1} = 0.$$

This is a contradiction. □

Example

Let $A = \begin{pmatrix} 2 & 1 & 1 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{pmatrix}$. Then A cannot be diagonalized.

Example

Let $A = \begin{pmatrix} 2 & 1 & 1 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{pmatrix}$. Then A cannot be diagonalized.

Solution. A is diagonalizable implies A is similar to a diagonal matrix D with diagonal entries $\{d_1, d_2, d_3\} = \{2, 2, 2\}$.

Example

Let $A = \begin{pmatrix} 2 & 1 & 1 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{pmatrix}$. Then A cannot be diagonalized.

Solution. A is diagonalizable implies A is similar to a diagonal matrix D with diagonal entries $\{d_1, d_2, d_3\} = \{2, 2, 2\}$. Hence

$$D = 2I \Rightarrow A = PDP^{-1} = 2I.$$

This is a contradiction. □

Theorem

Let A be an $n \times n$ matrix. If P is an invertible matrix such that

$$PDP^{-1} = \text{diag}(d_1, \dots, d_n),$$

then for $1 \leq i \leq n$, the i -th column of P is an eigenvector of A corresponding to d_i .

Proof. Let $P = [u_1, \dots, u_n]$.

Proof. Let $P = [u_1, \dots, u_n]$. Then

$$P^{-1}AP = D \Rightarrow AP = PD.$$

Proof. Let $P = [u_1, \dots, u_n]$. Then

$$P^{-1}AP = D \Rightarrow AP = PD.$$

Thus,

$$[Au_1, \dots, Au_n] = A[u_1, \dots, u_n]$$

Proof. Let $P = [u_1, \dots, u_n]$. Then

$$P^{-1}AP = D \Rightarrow AP = PD.$$

Thus,

$$\begin{aligned}[Au_1, \dots, Au_n] &= A[u_1, \dots, u_n] \\ &= AP = PD\end{aligned}$$

Proof. Let $P = [u_1, \dots, u_n]$. Then

$$P^{-1}AP = D \Rightarrow AP = PD.$$

Thus,

$$\begin{aligned} [Au_1, \dots, Au_n] &= A[u_1, \dots, u_n] \\ &= AP = PD \\ &= P \operatorname{diag}(d_1, \dots, d_n) \end{aligned}$$

Proof. Let $P = [u_1, \dots, u_n]$. Then

$$P^{-1}AP = D \Rightarrow AP = PD.$$

Thus,

$$\begin{aligned} [Au_1, \dots, Au_n] &= A[u_1, \dots, u_n] \\ &= AP = PD \\ &= P \operatorname{diag}(d_1, \dots, d_n) \\ &= [d_1u_1, \dots, d_nu_n]. \end{aligned}$$

Proof. Let $P = [u_1, \dots, u_n]$. Then

$$P^{-1}AP = D \Rightarrow AP = PD.$$

Thus,

$$\begin{aligned} [Au_1, \dots, Au_n] &= A[u_1, \dots, u_n] \\ &= AP = PD \\ &= P \operatorname{diag}(d_1, \dots, d_n) \\ &= [d_1u_1, \dots, d_nu_n]. \end{aligned}$$

Or equivalently,

$$Au_i = d_iu_i, \text{ for } 1 \leq i \leq n.$$

Proof. Let $P = [u_1, \dots, u_n]$. Then

$$P^{-1}AP = D \Rightarrow AP = PD.$$

Thus,

$$\begin{aligned} [Au_1, \dots, Au_n] &= A[u_1, \dots, u_n] \\ &= AP = PD \\ &= P \operatorname{diag}(d_1, \dots, d_n) \\ &= [d_1u_1, \dots, d_nu_n]. \end{aligned}$$

Or equivalently,

$$Au_i = d_iu_i, \text{ for } 1 \leq i \leq n.$$

As P is invertible, u_1, \dots, u_n are linearly independent.

Proof. Let $P = [u_1, \dots, u_n]$. Then

$$P^{-1}AP = D \Rightarrow AP = PD.$$

Thus,

$$\begin{aligned} [Au_1, \dots, Au_n] &= A[u_1, \dots, u_n] \\ &= AP = PD \\ &= P \operatorname{diag}(d_1, \dots, d_n) \\ &= [d_1u_1, \dots, d_nu_n]. \end{aligned}$$

Or equivalently,

$$Au_i = d_iu_i, \text{ for } 1 \leq i \leq n.$$

As P is invertible, u_1, \dots, u_n are linearly independent. Hence, (d_i, u_i) , for $1 \leq i \leq n$, are eigen-pairs of A . This proves the theorem. \square

From the proof of the above theorem we obtain the following theorem

Theorem

Let A be an $n \times n$ matrix. If A is diagonalizable, then A has n linearly independent eigenvectors.

Its converse is also true.

Theorem

Let A be an $n \times n$ matrix. If A has n linearly independent eigenvectors, then A is diagonalizable.

Proof. Let $\{u_1, \dots, u_n\}$ be n linearly independent eigenvectors of A corresponding to eigenvalues $\lambda_1, \dots, \lambda_n$.

Proof. Let $\{u_1, \dots, u_n\}$ be n linearly independent eigenvectors of A corresponding to eigenvalues $\lambda_1, \dots, \lambda_n$. Then $V = [u_1, \dots, u_n]$ is non-singular and

Proof. Let $\{u_1, \dots, u_n\}$ be n linearly independent eigenvectors of A corresponding to eigenvalues $\lambda_1, \dots, \lambda_n$. Then $V = [u_1, \dots, u_n]$ is non-singular and

$$AV = [Au_1, \dots, Au_n]$$

Proof. Let $\{u_1, \dots, u_n\}$ be n linearly independent eigenvectors of A corresponding to eigenvalues $\lambda_1, \dots, \lambda_n$. Then $V = [u_1, \dots, u_n]$ is non-singular and

$$\begin{aligned} AV &= [Au_1, \dots, Au_n] \\ &= [\lambda_1 u_1, \dots, \lambda_n u_n] \end{aligned}$$

Proof. Let $\{u_1, \dots, u_n\}$ be n linearly independent eigenvectors of A corresponding to eigenvalues $\lambda_1, \dots, \lambda_n$. Then $V = [u_1, \dots, u_n]$ is non-singular and

$$\begin{aligned} AV &= [Au_1, \dots, Au_n] \\ &= [\lambda_1 u_1, \dots, \lambda_n u_n] \\ &= V\Lambda, \end{aligned}$$

where $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$.

Proof. Let $\{u_1, \dots, u_n\}$ be n linearly independent eigenvectors of A corresponding to eigenvalues $\lambda_1, \dots, \lambda_n$. Then $V = [u_1, \dots, u_n]$ is non-singular and

$$\begin{aligned} AV &= [Au_1, \dots, Au_n] \\ &= [\lambda_1 u_1, \dots, \lambda_n u_n] \\ &= V\Lambda, \end{aligned}$$

where $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$. Therefore,

$$V^{-1}AV = \Lambda.$$

This implies that A is diagonalizable. □

Example

Show that the following matrix is diagonalizable:

$$A = \begin{bmatrix} 1 & 0 & -1 \\ 1 & 2 & 1 \\ 2 & 2 & 3 \end{bmatrix}.$$

Solution: First, we shall find the eigenvalues of A .

Solution: First, we shall find the eigenvalues of A . The characteristic equation of A is given by:

$$|A - \lambda I| = \begin{vmatrix} 1 - \lambda & 0 & -1 \\ 1 & 2 - \lambda & 1 \\ 2 & 2 & 3 - \lambda \end{vmatrix} = 0$$

Solution: First, we shall find the eigenvalues of A . The characteristic equation of A is given by:

$$|A - \lambda I| = \begin{vmatrix} 1 - \lambda & 0 & -1 \\ 1 & 2 - \lambda & 1 \\ 2 & 2 & 3 - \lambda \end{vmatrix} = 0$$

$$\Rightarrow (1 - \lambda)(2 - \lambda)(3 - \lambda) = 0$$

Solution: First, we shall find the eigenvalues of A . The characteristic equation of A is given by:

$$|A - \lambda I| = \begin{vmatrix} 1 - \lambda & 0 & -1 \\ 1 & 2 - \lambda & 1 \\ 2 & 2 & 3 - \lambda \end{vmatrix} = 0$$

$$\Rightarrow (1 - \lambda)(2 - \lambda)(3 - \lambda) = 0$$

$$\Rightarrow \lambda = 3, 2, 1.$$

$$\underline{\underline{\lambda_1 = 3}}$$

The eigenvector corresponding to $\lambda_1 = 3$ is the non-zero solution of the following matrix equation:

$$(A - 3I)X = 0$$

$$\underline{\underline{\lambda_1 = 3}}$$

The eigenvector corresponding to $\lambda_1 = 3$ is the non-zero solution of the following matrix equation:

$$(A - 3I)X = 0$$

$$\Rightarrow \begin{bmatrix} 0 & 0 & -1 \\ 1 & 1 & 1 \\ 2 & 2 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\underline{\underline{\lambda_1 = 3}}$$

The eigenvector corresponding to $\lambda_1 = 3$ is the non-zero solution of the following matrix equation:

$$\begin{aligned} (A - 3I)X &= 0 \\ \Rightarrow \begin{bmatrix} 0 & 0 & -1 \\ 1 & 1 & 1 \\ 2 & 2 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} &= \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \\ \Rightarrow \begin{bmatrix} -z \\ x + y + z \\ 2x + 2y + 2z \end{bmatrix} &= \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \end{aligned}$$

This gives

$$\begin{cases} -z = 0 \\ x + y = 0 \end{cases}$$

This gives

$$\begin{cases} -z = 0 \\ x + y = 0 \end{cases}$$

If we take $x = 1 \Rightarrow y = -1$.

This gives

$$\begin{cases} -z = 0 \\ x + y = 0 \end{cases}$$

If we take $x = 1 \Rightarrow y = -1$.

Hence, the corresponding eigen-vector

$$u_1 = \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$$

Similarly, the eigenvector corresponding to $\lambda_2 = 2$ is given by:

$$\begin{bmatrix} -1 & 0 & -1 \\ 1 & 0 & 1 \\ 2 & 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} -x - z \\ x + z \\ 2x + 2y + z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Similarly, the eigenvector corresponding to $\lambda_2 = 2$ is given by:

$$\begin{aligned} & \begin{bmatrix} -1 & 0 & -1 \\ 1 & 0 & 1 \\ 2 & 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \\ \Rightarrow & \begin{bmatrix} -x - z \\ x + z \\ 2x + 2y + z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \\ \Rightarrow & \begin{cases} x + z = 0 \\ 2x + 2y + z = 0 \end{cases}. \end{aligned}$$

Hence, the corresponding eigen-vector

$$u_1 = \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix}$$

Hence, the corresponding eigen-vector

$$u_1 = \begin{bmatrix} 2 \\ -1 \\ -2 \end{bmatrix}$$

Put

$$P = \begin{bmatrix} 1 & -2 & 1 \\ -1 & 1 & -1 \\ 0 & 2 & -2 \end{bmatrix}$$

Now,

$$P^{-1}AP = \frac{1}{2} \begin{bmatrix} 0 & -2 & 1 \\ -2 & -2 & 0 \\ -2 & -2 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 \\ 1 & 2 & 1 \\ 2 & 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ -1 & 1 & -1 \\ 0 & 2 & -2 \end{bmatrix}.$$

Now,

$$P^{-1}AP = \frac{1}{2} \begin{bmatrix} 0 & -2 & 1 \\ -2 & -2 & 0 \\ -2 & -2 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 \\ 1 & 2 & 1 \\ 2 & 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ -1 & 1 & -1 \\ 0 & 2 & -2 \end{bmatrix}.$$

Then we see that

$$P^{-1}AP = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}.$$

Now,

$$P^{-1}AP = \frac{1}{2} \begin{bmatrix} 0 & -2 & 1 \\ -2 & -2 & 0 \\ -2 & -2 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -1 \\ 1 & 2 & 1 \\ 2 & 2 & 3 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ -1 & 1 & -1 \\ 0 & 2 & -2 \end{bmatrix}.$$

Then we see that

$$P^{-1}AP = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}.$$

Thus, A is diagonalizable.

Thus, from our work above, if there is a basis of \mathbb{R}^n consisting of eigenvectors of A , then A is similar to a diagonal matrix D and so A is diagonalizable.

Thus, from our work above, if there is a basis of \mathbb{R}^n consisting of eigenvectors of A , then A is similar to a diagonal matrix D and so A is diagonalizable.

On the other hand, if at least one of the eigenvalues of A is deficient, then A will not have linearly independent eigenvectors.

Thus, from our work above, if there is a basis of \mathbb{R}^n consisting of eigenvectors of A , then A is similar to a diagonal matrix D and so A is diagonalizable.

On the other hand, if at least one of the eigenvalues of A is deficient, then A will not have linearly independent eigenvectors. Hence we will not be able to construct an invertible matrix P whose columns are eigenvectors of A . In this case, we say that A is not diagonalizable.